CONFERENCE REPORTS...

Ulan Bator Welcomes Phosphorites Experts

IGCP Project 156: Third International Seminar

Group photograph of participants in the 1980 international meeting of IGCP Project 156 held in Ulan Bator, capital of the Mongolian Peoples' Republic. From left to right, sitting: W. Burnett, D. Tamdin, R. Barzbold, O. Tomoor-Togoo, P. Southgate, G. Ratnikova, P.F. Howard, D. Banerjee, A. Ziegler; standing: P. Cook, N. Yudin, R. Choudhury, E. Eganov, R. Rogers, I. Lozovaya, A. Yanshin, (the next three persons not identified), I. Shergold, R. Sheldon, A. Pant, Y. Zanin, S. Riggs, V. Pokryshkin, P. Pritchard, B. Barjin, Y. Popov, B. Luvsandansan, K. Muller, D. Chinbat, M. Batchulun, B. Gurelhajav. (Photo courtesy F. Zambrano).

The Khubsugul Basin phosphorites in northern Mongolia are important to an understanding of Cambrian world phosphogenic events, and hence their study is of significance to the geoscientific community as a whole. This was one of the main conclusions drawn by participants in the Third International Field Workshop and Seminar held from June 17 - 28, 1980 in the Mongolian Peoples' Republic (M.P.R.) and sponsored by IGCP Project 156 ("Phosphorites") in collaboration with the M.P.R. Academy of Sciences, which hosted the meeting.

In endorsing the view that more detailed geological studies of the Khubsugul deposits are desirable, participants made the following specific recommendations:

- While the Khubsugul phosphorites have been considered to be of Vendian to Cambrian age, the group considers an Early Cambrian age to be more likely, and concludes that further research is necessary, particularly since deposits of this age are increasing in importance as sources of phosphate around the world;
- A clearer understanding of the detailed chronology and paleogeography would facilitate further exploration for phosphorites in the region;
- The environment of deposition appears to fluctuate within an epicontinental shelf environment. Consequently, there are likely to be facies changes (some may be rapid) which must be delineated through detailed sedimentary facies analysis and development of depositional models;
- Geological research should continue on the phosphogenic sequences throughout the region using a wide range of techniques such as geochemistry (organic, inorganic, isotopic,

Figure 1. Map showing the location of the 1980 field workshop held by IGCP Project 156.

CONFERENCE REPORTS (continued)

and the redox state of selected elements), lithogenesis, etching for biological material, magnetostratigraphy and geophysics;

. Based on what the group was shown, the Burin-Khan deposits (Fig. 1) appear likely to come into production before the Khubsugul deposits for geological, logistic and environmental reasons, although future research may change this evaluation; . Intensive surface investigations have already been completed at the Burin-Khan deposits, but at present no detailed information is available on lateral continuity. Such information is essential for accurate reserve/resource evaluation and will require: (i) deep trenching followed by intensive stratigraphic drilling; (ii) detailed logging (lithological, petrological, geochemical, gamma ray resistivity, self potential and neutron activation);

. The Burin Khan deposits will require detailed knowledge of the mineralogical, chemical and physical characteristics of the phosphorites in order to establish the most appropriate beneficiation techniques for both high grade and low grade materials;

. The structurally complex area will necessitate detailed structural analysis using appropriate geophysical and geological techniques;

. Since phosphorites provide valuable elements as by-products of fertilizer manufacture, extensive analysis for elements such as U, V, and rare earth, and studies of concentration and mode of occurrence should be undertaken. Mn and bauxite deposits are also associated with these phosphatic sequences;

. Some phosphogenic sequences are important source rocks for petroleum, which further justifies study of the Khubsugul sequence.

Most of the papers presented were devoted to the Late Precambrian-Cambrian phosphogenic period which has now been recognized in Asia, Australia, and elsewhere. Reports were given on the Paleozoic paleogeographic reconstructions with implications for phosphogenesis, paleo-oceanographic considerations of phosphate deposition, primary versus re worked origin of phosphate pellets, structural controls on phosphorite formation, and past excursions in seawater which may have resulted in enhanced phosphate deposition.

The general geology and Late Proterozoic-Cambrian stratigraphy of the Khubsugul Basin were examined in detail. Fossiliferous sequences containing trilobites, archeocyathids, hyolithids and algal structures were inspected with particular interest because of the importance of establishing the age of the phosphorites. Project paleontologists collected samples which will hopefully clarify age relationships.

The first phosphate deposit in the Khubsugul region was discovered in 1963 in the mountains of the western side of the Khubsugul rift valley which is an extension of the Baikal rift valley, one of the great rift systems of the world. Since that time, further work by the Joint Soviet-Mongolian Scientific-Research Geological Expedition has resulted in the discovery of phosphate resources totalling several billion tonnes of phosphate rock. Eight deposits have been found in the Basin, the Khubsugul deposits being probably the largest, underlying an area of about 500 km² adjacent to the western shore of Lake Khubsugul. The area is very mountainous with phosphate outcrops extending to as much as 1500 m above the lake level. The main phosphate unit is 60 to 90 m thick with up to five phosphorite horizons interlayered with dolomites. The most persistent is the lowest horizon averaging 10 m thick with 20-22% P₂O₅. Common lithologic associations include black cherts, siltstones, carbonates, and iron-manganese ores.

The Burin Khan deposits are located approximately 100 km south of the Khubsugul deposits in relatively low-lying terrain. Broken into several parts by faulting and by a large syenite intrusion, they are thicker and have a higher silica content than the Khubsugul deposits. The middle part of the series is richest in phosphate with 7 phosphorite horizons providing a total thickness of up to 55 m.

Plans are now underway to publish the abstracts, full papers, and scientific results of the meeting in Mongolian, Russian and English later this year.

B. Luvsandandsan, Geological Institute, M.P.R.
A.V. Ilyin, Institute of the Lithosphere, U.S.S.R.
A.L. Yanshin, Institute of Geology & Geophysics, U.S.S.R.
in collaboration with
P.J. Cook and J.H. Shergold,
Leaders, IGCP Project 156

From left, Dr. Luvsandandsan (M.P.R.), Dr. Tomor-Toogool (M.P.R.) and Dr. Zaitsev (U.S.S.R.) of the Joint Soviet-Mongolian Scientific Research Expedition during the meeting of IGCP Project 156 in Ulan Bator, M.P.R.

A group examines Cambrian sediments in the mountains west of Lake Khubsugul.

Attending a rodeo (Mongolian style) at the conclusion of the field workshop.

CONFERENCE REPORTS (continued)

ion was best summarized by the Chairman of the Scientific Committee who said: 'If an International Geological Correlation Programme did not exist, then it would be necessary to initiate one as quickly as possible'.

It is impossible to sit down and consider the achievements of the Programme, to read the vast volume of papers that it has generated, to scan the bibliography recently produced by the Secretariat, or to talk to participants in many of its Projects without feeling that a very powerful force for science and humanity has been generated."

Dr. McLaren went on to underline several ways in which the merits of IGCP could be made better known and the Programme's results be put to best possible use:

"1. Member countries of the Programme must be made more aware of its potential at the administrative and political levels. One step we can take immediately is to make our meetings more visible. For a start, at the meeting of the Scientific Committee and Board during the last two weeks of February next year, we hope to hold a special session and invite all member countries to send scientific representatives to a presentation on the Programme. We hope that scientific advisers from the many embassies in Paris will be able to attend.

2. IGCP must be identified in your own country. The scientists who now take part should be encouraged and supported to collaborate with their fellows elsewhere, thereby increasing the opportunity for the flow of knowledge and technology across borders.

3. The parent organizations, Unesco and IUGS, the Secretariat in Paris, the Board and Scientific Committee, the National Committees and Project participants within each country, must make every effort to demonstrate by specific examples the importance of collaboration between like-minded scientists and the ease of technology transfer when your own scientists are encouraged and supported.

4. Finally, there remains the question of support for projects in each country or region. The problem here is primarily financial. Access must be found to massive funding from

outside the Programme if its full potential is to be used for the benefit of development. Here, more than anywhere else, we need your support and advice. We need political awareness that the key to development or conservation of mineral, energy, and land resources rests within your own boundaries, and may be achieved by establishing a sound base of scientific and technical knowledge capable of benefiting from the major research effort that is now going on as a result of IGCP."

IGCP participants studying ring-dyke complex in the Coast Range Batholith, Peru.

Data in a Dynamic World

7th International CODATA Conference, Kyoto, Japan

The vast amount of data rapidly being collected by satellites, spectroscopy, biology and the earth sciences and the best use of these data bases in formulating new scientific concepts or discoveries constitute some of the most pressing challenges in science today.

In the first keynote address at the 7th International CODATA Conference in October, Dr. Philip H. Abelson (U.S.A.), editor of *Science* and a former president of IUGS, discussed the increasing impact of scientific data on science and society and the profound changes occurring in the creation, processing, storage, retrieval and transmission of scientific numerical data. Not only do new data bases facilitate the analysis of large volumes of data ("a single earth satellite provides more bits of information in a day than all of chemistry creates in years"), but they have also been useful in storing information on processes in nature which occur in a very short time, and have led to the current focus on picosecond chemistry. In addition new electronic instrumentation now permits measurements of weights as small as 0.1 picogramme of a hormone.

Prof. M. Kimura (Japan) outlined the way in which the DNA message contained in one chromosome set of 3.5×10^9 nucleotide base pairs long is determined. In his fascinating presentation, he stressed that during the last few years data on DNA sequences of various parts of one chromosome set (genome) in diverse organisms have been published at an accelerating rate which could now be termed "impossibly

Standing outside the Kyoto International Conference Centre are (left to right) Phyllis P. Glaeser, Executive Secretary of CODATA, Prof. M. Kotani, President of CODATA, and Prof. E. F. Westrum Jr., Secretary General, CODATA.

CONFERENCE REPORTS (continued)

explosive". While there is a great need for international cooperation to ensure that new sequences are made publicly available, it is also impossible to publish these in the formal sense. Hence, there is no alternative but to use a computer-based approach in solving this problem.

Dr. A. Ramachandran of the United Nations Centre for Human Settlements in Kenya described what has been done to plan data systems for development of human settlements. Physical indicators are combined with social, economic and cultural attributes in order to monitor the evolution of human settlements and provide some basis for a more aggressive or dynamic approach to effective human settlement policies and planning. It is hoped that this approach may be applied to three centres - in South America, Africa (Lagos) and India (Madras) - during the next few years.

The geosciences were strongly represented at the Conference. A thoughtful, almost philosophical presentation by Dr. Felix Chayes (U.S.A) compared the nature of phenomena with which the natural-historical sciences deal as opposed to the concepts of accuracy, precision and reliability of tightly controlled experiments in the physical sciences. Though much of the information in the geological base is literal and descriptive, scholarly control over the information available and data acquired more recently can be maintained.

Prediction of earthquakes and volcanic eruptions is a high-priority topic in Japan. T. Rikitake (Japan) reviewed the current monitoring network and techniques used to predict earthquakes of a magnitude greater than five; these include land deformation, seismic events, seismic wave velocity, changes in the geomagnetic and geoelectric fields, variations in earth resistivity, gravity, and levels of the water table, as well as behaviour of animals. In parts of Japan, legislation has been passed in order to deal with emergencies where the risk is high. In certain regions, however, issuing a warning to the public may result in an economic loss which could exceed the damage caused by the earthquake itself.

Precursors to volcanic eruptions may also cause damage, and D. Shimozuru (Japan) described case histories which now permit recognition of the rates of change in certain precursors of potentially damaging volcanic eruptions. Precursor earthquakes are seldom recorded prior to phreatic eruptions so other phenomena such as temperatures of vents and fumaroles or release of gases are monitored.

An intriguing study by R. Sinding-Larsen (Norway) focussed on the methodology used in Norway to integrate information from multiple domains in assessing the mineral resources of the Grong mining district. By a simple technique of generating binary surfaces for each of the geochemical, geological and geophysical variables available within a province, it is possible to evaluate the relative weights calculated for each variable. The binary maps for the different domains are then combined into one map which expresses a degree of association between a given area unit and mineralization. This is used to compile a map indicating the economic potential for each cell within a region. A practical result of this study has been the identification of three target areas, one of which is currently being drilled.

D. Gill (Israel) described a program to assess the areal value of the natural resources endowment of Israel. The comprehensive study suggested that Israel has probably discovered most of its near-surface mineral deposits.

Data bases of particular interest to geologists included the file of igneous rocks of Japan (presented by S. Aramaki) containing 8,000 records of analyses of variable format and quality collected during the last ten years. The most complete group is the file for Quaternary volcanic rocks of the Japanese Islands. This data base will be conformable with the IGBA file of IGCP Project 163 ("Design and generation of a world data base for igneous petrology").

Dr. Aramaki also showed how the current data base can be used for tracing the nature of some magmatic activity during the Mesozoic and Cenozoic. Similarly, J.M. Monget (France) indicated how remote sensing from satellites has provided an operational data base for use in oceanography. Though much can be done using available data, there is still the problem of standardizing data recorded from different satellites and a need to design improved data structures for processing in order to properly assess the various theoretical models proposed for ocean movements. K. Wadatsumi (Japan) described the GEODAS - an interactive graphic data entry and display system for geological borehole data.

Representatives of 23 countries attended the CODATA Conference. Proceedings will be published shortly and will be dedicated to the memory of Prof. T. Shimanouchi, who was chairman of the organizing committee until his untimely death in May 1980.

W.W. Hutchison
Co-Chairman, 7th CODATA Conference

CODATA Conference delegates visit the elaborate Kasuga Shrine at Nara, Japan. The shrine is famous for its rows of 3,000 stone lanterns of various shapes and sizes.

Chairman of the Organizing Committee, 7th CODATA Conference, Prof. Y. Mashiko with Mrs. Mashiko.

Exploiting Antarctica's Natural Resources

XVI SCAR Meeting, Queenstown, New Zealand
October 20 - 24, 1980

What scientific problems are related to the impending exploitation of Antarctica's living resources, as well as the possible future exploitation of her mineral resources? At the XVI meeting of the Scientific Committee on Antarctic Research (SCAR), a symposium dedicated to research in the Ross Sea region divided its discussion of the problems into three sections: marine and biological sciences, glaciology, and geosciences.

G.A. Knox (N.Z.) reviewed the oceanography of the Ross Sea, describing the five main water masses, their movement patterns, their abundant supercooled water and high nutrient levels, and their rich biological communities. Recent studies of Ross Sea fishes, birds, and seals were discussed by A.L. DeVries (U.S.A.), E.C. Young (N.Z.), and D.B. Siniff (U.S.A.) respectively. Work on the morphology, internal structure, and thickness of the ice sheet and the Ross Ice Shelf by the Scott Polar Research Institute was summarized by G. de Q. Robin (U.K.), with emphasis on airborne radio sounding studies since 1967. The main results of post-IGY research on the Ross Ice Shelf were reported by C.R. Bentley (U.S.A.) in a paper given by K. Jezek.

In the geosciences section, T.J.H. Chinn (N.Z.) discussed hydrology and climate, relating the hydrological regime to (i) ice sheet outlet glaciers, (ii) local alpine glaciers, and (iii) inland advances of the Ross Ice Shelf, noting that the dry valley lake levels are currently rising. T. Torii (Japan) described the varied geochemistry and high salinity of the dry valley lakes (suggesting multiple sources for the salts) with emphasis on the atmospheric component. I.B. Campbell and G.G.C. Claridge (N.Z.) reviewed the cold desert soils of the region, including some that are possibly as old as Miocene, and concluded that soil differences are related to age, climate, and parent rock composition.

The Cenozoic history of the area was reviewed by P.N. Webb (U.S.A.) who reported that in spite of important advances made since 1973 from four drilling projects, knowledge of the Paleogene still remains very meager. P.J. Barrett (N.Z.) spoke about the Taylor Group (Devonian) and the Victoria Group (Permian-Triassic), which together comprise the Beacon Supergroup - the Gondwana sequence in Antarctica, while M.G. Laird (N.Z.) discussed the lower Paleozoic successions (both about 8000m thick) in northern Victoria Land and the central Transantarctic Mountains. He speculated on their relationships to lower Paleozoic strata in New Zealand and SE Australia.

Precambrian rocks, representing at least two orogenic cycles, were reviewed by G.W. Grindley (N.Z.). Recent work on the Precambrian and Paleozoic rocks of northern Victoria Land, including some new interpretations, was described by F. Tessensohn (F.R.G.). Geophysical studies in the region were treated by F.J. Davey and D.A. Christoffel (N.Z.), including the present knowledge of Tertiary sedimentary sequences. D.P. Elston (U.S.A.) reported on the results of magnetic polarity studies on cores from several drill holes in the dry valleys and in McMurdo Sound, stressing the problems of establishing correlations between the drill sites.

A SCAR program called Biological Investigations of Marine Antarctic Systems and Stocks (BIOMASS) received much attention, especially its component projects on birds, seals, squid, and krill. A number of workshops have been tentatively planned for the next three years, and a special effort will be made to include scientists from developing countries in BIOMASS cruises. In addition, SCAR member nations will be asked for a special contribution to aid in establishing and maintaining a BIOMASS secretariat for the next decade.

After an introspective review of its structure, role, and future, SCAR modified its constitution. The SCAR executive committee was expanded to five members by adding the past president and a second vice-president. Each nation, although retaining a single vote, will be allowed to select a second scientist as an alternate delegate. New specialist groups created at the meeting included one on Environmental Implications of Possible Mineral Exploration and Exploitation and another on Antarctic Climate Research; the latter will plan the Antarctic component of the World Climate Research Program. The specialist group on Late Cenozoic Studies was disbanded, and its responsibilities were transferred to the Geology Working Group. The biology and logistics working groups met during the week preceding the SCAR meeting and an interdisciplinary symposium was held in conjunction with this meeting.

Concern was expressed about problems stemming from recent increases in the numbers of Antarctic tourists and adventurers. These visitors pose both a hazard to specially protected sites, and a nuisance (or worse) at small stations during mass visitations, as well as an unwelcome worry for search-and-rescue personnel. Controls or guidelines are needed, and SCAR is planning to publish "A Visitor's Introduction to the Antarctic and its Environment" to address these problems.

SCAR delegates during a cruise on Lake Wakatipu.

Field trip stop at Lake Hayes, five miles E of Queenstown, N.Z.

CONFERENCE REPORTS (continued)

Future SCAR Symposia were authorized as follows: on logistics - Leningrad, July, 1982; on earth sciences - Adelaide, August, 1982; on late Cenozoic coolings of the Southern Ocean - Moscow, 1982; and on biology - South Africa, 1983. A Polish proposal for an international program to study Antarctic plate boundaries was referred to the Geology and Solid-Earth Geophysics working groups. The term "Southern Ocean" was adopted for the circumpolar seas for use in SCAR documents, but its northern boundary was not defined. R.M. Martinez Abal (Argentina) was elected Vice-President of SCAR for the 1980-84 period. Subject to formation of a national Antarctic committee, the application from the German Democratic Republic for SCAR membership was approved. Two observers (a geologist and a biologist) from the People's Republic of China also attended the XVI SCAR meeting.

During the field trip organized on October 19, a traverse was conducted eastward to Cromwell over the upper Paleozoic (?) Haast Schist Group, ending at the construction site of the first dam for the ambitious Clutha Valley hydroelectric scheme. Included also was a short visit to Arrowtown, an important center during the gold rush more than a century ago.

The Royal Society of New Zealand plans to publish the symposium papers in a special volume.

Campbell Craddock
IUGS Delegate to SCAR

Field trip stop at Clyde dam site, first of five dams to be built in the Chitha Valley development scheme.

Hollis Hedberg Honoured by GSA

GSA's 93rd Annual Meeting, Atlanta, U.S.A., November 17 - 20, 1980

Former Chairman of the International Geological Correlation Programme (IGCP) and Past Vice Chairman of the IUGS Commission on Stratigraphy, Prof. Hollis D. Hedberg received in November the prestigious gold Penrose Medal from the Geological Society of America for his outstanding contributions to stratigraphy and petroleum geology.

Prof. Hedberg began his professional career in 1926, working as a sedimentary petrographer, micropaleontologist and stratigrapher for Gulf Oil Corporation in Venezuela where he lived for 20 years. He returned to the U.S.A. in 1946 to manage worldwide petroleum exploration for Gulf Oil, and became Vice-President of the Corporation in 1957. His long-standing association with Princeton University where he is now Professor Emeritus goes back to 1959. His research work on petroleum genesis, and particularly his papers on high-wax oil (1968) and on geothermal methane generation from organic matter (1974) are well known.

Equally commendable have been his contributions to international geology. Aside from his activities in IUGS, he has

been involved in the World Petroleum Congresses, the JOIDES Safety Panel, and the Law of the Sea. The "International Stratigraphic Guide" which he edited is now a best-seller, and he has recently completed collaborative work on a magnetostratigraphic classification chapter for the Guide.

Several major symposia drew large crowds at the 93rd GSA Annual Meeting. The Society of Economic Geologists (SEG) convened one entitled "Challenges of Future Mineral Resources," chaired by Brian J. Skinner. The need to make greater use of coal to attenuate the transition from hydrocarbon fuels to hydrogen or uranium fuels was stressed. In L.J. Miller's opinion, there is "an almost infinite supply of mineral deposits to supply continued growth of civilization." In addition to conventional sources, lower grade (but as yet uneconomic) sources were also described - lead-zinc in black shales and evaporite basins, for example. One conclusion was that significant challenges lie ahead not only in discovering additional reserves but also in improving the efficiency of the energy technology needed to mine deposits of increasingly lower grade.

A symposium on processes in karst hydrogeology (co-sponsored by IAH) was dedicated to V.T. Stringfield for his contributions to the study of karst hydrogeology in the U.S.A. Aside from an analysis of the nature of and flow in aquifers of the Dinaric karst in Yugoslavia, the karst regions described were mainly those of continental North and Central America. Common problems summarized by H.E. Legrand were the uneven distribution of ground-water sources, lack of soils, common lack of perennial streams, instability of the ground, and the unreliability of karst for waste disposal. The significance of the salt-water mixing zone in carbonate aquifers was summarized by B. Hanshaw and W. Back, who stressed that dolomitization, chertification and dedolomitization can all occur in this zone. In addition, it was suggested that "zones of high secondary porosity and permeability may result from dissolution processes" in response to sea level changes. It was concluded that such processes may account for some "paleokarsts" instead of surficial weathering.

Other major symposia included the David Johnston Memorial Symposium on the results of scientific studies of Mt. St. Helens, a public symposium which started with the presentation of a set of unique slides of the eruption on May 18, 1980.

Professor Hollis D. Hedberg, recipient of the Penrose Medal, and Mrs. Hedberg.

CONFERENCE REPORTS (continued)

Key-note speaker at the Annual Dinner, Michel T. Halbouty, renowned petroleum engineer and successful "wildcatter", outlined the plans for the "Decade of North American Geology" - a broad program of regional syntheses, map compilations of the continent, special publications - all dedicated to summarizing the latest knowledge on the geology of the North American continent which will form a basis for research and future exploration for mineral resources.

Held in Atlanta, Georgia, the GSA Meeting was attended by over 5,000 geologists and students.

Michel T. Halbouty

Publishing Today: Survival of the Fittest

AESE: 14th Annual Meeting, October 19-22, 1980, Halifax, Canada

In the face of rapidly rising costs of paper, printing, postage, ever-increasing inflation, drastic reduction of librarians' budgets, how can one rationalize the mysterious and continuous increase in the numbers of journals, books and technical reports being published in the earth sciences today? Who can afford them? What should earth science editors be doing about it?

According to Don Deneck of John Wiley & Sons, Inc., publishing in the 1980's will increase in proportion to the increases in global population and the spread of technology, which will demand dissemination of scientific information of all kinds: "The communication, quick and accurate, of salient technical advances is absolutely critical," he stressed.

But, cautioned Fred Spilhaus, Jr. of AGU, "there is no free lunch.... If any of us are to remain relevant to the problem of communication of scientific information, we must provide researchers with the information they need quickly and at low cost.... The road to relevance... lies in providing quality.... Publishers must put emphasis on what they accept in their journals. Libraries must put emphasis on quality publications in their collections. We must all establish mechanisms to judge what scientists need."

His provocative review of publishing economics reproached the library community for abetting the publication of marginal scientific journals at the time when budgets were "fat". Marilyn Guin of Oregon State University, expressing the librarian's point of view, retorted that it was up to the existing and new journals to re-examine their review processes, streamline their productions, curtail redundancy and exercise a more aggressive judgement on the worth of the published material.

The message that reverberated throughout the 14th Meeting of the Association of Earth Science Editors (AESE), held jointly with the International Association of Marine Science Libraries and Information Centers (IAMSILIC), was that literature today had to be relevant and affordable. Anyone approaching publishing in the next decade with a "non-profit attitude", warned Ronald Hart of AAPG, will soon run a deficit.

The editor, in short, can no longer afford to bury his head in technological and editorial expertise alone: a solid understanding of the rudimentary economics of publication is required.

There were many pragmatic suggestions. Gary Howell of ERICO, President of AESE, outlined the economic advantages of using editorial consultants over salaried staff, particularly when workloads become heavy. Claude Bishop, Editor-in-Chief at the Canadian National Research Council noted the effectiveness of using citation analysis for journal evaluation. Other sessions delved into the successes and failures of various manuscript review processes, of certain newsletters and newsmagazines, and a workshop was held on technical writing. Many of the discussions-from-the-floor turned to

the pros and cons of the microfiche concept of publishing, with most participants supporting it as a viable idea for the future, yet one that will still run into strong resistance from readers in the next decade.

An excellent, thought-provoking, sobering meeting, well organized by program committee members Ronald Hart of AAPG and Michael Latremouille of the Bedford Institute of Oceanography, in collaboration with Host Committee members Alan C. Grant and Bosko D. Loncarevic, also of the Bedford Institute.

AESE and IAMSILIC delegates at the lobster banquet. (Photo courtesy H.F. Wiele)

Newsletter session panelists (left to right): Peter J. Smith (U.K.), editor of *OPEN EARTH*; Judy C. Holoviak (U.S.A.), past editor of *Blueline*; Richard V. Dietrich (U.S.A.) session commentator; Patricia W. Dickerson (U.S.A.), session chairperson; Wendell Cochran (U.S.A.), editor of *GEOTIMES*; Vera Lafferty (Canada), editor of *EPISODES*. (Photo courtesy H.F. Wiele)

PIPELINES TO PEOPLE

IN UNIVERSITIES: More than 6500 faculty members at geoscience departments in the United States and Canada. **NEW** in this edition—specialty of faculty members will appear beside each name. Specialty Codes have been revised and expanded. Geographic locations of each college. Separate indexes of faculty—an alphabetical list and a specialty list based on the *new* specialty codes. Latest department telephone numbers, new or continuing chairman, list of departments offering field courses/field camps.

IN GOVERNMENT: Your direct line to the more than 2000 employees in the Geologic Division of the U.S. Geological Survey, listed alphabetically, with pictures, telephone numbers, current projects and research specialties. Includes USGS branch/office addresses and personnel for Reston, VA., Denver, CO., Flagstaff, AZ., Menlo Park, CA., as well as other domestic and foreign offices. Includes a separate roster of personnel in each branch or office of the Geologic Division.

Order From: American Geological Institute
Publications Department, One Skyline Plaza
5205 Leesburg Pike, Falls Church, VA 22041

Gentlemen:
Please send me the following:

_____ copies of the Directory of Geoscience Departments, 19th ed.
@ \$12/copy

_____ copies of the Directory of Geologic Division: U.S. Geological
Survey @ \$6/copy

Payment enclosed in US\$ _____ Invoice needed. Pur. Ord. No. _____

Ship To:
Name: _____

Address: _____

City/State: _____ ZIP: _____

AGI publication prices are subject to change without notice.

Charge to my credit card:

_____ Master Card _____ VISA

Acct. No. _____

Exp. Date _____

Signature (required) _____

Code: DGD2 Geo