Conference Reports . . .

Ground Truth and Remote Sensing Reviewed in Kenya

IGCP Project 143 Workshop, October 13 - 24, 1980
Nairobi, Kenya

Starting with a thorough review of remote sensing systems (particularly Landsat) and their applications to geological mapping and exploration, the IGCP Project 143 Workshop on Remote Sensing and Mineral Exploration concentrated on visual interpretation of a Landsat scene of the Lake Magadi area (East African Rift Valley), SW of Nairobi. Drainage, bedrock, and lineament overlays were made so that workshop participants could become familiar with the geology of the area.

Another day was devoted to a field trip to Lake Magadi, where participants visited a soda ash roasting and bagging plant. A dredge floating on the lake is joined to the plant by a slurry pipeline several km long. The history, development, and current production from Lake Magadi has been described by Birtwistle (1971): annual production of about 200 000 metric tons of ash is mostly exported, with the Philippine glass industry being the largest purchaser. More recent studies on the hydrogeochemistry and mineral reactions of the Magadi deposits have been described by Surdam and Eugster (1976) and Jones *et al.* (1977). The mineralized surface near the lake consists of vertical, closely packed crystals which felt like a thickpile rug underfoot. Large polygonal plates several m across show overthrusting of several cm along their margins, suggesting considerable expansion of the surface as the minerals crystallize. Algal blooms are abundant, creating a pink hue in both the water and dry mineral surface.

During a two-day safari to Voi and East Tsavo National Park, participants observed a major Holocene basalt plateau which fills an ancient channel of the Athi River. On the Landsat scene covering the area, this particular formation stands out in marked contrast to the surrounding Precambrian terrain. Along the road west of Voi, the group saw outcrops of white crystalline dolomitic limestone which in one locality is worked for cement. The upper surface of the dolomite is marked by a tan, oolitic, fragmental conglomerate(?) at least 1m thick. It was sampled in two places for closer examination and analysis for phosphate.

On the Taita Hills trip, led by R.E. Okelo of the Kenya Geological Survey, the group found that the Precambrian rocks are made up of the Lower Kuruse Group composed of thickly layered biotite gneiss. The strata (strike E-W, dip 10-15° N) are conformably overlain by the Kasigare Group, composed predominantly of quartz-feldspar gneiss. The units form prominent, steeply sloped hills above the surrounding, gently rolling plains. The hills are, in many places, soil covered and farmed, and foster the growth of a wide variety of vegetables and fruit.

This field trip included a visit to a locality in which tsavorite, a green garnet, was being mined from narrow pegmatite veins in biotite gneiss and schist. (Tsavorite is a green garnet and a chromium-rich member of the grossularite group.) The garnet vein strikes north and dips 45°E, and is being mined to a width 1 m. Hand cobbing is done underground. Waste rock is backpacked to the top of the workings and dumped, after which it is again given careful scrutiny.

The final three days of the workshop were devoted to discussion of case histories using Landsat data. W.D. Carter outlined how the "Photogeologic Map of Botswana" (Mallick *et al.*, 1978) was an excellent example of new map products and useful symbology. A. Stancioff (USAID/Zaire) presented an overview of the geology and mineral resources of Zaire; he reported that a preliminary mosaic of Zaire had been completed, and that Landsat data were being used to study the Congo Basin and the Precambrian rocks of the SE, where the copper belt as well as tin, lead, and zinc occur near the western boundary of the East African Rift. The Shaba Plateau of SE Zaire contains a tin-bearing pegmatite, 12-14 km in diameter, where 200 000 tons of cassiterite, spodumene, and columbite - tantalite have been mined. The NE plateau also has potential for gold. To the west on the continental shelf, offshore drilling for oil is being conducted by Gulf Oil Co. and Texaco.

Stancioff also reported on the successful exploration for zinc deposits using Landsat data in northern Newfoundland, Canada, in which 100 000 metric tons of 4% zinc were found in the Lower Ordovician St. George Formation, a stromatolitic dolomite. The deposit, however, is considered too small and of too low a grade to be mined. Stancioff presented a third case history in which areas for exploration of placer diamonds were located by correlating areas of uplift to changes in drainage patterns in the Central African Republic.

A. Fernandez (UNDP, Mozambique), former Director of the Servicio Geologico of Bolivia, spoke on Bolivia's extensive use of Landsat for geological mapping, mineral exploration, land use, and industrial planning (Brockmann *et al.*, 1979). His presentation was especially useful, because development in Bolivia has closely resembled the approach that many African nations may wish to use in analysing Landsat images.

W.D. Carter discussed the use of interactive digital analysis of Landsat images for geological investigations by demonstrating an application relating to the spectral measurement of chromite-bearing ultramafic rocks near Mysore, south-central India.

Workshop participants study images of the Serengetti Plains south of Lakes Magadi and Natron.

Lake Magadi: overthrust plates of the crystalline natron surface layer reflect expansion of surface by polygonal fracturing and crystal growth during desiccation.

CONFERENCE REPORTS....

W. Bekele described the general geology and mineral resources of Ethiopia. Placer deposits of platinum and gold associated with belts of ultramafic rocks are mined by local residents near the Sudan border and in south-central Ethiopia. Recently, Canadian firms discovered 70 million tons of nickel ore (1%) using satellite images. In northern Ethiopia, Nippon Exploration and the Ethiopian Geological Survey are exploring a Cu/Pb/Zn belt, while in the Ogaden Basin, Tenneco completed eight drill holes in 1972-73 with shows of petroleum and discovered a small gas field of 45 million m³. Bekele spoke of high fluorine content in geothermal waters which may have some commercial value. Potash (16 million tons) has been developed by the Parsons Co., U.K.

Prof. F. Ahmed (Sudan) described the geology and mineral resources of his own country, the largest in Africa, covering over 2.5 million km². Chrome-nickel deposits occur along its NE border in ultramafic rocks that are thought to represent old subduction zones. Late Proterozoic granites and pegmatites contain mica-bearing pods and gold-bearing quartz veins. Tin/tungsten deposits occur in 60 ring-dike complexes that have been defined thus far. Mesozoic (Triassic) sedimentary rocks consisting of sandstone, mudstone, limestone, and oil-bearing shales have been identified by Chevron Oil Co. A remote sensing centre is to be established in Khartoum by the National Research Council of Sudan.

L.C. Rowan referred to the work of Raines et al. (1978) in conducting base-metal exploration in Sonora, Mexico, in cooperation with the Mexican Government. Following analysis of Landsat images of the basin and range areas of northern Sonora, Raines confirmed Salas' earlier assertions that important sets of NE- and NW-trending lineaments, spaced approximately 40 to 50 km apart, exist in the area. Geochemical sampling (Raines et al., 1978) and airborne and ground geophysical surveys based on the lineament patterns, revealed anomalous ground at one of the intersections. Subsequent drilling revealed mineralized rock to a depth of over 200 m. Exploratory development drilling on the deposit continues.

J.C. Rivereau (France) described the AFRICASAT Study, which covered 5 million km², encompassing 250 Landsat scenes compiled into mosaics at 1:1 million scale and reduced to 1:2 million. This is a lineament and bedrock map study of northern Africa extensively supported by bibliographic references. It distinguishes between "active rift" in Ethiopia and "passive grabens" further south. Completion of the structural model and map synthesis took 8 months by three geologists and could have been done only with Landsat data. Rivereau also presented a detailed lineament map of NW Africa and S Spain showing the extension of linear features across the floor of the Mediterranean Sea (Rivereau, 1977).

P. Winani (Kenya) lecturing on the geology and mineral resources of Kenya, described airborne geophysical surveys, conducted in 1977, that were successful in finding graphite deposits. Four kimberlite pipes were also discovered on the east side of Lake Victoria but have not yet been tested for diamonds. Fluorspar, soda ash, and salt are the main mineral products of the Rift Valley. A geothermal study was conducted under UNDP auspices in 1969-71. Japan is now providing technical aid to explore and evaluate geothermal possibilities in the area just west of Lake Naivasha. The U.K. Institute of Geological Sciences has just signed a U.S. \$10 million agreement with the Kenya Geological Survey to map mineral resources in a large area east of Lake Turkana. A French/Finnish project is also underway to evaluate and explore industrial mineral occurrences.

Participants were given the opportunity to evaluate the workshop. The interpretation exercise and field trips were found to be most useful, and many commented on the value of the national case histories. Almost all felt that the workshop should have been longer, and that it should be repeated elsewhere.

References

- Birtwistle, D., 1971, Lake Magadi: Magadi Soda Company Reprint updated in 1977, 6p.
- Jones, B.F., Eugster, H.P. and Rettig, S.L., 1977, Hydro-geochemistry of the Lake Magadi Basin, Kenya: *Geochemica et Cosmochemica Acta*, v. 41, p. 53-72.
- Mallick, D.I.J., Habgood, F. and Skinner, A.C., 1978, Photogeologic map of Botswana: Institute of Geological Sciences, London, England. (Map Scale 1:1 000 000, 2 sheets)
- Raines, G.L., Theobald, P.K., Kleinkopf, M.D., Lee Moreno, J.L., and del la Fuente Duch, M.F., 1978, Base Metal Exploration Case History, in 5th Symposium of the International Association on the Genesis of Ore Deposits: Program and Abstracts, Snowbird, Utah, p. 148.
- Rivereau, J.C., 1977, Contribution of Landsat imagery to geological prospecting over the western Mediterranean areas, in 3rd W.T. Pecora Symposium: Abstracts, Sioux Falls, S. Dak., October 1977. (See also "The Alboran Sea Structural Pattern; an offshore extrapolation of Landsat Imagery Interpretation", Bureau d'Etudes Industrielles et de Cooperation de l'Institut Française du Pétrole (Beicip) Paris, France, 3 p., 4 figs.)
- Surdam, R.C., and Eugster, H.P., 1976, Mineral reactions in the sedimentary deposits of the Lake Magadi region, Kenya: *Geological Society of America Bulletin*, v. 87, p. 1739-1752.

**W.D. Carter and L.C. Rowan (U.S.A.)
Co-Leaders, IGCP Project 143**

Dredge working soda ash at Lake Magadi.

The Pan African

**11th Colloquium on African Geology, The Open University,
Milton Keynes, U.K. April 13 - 15, 1981**

If a consensus of opinion emerged from the many papers devoted to the Pan African at the 11th Colloquium on African Geology, it was that there was no uniform development on the continent, that in parts a fully developed Pan African "Orogeny" can be traced, whereas in others a less distinct Pan African "Event" dominated. Plate tectonics seem to have played a more prominent role than during the earlier time units.

About 130 participants listened to the papers delivered in parallel sessions during the three days of the colloquium. Papers were divided into the following sections: Pan African and basement in Saudi Arabia and NE Africa (17 papers); Geophysical studies of the African crust (14); Hot Spots (2); Pan African of Namibia (4); Mobile Belts of Southern Africa (3); Pan African of Mali and Hoggar (7); Pan African and basement in West Africa (4); Precambrian of Central Africa (3); Young Volcanic Rocks (6); Sedimentary Basins (2); Neotectonics (4); Ring Complexes (9); Economic Geology (6). A total of 9 contributions devoted to mineral resources, a poster session and a one-day excursion to the "Pan African"

CONFERENCE REPORTS...

of southern England (at Charnwood Forest) completed the agenda.

The colloquium was aimed mainly at exchanging information among European researchers working in Africa; the number of participating African geoscientists, except for African

postgraduates currently in Europe, was unfortunately quite limited.

The 12th Colloquium in 1983 will be organized by the Royal Museum for Central Africa in Belgium.

Dr. G. Matheis, F.R.G.

Afrikagruppe deutscher Geowissenschaftler

Formation and Evolution of the Hellenic Arc and Trench

International Symposium on HEAT, April 8-10, 1981

Athens, Greece

At 20:57 hrs. GMT on February 24, 1981 an earthquake of magnitude 6.6 on the Richter scale occurred in the vicinity of the Greek islands of the Alchiomides. At 02:36 hrs. GMT the following day, an after-shock of magnitude 6.3 struck the same area. Nine days later, at 21:58 hrs. GMT, a magnitude 6.2 earthquake occurred 20 km away near the town of Napareli, and was followed by a 5.9 magnitude after-shock on March 5.

As might be expected, these major events received much attention at the International Symposium on the Hellenic Arc and Trench (HEAT) held in Athens the following month. Organized by the National Technical University of Athens and various research teams working on the Hellenic Arc and Trench, the symposium was held under the auspices of the Ministry of Culture and Science of Greece. Co-sponsors were the Geophysical Committee of Greece, IUGS, and the IUGS/IUGG/ICSU-sponsored Inter-Union Commission on the Lithosphere (ICL). The well-organized sessions held in the auditorium of the Chamber of Commerce and Industry were attended by more than 250 scientists representing 25 countries.

The symposium was dedicated to the formation and evolution of the Hellenic Arc and Trench and their relationship with adjacent and similar areas, particularly the Tyrrhenian Sea and Pannonian Basin areas. As Professor S.S. Augustithis (President of the Symposium's Organizing Committee) said in his opening address, the emphasis was on "deep lithospheric and asthenospheric processes responsible for the dynamics of the area...." The main questions addressed during the three-day meeting, therefore, included whether the Hellenic Arc and Trench were formed from the subduction of the African plate beneath the Aegean plate, and if the area was one of active subduction.

Following an excellent summary by X. Le Pichon (France) of the HEAT Project (see EPISODES, Vol. 1979, p. 3-6), E.A. Flinn (U.S.A.), Secretary General of ICL, outlined the objectives and organization of the Lithosphere Program. The many excellent papers that followed generally concluded that the Hellenic Trench system is a well-developed subduction zone produced by the collision of the African and Aegean plates.

In a paper by B.C. Papazachos and P. Comminakis (Greece) on deep tectonics and earthquake prediction, data were presented on intermediate focal depths that confirmed the presence of a well-developed Benioff zone. X. Le Pichon pointed out that strain patterns observed from Sea Beam data indicated that the axis of the Hellenic trench forms a sharp boundary between strain patterns of different character - compressional on the south side, and extensional on the north. These strain patterns are likely influenced by the Libyan Promontory which may be considered a prelude to continental collision.

This view is supported by G. Anastasakis (Greece) and G. Kelling (U.K.) who have described the area south of the trench as one exhibiting compressional stresses characterized by thrusting and reverse faulting, while the fore-arc (northern part of the Ptolemy Mountains) is one of extensional stresses characterized by vertical tectonics (for example, the uplift of Crete) with sub-parallel faults and grabens immediately north of the perched basins area.

Forty-six high precision gravity base stations have been established in the Aegean region to form a precise network for observing gravity variations with time. W. Torge (F.R.G.) and colleagues believe it is possible to measure changes in elevation of 3-5 cm, thus providing a long term method of measuring shifts of vertical masses. An evaluation of the 18 000 gravity stations in the Aegean region and additional marine gravity data has been made by Y.J. Makris (F.R.G.) and colleagues who applied a polynomial surface to fit their data in an attempt to separate the regional field from local effects. The authors' interpretation indicated that the Hellenides are spreading outwards toward the Ionian Basin due to plastic deformation of the continental crust, controlled by the upper mantle lithospheric system and not by subduction.

Paleomagnetic studies of the Miocene-Pliocene series of the Hellenic sedimentary arc, undertaken by M. Jamet (France) and others, was directed toward determining possible rotational deformation. The investigators' results showed that the pole position for Crete was essentially the same as for

Left to right: Prof. S.S. Augustithis (Greece), President of the Organizing Committee of the International Symposium on HEAT, confers with Prof. X. Le Pichon (France), a member of the Symposium's International Scientific Committee.

Symposium participants include, in the front row (left to right): Prof. G. Parisakis, Vice-Rector of the National Technical University of Athens, Dr. C.C. Weber, Secretary General of IUGS, Mrs. Trumpy and Prof. R. Trumpy, Past President of IUGS, and Prof. S. Katrakis, Emeritus Professor, National Technical University of Athens.

CONFERENCE REPORTS...

Africa, indicating no rotation during the past 7 Ma (since Tortonian time). The situation seemed to be the same for Rhodes, but here the lake bed sediments measured were younger (4-5 Ma). Corfu, on the other hand, showed a definite clockwise rotation of approximately 25 degrees since middle Tortonian time (8-10 Ma) and a current rate of 3 degrees per Ma.

Several timely papers were presented on the two recent seismic events in which deformation in the Corinthian Gulf

area produced significant damage to class-C masonry. The two sets of faults observed trend generally E-W for a distance of 9-12 km, forming a graben in between. The faults are right-lateral normal and the relative motion between the blocks is opposite. Smaller structures observed indicated that the deformation is not pure extension and that some rotation is present.

Gordon Andreassen, U.S.A.
Geophysicist, USGS

European Geoscientists: Multiplicity of Approaches

European Union of Geoscientists: First Meeting, April 13 - 16, 1981, Strasbourg, France

The originality of modern earth (and planetary) sciences is without doubt the result of a duality. This duality rests on both the multiplicity of the approaches (going from the frontiers of biology to mathematics and astronomy, sweeping through most of the physical and chemical sciences, relying on theoretical computations, on laboratory experiments, on field work), and on the extraordinary convergence of the goals which must be attained: unity in diversity.

The thoughts expressed by Claude J. Allègre, President of the new European Union of Geosciences (EUG) in his welcoming address received the full support of 600 enthusiastic geoscientists attending the Union's first meeting this spring. The 400 or so papers presented during ten symposia and even the eight additional sessions on specialized themes which were added to the original schedule were a testimony to the multidisciplinary spirit that prevailed.

Some highlights and achievements of the meeting follow.

- The fundamental similarities and differences between ophiolites and greenstone belts are becoming slowly better understood, and a consensus was reached regarding greenstones probably being indicative of huge volumes of ultramafic magmas formed through the hitherto unknown, but radically different, processes of Archean plate tectonics. Rare modern komatiites within ophiolites (Gorgona, East Pacific) are providing useful clues for such studies.
- The subject of earthquakes received much attention, the focus being on the recent major Algerian (El Asnam, 10-10-80), Italian (Campania, 11-23-80) and Greek (Korinthos, 02-24-81) earthquakes, but also on the less destructive Sierentz (07-15-80) earthquake in Alsace. No major breakthrough was reported on earthquake prediction methods, although M. Caputo (Italy) did predict, on the basis of a statistical/seismic-gap method, one or two more major earthquakes in Southern Italy during the next few years.
- Recent biochemical studies of organic molecules have shown that isomerization and aromatization depend as much on the temperature reached as on the heating rate. Geophysicists have simultaneously shown that the geothermal gradient was higher during the first 150 Ma of a basin and dependent on the rate of extension during that period. These two approaches yield a maturation index based on molecular reactions, which is considered better than the indexes currently used commercially.
- The symposium on the early development of the Earth and planets provided an opportunity to improve the present models for accretion and outgassing of the Earth, a planet with an original Venusian-type atmosphere, to compare the data and models available for various planetary bodies throughout the solar system, whether observational or experimental.
- A session on hydrothermal processes focussed mainly on the "black smokers" (superheated steam depositing sulfides) of the East Pacific Rise, and on related metal-rich deposits; hydrothermal activity away from the ridges (for example, the alteration of basalts) was also discussed.
- Contributions to the study of magma generation and segregation dealt with mixing, partitioning, and the like, and with the dynamics of magma transfer to the surface, both from a mechanical point of view and on the basis of interpretation of isotopic (Th) geochemistry.

- Episodicity in accretion is now well established, and when averaging opening rates throughout time intervals shorter than two million years, the postulate of rigid plates does not hold any more. For collisions too, the episodicity depends on the scale of observation: an orogeny as a whole may seem to be a continuous process, though the complex shape of the continental margin may yield successive discontinuous tectonic events at a regional scale.
- Chemical transfers as analyzed through isotope geochemistry now yield not only static models of theoretical reservoirs, but truly dynamic models of the Earth - hence, a whole new field referred to as "chemical geodynamics" by C.J. Allègre.
- A P-T plane symposium broadly reviewed the problems related to geotherms, with geochemists admitting the often neglected importance of cooling rates and other time effects, and geophysicists presenting both theoretical and field evidence for the actual thermal regimes.
- Paleoclimates throughout the whole span of the geological ages were discussed, with special emphasis on relatively recent times, culminating in some short-term predictions: for example, very hot weather lies ahead of us for the end of the century, both on the basis of recent studies of geomagnetic precursors and of statistical studies of dry Sahel cycles.
- A symposium on geological and planetary phenomena and the evolution of species led to debates on what caused the apparent sudden extinction of the dinosaurs, with most concluding that the paleontological record is not as yet sufficient for a decision on whether this extinction was really synchronous all over the planet and how long the event lasted; the hypothesis of a simple meteorological impact was rejected on the basis of its inability to explain certain enrichments other than Ir.
- A unique session devoted to zircon chronology demonstrated how crystal isochrons may be obtained through "microsurgery" of a zircon grain.

The sessions were varied and stimulating and generally well attended, with the exception perhaps of some of the more "classical geology" ones. The quality of the Union's first meeting will serve as an inspiration to geologists in all fields contemplating attending the next one.

Jean Claude Mercier, France
Member, Founding Council, EUG

Claude J. Allègre,
President of the EUG.

CONFERENCE REPORTS... Calgary Hosts Three Associations

GAC/MAC/GSU Joint Annual Meeting, Calgary, Canada

May 11-13, 1981

International geoscience constituted an important recurring theme at the annual meeting of the Geological Association of Canada (GAC), held in conjunction with the Mineralogical Association of Canada (MAC) and the Canadian Geophysical Union (CGU). For example, Prof. R.A. Price, President of the Inter-Union Commission on the Lithosphere, stressed that although the study of oceanic lithosphere will represent a significant component of the work planned for the new program, it is the continents and the interaction between the continents and oceanic lithosphere that will be the principle focii. With the oldest sediments in deep oceans being younger than 200 Ma (less than 5% of the Earth's history) the continents must be studied to unravel the remaining 95% of the Earth's history. He drew attention to the International Geological Correlation program and some of its more successful projects, as well as the recently launched Decade of North American Geology, a project of the Geological Society of America involving various organizations in Mexico, U.S.A., and Canada.

In a stimulating keynote address, D. W. Strangway briefly reviewed the role of fluids in the geological processes of planetary bodies other than the Earth. He described the lack of fluids (and atmosphere) on the moon, which results in vacuum welding of the shattered surface layers, giving the moon quite different seismic properties from those of the Earth. The lack of evidence of fluid in moon rocks suggests that any economic mineralization will be primarily related directly to magma segregations; deposits such as those resulting from hydrothermal activity are presumably lacking.

In many ways, the planet Mercury is quite similar to the Moon, while the planet Mars is strikingly different, with features such as braided streams which may have resulted from flash floods, mud flows, and the reddening of surface materials due to oxidation and hydration. Venus, with its highly dense atmosphere, still remains very much of a mystery, but its present day surface temperature is so high that it is unlikely that any water can be present. By contrast, some of the moons of Jupiter appear to be largely composed of ice resulting in very curious surface features. The fact that most of the Earth's surface has remained within a temperature variation of less than 100°C for possibly over 3.5 Ga is therefore remarkable.

T. Urabe (Japan) and S.D. Scott (Canada) set the stage for the symposium on new exploration guides for Kuroko-type massive sulphide deposits with a comparative analysis of Kuroko and Archean massive sulphide deposits. These volcanogenic massive sulphide deposits were formed within island arcs in association with bi-modal, rhyolitic-basaltic volcanism. Syn-volcanic acidic plutons or dykes are also common and are considered to have played significant roles in the formation of the deposits both as sources of heat and the critical metallic elements. Studies of the host rocks using oxygen isotopes has given evidence for the size of the hydrothermal fluid circulation system which was a major cause for some of the alteration haloes that extend as far as 1 500 m or more from some orebodies.

Of special interest too were the various presentations on metallogeny and tectonics of the North American Cordillera, and in particular, parts of the Yukon and the Northwest Territories. A spirited presentation by D.J. Templeman-Kluit (this year's recipient of GAC's Past President's Medal for outstanding work in the earth sciences by a young Canadian earth scientist) described the two geological elements - the Paleozoic and older sedimentary strata deposited under relatively stable conditions near the western edge of ancient North America, and, to the SW, those strata dominated by Mesozoic sedimentary and volcanic rocks and including some older metamorphic strata. The SW block is conceived to have accreted to the North American continent and thrust NE above it during the Late Jurassic, thus forming a line of

junction called the Teslin Suture. The economic significance of this interpretation is critical because it has led, in the E belt, to the recognition of conformable lead-zinc deposits restricted to North America which occur in three stratigraphic associations around what is known as the Selwyn Basin. R.W. Macqueen and J.F. Barker focussed on the organic geochemistry of the shale-hosted, lead-zinc deposits of the Basin. The sulphide-rich zones generally have lesser amounts of organic matter than some of the enveloping rocks.

The organic carbon ranges from 2 to 14%; C13/C12 ratios of associated calcite ranges from -3 to -9‰, thus suggesting some biological interaction with the organic matter; pyrolysis yields from organic matter are abundant over a temperature range from 350-700°C. The data suggest that the temperature range for the sulphide deposits has been relatively high, possibly above 200°C, which aids, in combination with the data on the organic matter, in interpreting the sulphide mineralization process. It is believed that some of these deposits may become major sources of lead and zinc in North America.

A key paper on late Precambrian plate-tectonics by P.F. Hoffman focussed on the complex series of faults resulting from the terminal continental collision of early Proterozoic age in the Bear Province of the Canadian Shield. He described how the north-central part of the Wopmay orogen is broken into overlapping crustal shingles about 70 km wide and is bounded by a network of conjugate transcurrent faults and thrusts dipping west. These features represent the youngest compressional event in the orogen (believed to be younger than 1.87 Ga). Detailed analysis of the faults indicates that 9 km of convergence can be documented between two of the main shingles. In addition, backsliding on the thrusts during post-orogenic extension is believed to have resulted in the formation of graben-like basins. The shingle-like structure evident in this part of the Precambrian appears to be similar to shingle structures forming today on a large scale in south-central Asia.

W.W. Hutchison, Canada
Past Secretary General, IUGS

Left to right: GAC President W.G.E. Caldwell, W.S. Fyfe, recipient of the GAC Logan Medal, and N.J. Allman, incoming President of GAC.

Left to right: D.J. Templeman-Kluit with D.F. Sangster, recipient of GAC's Duncan Derry Medal.

Exploration for Gas Fields

UN/ECE Seminar, Exploration for Gas Fields
May 18 - 22, 1981, Hannover, F.R.G.

The United Nations/Economic Commission for Europe-sponsored Seminar on the Exploration for Gas Fields in the ECE Region started off with a comprehensive, well illustrated review of the geology of the gas-bearing provinces of the ECE region by H.-A. Hedemann (F.R.G.) - excellent background for geological, geochemical, and geophysical reviews that followed. Another introductory lecture (by B. Tissot of France) on the origin of natural gas in the ECE region supported the modern concept of maturation in spite of some strong reservations among some of the participants who prefer the "multiple working hypothesis".

The Working Session on Geology/Geochemistry was chaired by H.A. van Adrichem Boogaert (Netherlands), who summarized and discussed the papers that were to be presented, accepted the maturation concept as the explanation of the origin of gas, and stressed the importance of maturity of the source beds. His views were supported by at least two of the geochemical papers. Informative case histories were presented on the Namur Basin, the Northern Alpine Thrust Belt, the Vienna Basin, Czechoslovakia, the deep Dnieper-Donetz Basin, the North Sea area, and the new East Hannover Gas Province. These were examples of the success of the exploration efforts in these countries and it was made clear that exploration has yielded good results elsewhere in the ECE region. As the Chairman pointed out, exploration is being directed more and more toward deeper targets, and a thorough study of the geology, both stratigraphical and structural, is needed in order to justify drilling of costly ultra-deep wells. The experience in the U.S.S.R. and the U.S.A. in this regard has been favourable.

The Working Session on Geophysics was chaired by H.J. Dürbaum (F.R.G.); the majority of the geophysical papers dealt with the seismic method. The Chairman stressed the importance of the magnetic and gravity methods and how these have been improved by the computer age. Another non-seismic method - the magnetotelluric method, which uses naturally-induced electromagnetic disturbances, permits deep penetration. One paper reviewed the results of the application of this method in the pre-Zechstein Basin of NW Germany and in an overthrust area at the southern boundary

of the Po plain. Due to the development of the remote-reference method, which greatly improves the signal-to-noise ratio, it is predicted that the magnetotelluric method will find more application in the future. Dürbaum stressed the need for improvement of the signal-to-noise ratios as the most important problem facing the seismic method. Stacking and deconvolution are two main methods currently in use.

A tribute was paid to Peter Vail and his co-workers for their pioneering work on seismic stratigraphy. In this session, papers dealt with slant stacking, a study of the Frigg gas sands in the North Sea, 25 years of looking for stratigraphic traps in the Po Basin during which 14 new fields were discovered (up to 1964) with total recoverable reserves of 35BCM, and field data from NW Germany.

Direct detection of hydrocarbons by recognition of "bright spots" and "flat spots" created great enthusiasm, but there are also many pitfalls to this method. Three papers dealt with this problem competently and improvements in methods were suggested. A novel idea was the application of the method to monitor the Hähnlein gas storage reservoir in the Rhine Graben, beginning before the injection of any gas. The Seminar succeeded in stimulating exchange of scientific ideas and experiences, especially in the fields of bright-spot interpretation and seismic stratigraphy, and it was recommended that such exchanges be continued and extended to include the magnetotelluric method as applied to exploration for gas at great depths.

A paper on a recent survey of undiscovered gas resources of the U.S.A. gave a bright outlook for the future and a stimulating example for the ECE countries.

The Seminar was complemented by two enjoyable Study Tours - to the old established fields of the Oldenburg South District, north of Hannover, and to the newly discovered gas fields east of Hannover. A fascinating exhibit was available, commemorating Alfred Wegener's life and his contributions to the geological sciences.

W.C. Gussow, Canada
Petroleum Consultant

Mapping Half the World

Circum-Pacific Map Project: 7th Annual Meeting, May 27-30, 1981
Menlo Park, California, U.S.A.

More than 50 geoscientists from 10 countries gathered in California in May to review the status of map compilation and to finalize guidelines for future maps in the Circum-Pacific series. With the May release of the two eastern quadrant maps in the Plate-Tectonic Series, the Map Project is moving from the compilation stage to completion and publication of the thematic map series.

To date, 14 of the 44 geographic and geological maps have been published, the 5-map 1:10 000 000 scale Plate-Tectonic Map Series will be completed during August 1981, the Panel Chairmen have reviewed proofs of the NW and SW Quadrant map sheets, a working copy of the 1:20 000 000 scale Plate-Tectonic Map, being compiled by Kenneth J. Drummond, NE Panel Chairman, has also been reviewed, and the basin-wide map is scheduled for publication by year-end 1981.

It was decided at the Annual Meeting to prepare and publish a short paper explaining the plate-tectonic maps and the development of the various data sets to accompany the map series. Project work over the next 12 months will focus on the Geologic and Tectonic Map Series.

A review of experimental proofs of the equatorial Pacific/-Caribbean region of the NE Quadrant Geologic Map led to the decision that experimental depiction of seabed geology in concert with sea floor sediments be further improved.

Panel Chairmen reported that compilation of land geology for all map sheets has essentially been complete and final

cartography of the NE and SE Quadrants was scheduled for completion by mid-1982. Reconsideration of guidelines for the Tectonic Map Series, begun during the meeting and dealt with in working group meetings, will be completed by the end of July 1981. The Panel Chairmen agreed to have these maps compiled prior to May of 1982.

Compilation of the Minerals Resources Map of the NE Quadrant was illustrated by a proof of the land areas. It was decided that for land areas, data from the Geologic Maps will be used as background for the mineral deposit symbols with selected tectonic elements to be added, as needed, by the Panel Chairmen. Bathymetry and possibly generalized sediment distribution will be used as background for the manganese nodule abundance and chemistry data for ocean areas. The NE Quadrant map was scheduled for completion by December 1, 1981.

Proofs of the NE and SE Quadrant Fossil Organic Fuels Resources Map were displayed during the meeting. A special working group considering this series of maps recommended extensive revision of compilation guidelines, including emphasis on sedimentary basins, geological age of producing strata, and inclusion of representative stratigraphic columns for basins with "giant" status.

Discussion of the Geodynamics and Geothermal Map Series led to the designation of a panel of specialists to deal with gravity coverage for the map series. The group will start the

CONFERENCE REPORTS...

gravity compilation in early 1982.

The meeting concluded with a field trip to the Oakland Museum to view newly developed plate-tectonic exhibits including a large, world-wide fiber optics model of active plate boundaries, seismicity, volcanism, and mineral deposits.

Next year's meeting will be held during the first week in May, again in California. A report, similar to last year's USGS Open-File Report 80-1059, summarizing the meeting and Project status will be issued later this year. Copies of present and future reports are available on request from the Circum-Pacific Map Project, 345 Middlefield Road, MS 52, Menlo Park, California 94025, U.S.A.

W.O. Addicott, U.S.A.
Deputy Chairman, Circum-Pacific Map Project

Index map of the Pacific Basin showing Circum-Pacific region and map sheets of the Circum-Pacific Map Project.

Meeting participants including W.O. Addicott, Map Project Deputy Chairman (front row, left); C. Craddock, Antarctica Panel Chairman (front row, second from left); C. Nishiwaki, NW Panel Chairman (second row, third from right); C. Martin, SE Panel (second row, second from right); W.Y. Chang, NW Panel (third row, third from left); Frances Delany, CGMW (third row, fourth from left); H.F. Douth, SW Panel Chairman (third row, third from right); M.T. Halbouty, Chairman, Circum-Pacific Council (third row, right); E. Simpson, SCOR (fourth row, third from left); J. Corvalan, SE Panel Chairman (fourth row, second from right); G. Salas, NE Panel (fourth row, right); J.A. Reinemund, Map Project General Chairman (fifth row, second from left); K.M. Dawson, NE Panel (top row, fourth from right).

Energy Bridges to the Future

AAPG 66th Annual Convention, San Francisco, U.S.A.
May 31 - June 3, 1981

Global energy needs, developments in energy minerals, new exploration directions - these were the major concerns discussed at the Annual Meeting of the American Association of Petroleum Geologists (AAPG). A key session was devoted to the search for "subtle petroleum traps" formed by facies variations, unconformities or paleogeomorphic features. M.T. Halbouty set the tone for the session by stressing that these features had not been sufficiently studied previously because of the lack of tools to study them effectively at depth. Concepts developed in the last few years have made such study more acceptable to exploration managers, and he contended that greater effort by explorationists in seeking traps could offset not only the annually declining number of fields being found, but also the economic worth of the fields themselves.

Evidence was presented by D.L. Baars and G.M. Stevenson that reservoir reefs of algal bioherms of Pennsylvanian age grew over subtle basement controlled paleostructures. Although present-day seismic techniques cannot locate these thin reservoirs themselves, seismic mapping of the paleostructures can detect locations where these reefs may occur. E. Briseid, D.I. Milton and O. Skarpnes summarized the search for the Statvik field in the Norwegian sector of the North Sea. During early exploration, the crucial problem was whether the Jurassic reservoir rocks had been eroded during a late Kimmerian event. Although early seismic work showed

no reflectors below the unconformity, a later seismic reflection survey shot in 1974 indicated the presence of a thick underlying Jurassic sequence, confirmed by drilling in 1978. This was followed in 1979 by a program of 3-D seismic surveying which was interpreted as drilling proceeded and greatly speeded up evaluation of the field. Before drilling, the probable recoverable reserves were calculated to be 1.6 billion barrels; following drilling, the estimate was revised slightly downward to 1.2 billion barrels.

Two papers focussed on how strata older than the source rocks acted as host for petroleum derived from much younger strata. For example, H.J. Watson described the Casablanca field offshore Spain, where the reservoir rock is weathered and fractured Upper Jurassic carbonate rock the porosity of which was greatly increased during weathering in the early Tertiary. A combination of the secondary porosity and the extensive fracture network converted the dense carbonate rock into a commercially exploitable reservoir. The carbonate reservoir forms a ridge bounded by Miocene faults and capped by Miocene and younger shales. The oil source is the Middle Miocene organic shales.

G. Zhai referred to various types of so-called "buried-hill-type" oil fields in the northern part of China. In certain places the faulting has given rise to horsts and grabens in what appears to be a compressional environment. In one