Conference Reports . . .

Upper Cretaceous Standard Sections and Biozonation

3rd Working Session, Working Group on the Coniacian-Maastrichtian Stages
March 20 - 29, 1981, Tremp, Spain

The Working Group on the Coniacian-Maastrichtian Stages (WGCM) is part of the IUGS Subcommittee on Cretaceous Stratigraphy with Regional Working Groups for different parts of Europe (Fig. 1). Each group attempts to establish regional standard sections using the vertical range of fossils in order to define significant stratigraphic boundaries.

Figure 1. Location of the Regional Working Groups of the WGCM: a) SW France, b) Paris Basin, c) Netherlands, d) NW Germany, e) Denmark, f) SE England, g) N Ireland, h) S Germany, i) Austria, j) Switzerland, k) N Italy, l) N Spain.

The 3rd field working session of the WGCM focussed on the Upper Cretaceous of the Montsec, Flamisell/Pallaresa, and the Basque-Cantabric regions of Spain. The Sierra del Montsec is a 40 km long range of well exposed Mesozoic strata overthrust onto the Cenozoic strata of the Ager syncline in the southern region of the Tremp Basin. The Lower Cretaceous here is much reduced, discontinuous, with interlayered zones of continental (with erosional surfaces) and marine facies. The Upper Cretaceous marine transgression is represented by marls and limestones of Cenomanian and Turonian age. Overlying limestones and nodular marls belong to the Coniacian and Santonian. During the Campanian a carbonate platform developed which displays sporadic deposition of terrigenous material increasing upwards into the Maastrichtian continental sediments.

The oldest Cretaceous sediments in the northern part of the Tremp basin (Pradavnit) are dark micritic, thick-bedded Aptian limestones whose base is in fault contact with Triassic strata. Overlying Albian marls grade into shallow water, neritic Cenomanian limestone. Turonian marls in turn grade upwards into a Coniacian reef complex followed by evidence of basin deepening in the Santonian where the typical *Micraster* marls were deposited. In the Lower Campanian, flysch-like sediments were deposited and in the higher Puimanyons turbidites, mud flows and olistostromes are common. In higher sections the sedimentation changes to grey homogeneous marls until the lower unit of the Maastrichtian Areny Sandstone Formation records a regression comparable to that of the Montsec section culminating with

the sedimentation of continental material of the Upper Maastrichtian Tremp Formation.

Stage boundaries in the Upper Cretaceous are well-exposed in the Basque-Cantabric or Vascogotic Ranges of N Spain. For example, the disputed Turonian-Coniacian boundary appears well-defined in the Estella region and may resolve some of the previously identified stratigraphic problems. Similarly, the Coniacian, Coniacian-Santonian boundary and Lower Santonian are well exposed and fossiliferous in the W and SW Vascogotic Ranges. Sections of the Upper Santonian and Campanian are best represented in the Victoria plain, the Barranca of the Rio Araguil or in the Estella basin, while the well exposed Upper Campanian, Maastrichtian and the Cretaceous-Tertiary boundary are well defined in the flysch trough.

Part way through the field study, a two-day seminar was held at Tremp. Each speaker described critical stratigraphic information (mainly from classic localities) required for the broader regional correlation.

M. Feist (F.R.G.) and F. Colombo (Spain) reported on non-marine Maastrichtian Charophytes at Tremp which occur between marine intercalations. In particular, the presence of *Peckichara sertulata* from among the ten species represented confirms the position of the Rognacian within the Lower Maastrichtian. These new collections will greatly assist in locating precisely the Cretaceous-Tertiary boundary in the non-marine formations.

R. Höfling (F.R.G.) compared benthic foraminiferal assemblages from the Senonian Gosau Beds of the Eastern Alps with megafossil assemblages and sedimentological data to derive information about ancient marine environments - bathymetry, biotopes, and water energy in sections consisting of well-bedded silty and sandy micaceous marls intercalated with hard carbonate-cemented fine-grained sandstones. In the Moosenalm section (Bavaria) the plankton/benthos ratio ranges from 1:5 to 1:25 in its upper part, which indicates an outer-shelf fauna (ranging in depth from 100/200m to 50/100m).

H. Summersberger and H.A. Kollman (Austria) recounted how the finding of *Texanites* in the Grabenbach Formation and in the underlying Streieck Formation required the sequence to be re-defined as Santonian. Here, the ammonite and gastropod fauna is of great importance for the paleobiogeography of the Upper Cretaceous: affinities to faunas of the Circum-Pacific realm, as well as to Westphalia are evident. The

Figure 2. Lunch break in the field during the Montsec excursion.

CONFERENCE REPORTS....

gastropods give a hint of a temperate climate in the Upper Santonian of the Alpine Gosau basins.

F. Proto-Decima (Italy) presented the Coniacian to Maastricht stratigraphy of some pelagic limestone sections of the Venetian Alps (Northern Italy) which were investigated by F. Massari, F. Medizza (Italy) and J.E.T. Channell (Switzerland) who studied the microfossil content of three Upper Cretaceous sections in the Scaglia Rossa Formation. The upper part of the Scaglia Rossa is characterized by stratigraphic gaps, generally marked by abrupt changes in lithology and/or hardgrounds. The more important unconformities can be followed on a regional scale and are located within the Campanian and at the Cretaceous-Tertiary boundary. This information provided the basis for detailed magnetic-stratigraphic study which correlated the reversal sequences directly with planktic foraminiferal and calcareous nannofossil zones.

J.M. Pons, E. Caus, A. Cornella, J. Gallemi, E. Gili, and R. Martinez (Spain) traced the changes in paleoenvironment in the Montsec and Flamicell-Pallaresa regions of the Central Pyrenees. Limestones and nodular marls deposited during Coniacian and Santonian are overlain in the Campanian by a carbonatic platform within which there is sporadic deposition of terrigenous material; the latter increases during the Maastrichtian until the deposition (as a result of regression) of the continental material of the Tremp Formation (Maastrichtian to Paleocene).

E. Seibertz (F.R.G.) focussed on the problem of correlating the ammonite and inoceramid zones near the Turonian/Coniacian boundary. The Late Turonian/Early Coniacian divisions with inoceramids and ammonites of South and North America, U.S.S.R., Japan, Europe, North Africa, Madagascar, New Caledonia, and New Zealand were compared and uniform criteria for definition of zones and boundaries were presented for the inoceramid division. These criteria were discussed in

relation to the type locality near SE Münsterland, so that the Turonian/Coniacian boundary (in the traditional German sense) can be correlated with the international scheme.

A. Swiecicki (Scotland) described detailed foraminiferal analysis from 24 sections and boreholes in Campanian-Maastrichtian strata in S England, the North Sea, and the Fastnet Basin. The dominant planktonic fauna are characteristic of relatively shallow (300 m) shelf seas within the Transitional/-Boreal biogeoprovince. Meanwhile C.J. Wood (U.K.) demonstrated that the chalk of Yorkshire-Lincolnshire-North Norfolk has closer affinities both lithologically and faunally with correlative successions in Germany, Poland and the U.S.S.R. than with the chalk of the Southern or Anglo-Paris Province.

J. Wiedmann (F.R.G.) discussed the ammonite zonation of the Coniacian-Maastrichtian stages in N Spain. The ammonite diversity and evolutionary rate decrease in a simple manner from a maximum during the Mid-Cretaceous transgression at the Cenomanian toward the top of the Cretaceous. The youngest ammonite fauna (of dwarf size) is found just below the big faunal break at the top of the Cretaceous.

It was agreed that the 4th Working Session of the WGCM be held directly after the Munich meeting, June 7-11, 1982 and manuscripts will be published for presentation at the Subcommission's meeting, Oct. 18-21, 1983, in Copenhagen. The zonations and boundaries will be discussed then, and the discussions will also be published by the IGC, Aug. 4-14, 1984, in Moscow when it is anticipated that the Subcommission will be reorganized. Working Group members are invited to discuss correlations with the French type localities at the Colloque Sénonien, Sept. 26-30, 1983 at Marseille.

Professor Dr. F. Schmid (F.R.G.)
Convenor of 3rd Working
Session, WGCM

Dr. E. Seibertz (F.R.G.)
Secretary, WGCM

Automation of Geological Data

COGEODATA Symposium, Baku, Azerbaijan, U.S.S.R., April 6-11, 1981

As the need for mineral resources continues to rise, the problems of storage, automatic processing and retrieval of geological data are becoming increasingly more important around the world. In April, scientists from nine countries (Canada, Czechoslovakia, G.D.R., Hungary, Mexico, Norway, Poland, U.K. and U.S.S.R.) met in Baku, the capital of Azerbaijan, for a six-day symposium on these problems. Four main directions of research were emphasized: the structure of geological data, automatic processing of geological data on oil and gas deposits, automatic processing of geological data on ore deposits, and the processing of geological data obtained by remote sensing.

The wide application of modern computers and mathematical methods of data analysis during geological exploration has stimulated a number of countries to develop automated systems for geological data processing. Although the objective is to create logical structures of geological data bases to use as a foundation for developing powerful automated banks of geological data, the basic problems of ordering, formalization and standardization of data structure on defining properties, and the structure of natural and technological entities still remain unresolved. Discussion among participants did result in the following proposals: that the development of relational geological data bases will be of increasing concern in future and that logical-mathematical methods of geological data analysis - characterized by simplicity and in agreement with the traditional thinking of geologists - are becoming more necessary and widely used.

Because automatic data processing is used extensively by petroleum geologists and the oil industry, participants focused on the problem of creating automated systems of data processing capable of prognosis, surveying, exploration and exploitation of oil and gas deposits. There was a broad exchange of views on the application of different mathematical methods, algorithms and programs that have taken place. It was agreed that further development of automation in petroleum geology is connected with formulating accurate problems that take into account the information situations in different regions. As well, it was agreed that different methods of multivariate problem solutions should be developed, using mini- and micro-computers for geological data processing directly at the deposit sites, and effective man-machine systems be created for managing of geological prospecting for oil and gas.

Participants agreed that the present state of development and application of automated geological data processing systems on ore deposits is relatively advanced. Future investigations in this area should have three main objectives: creating sets of mathematical methods and algorithms, revealing the conditions for optimal applications of these methods and increasing the predictability and reliability of results for use in geological interpretations.

The COGEODATA symposium was organized by the Azerbaijan Academy of Sciences in collaboration with U.S.S.R. Academy of Sciences, the U.S.S.R. National Committee of

CONFERENCE REPORTS...

Geologists and the U.S.S.R. Ministry of Geology. Executive members of COGEODATA also met with the working group leader, Dr. S. Bie (Norway), and two representatives of the Azerbaijan Academy of Sciences, Dr. B.A. Bagirov and Dr. J. Djafarov, to discuss future COGEODATA activities.

Dr. D.A. Rodionov
Vice-Chairman
COGEODATA

Dr. R.I. Volkov
Secretary, U.S.S.R. National
Committee of Geologists

(STOP PRESS: COGEODATA member Dr. C.F. Burk, Jr. has just informed us that **Dr. Stephen C. (Binks) Robinson**, first Chairman of COGEODATA, died on September 13, 1981. The successful inauguration in January, 1967 of a formal group within IUGS to address the challenges presented by computers and new information technology in the handling to geological data was due primarily to the foresight and enthusiasm of Dr. Robinson. Having led an ad hoc committee of Canadian geologists during 1965-67 to develop the concept of a national system for geological data, Dr. Robinson recognized the impact that this new technology would have on international geology and accepted to organize and chair COGEODATA from 1967 to 1972. He retired in 1973 following a 25-year career with the Geological Survey of Canada.)

Participants in the COGEODATA Symposium included, front row (L to R): Dr. B.A. Bagirov, Institute on Problems of Deep-Seated Oil and Gas Deposits, Baku; Dr. V. Gavrishin, Polytechnical Institute, Novocherkassk; Dr. J. Bradacova, State Office for Geology, Czechoslovakia; Dr. R. Sinding-Larsen, Chairman of COGEODATA, Norwegian Institute of Technology; Dr. S. Sirotinskaya, Institute of Geology of Ore Deposits, Petrography, Mineralogy and Geochemistry, Moscow.

Crustal Movements in Africa

UNECA International Symposium, Addis Ababa, Ethiopia, May 5 - 16, 1981

How might an international plan for field operations in regions of Africa where crustal movements are active be implemented? This was the basic question underlying much of the discussion at the First International Symposium on Crustal Movements in Africa, organized by the United Nations Economic Commission for Africa (UNECA) and the IUGG/IAG Commission on Recent Crustal Movements (Africa).

Specific symposium objectives included the following:

- (a) to focus on the importance of monitoring and studying movements of the earth's crust in order to assess both the seismic risks involved in constructing large dams, buildings, and major pipelines and the effect of the deformations on geodetic and levelling networks;
- (b) to present the practical implications of research on continental drift and geological correlation, especially in interpreting the distribution and structure of mineral and hydrocarbon deposits.

The Symposium resulted in the adoption of four resolutions:

Resolution 1 on the seismicity of Africa and its impact on development planning recommends:

- (a) As a first step, a map be prepared showing all existing stations for the African and Arabian plates, together with a listing of these stations with coordinates, instruments and the international agencies to which they are currently reporting.
- (b) That up-to-date maps of earthquake epicentres and Quaternary faults be compiled, and a preliminary earthquake risk map be prepared on the basis of these two maps.
- (c) Inasmuch as the data available at present are insufficient for an accurate assessment of seismic risks, at least one simple seismic station (that is, with one short-period vertical seismometer) should be installed immediately in each country, preferably at a university or a government agency, and encouraged to report the data to the international seismological centres.
- (d) After assessment of the epicentre and fault maps, it will be possible to make recommendations concerning the siting of more sophisticated stations and the most suitable standardized equipment.
- (e) That the preliminary seismic risk map be periodically revised in the light of accumulated data.
- (f) Networks of local seismic stations should be established in connection with major constructions which are likely to be affected by earthquakes.

Resolution 2 on monitoring crustal movements and its social and economic implications noted the possible loss to human and economic resources which may accrue from the opening of fissures and cracks within the African rifts and similar phenomena, such as the breaching of the Nyiragongo lava lake in Zaire, and the need to complement allied geophysical investigations for the study of earth dynamics by geodetic work. It recommended that the geodetic study of crustal movements be taken up immediately in the African continent, especially in those countries affected by rifting, in cities affected by subsidence, and at large engineering structures.

Resolution 3 on the geodetic techniques for measuring crustal movements noted the lack of adequate geodetic networks for this purpose and recommended;

- (a) That each country immediately begin appropriate studies, using ground methods, on special geodetic networks designed on the basis of scientific and technological considerations in conjunction with seismic stations and allied geophysical measurements.
- (b) That the economic feasibility of using appropriate space techniques, when these are available, for regional studies (areas extending over 100 km) be studied.

Resolution 4 on volcanism, plate tectonics and mineral and energy resources noted the significance of plate tectonics and the study of the nature and composition of the magmatic rocks in the location of geothermal and mineral resources, including fossil fuels, and recommended the following action:

- (a) The preparation of adequate topographic maps at 1:250 000 or larger.
- (b) The establishment of a computerized data bank of geological, volcanological, geophysical, geochemical and metallogenic data.
- (c) The compilation of corresponding general thematic maps and specific metallogenic maps showing the distribution of mineralization having similar genesis, evolution and relation to the sedimentary history, metamorphism, magmatism and structural evolution.
- (d) The selection of appropriate areas in which to carry out further studies to guide exploration.
- (e) The use of the data bank for the assessment of the geothermal resources of the continent through the preparation of appropriate thematic maps and the selection of areas for detailed work.
- (f) That the attention of the African countries be drawn to

CONFERENCE REPORTS...

the importance of off-shore research on the distribution of sediments, the tectonomagmatic evolution and basin formation and the combined seismic, palaeomagnetic, radiometric and deep sea drilling programmes, as the understanding of the evolution of the palaeogeographic situation is the necessary first step toward the discovery and assessment of possible petroleum and other resources off the shelf area of Africa.

The Symposium requested UNECA to initiate and coordinate action to implement these recommendations in collaboration with African governments and the appropriate international and regional agencies, including the Geological Society of Africa, the Association of African Geological Surveys, IGCP and the International Centre for Recent Crustal Movements.

Plate Tectonics and the Proterozoic

International Proterozoic Symposium: Madison, U.S.A., May 18-21, 1981

The question of whether or not present-day plate tectonic processes can be extended back in time to the beginning of the Proterozoic is of considerable importance globally. Particularly important for the mid-continent region of North America, it constituted the main theme of a recent international symposium on the Proterozoic held in Madison, and organized by the Department of Geology and Geophysics, University of Wisconsin to celebrate Wisconsin's "Centennial of Geology".

Palaeomagneticians (such as D.J. Dunlop and J.D.A. Piper) envisage large Proterozoic continental plates or even supercontinents with little relative motion between cratonic domains separated by Proterozoic mobile belts. Some geologists (such as A. Baer, D. Bridgwater, A. Kröner, R. St. Lambert and X.-Y. Ma) try to reconcile this view with the plate tectonic scheme and develop models not requiring opening and closing of large oceans during the Proterozoic. Others (like K. Burke, P. Hoffman and B.F. Windley) argue strongly in favour of uniformitarian tectonics and see evidence of the present Wilson-cycle in many Proterozoic rock assemblages.

At the symposium, A. Kröner showed slides and a film of a late Precambrian ophiolite with sheeted dykes in Egypt, while P. Hoffman, in one of the most convincing lectures of the meeting, presented a detailed account of the evolution of the mid-Proterozoic Wopmay Orogen, NW Canada, in terms of the Wilson cycle. There remains little doubt, therefore, that, at least locally, contemporary plate tectonics were in operation as far back as about 2 Ga ago.

K. O'Nions demonstrated the usefulness of Sm-Nd isotopic systematics in Proterozoic sedimentary rocks for unravelling the still unknown mechanism of crustal growth. R. Taylor also showed that REE patterns in Proterozoic and younger sediments differ markedly from those in Archaean strata. Clearly, crustal growth mechanisms appear to have changed at the Archaean/Proterozoic boundary about 2.5 - 3 Ga ago,

Field party examines Archean greenstone east of Wakefield, Missouri. (Photo courtesy C. Craddock)

The technical program was arranged in 22 working sessions at which 48 papers were presented and discussed. In addition, panel discussions covered the following topics: (i) seismicity of Africa and its impact on development planning; (ii) geotectonic aspects of crustal movements; and (iii) volcanism, plate tectonics, and mineral resources.

Eighty-five geoscientists from 30 countries participated. Symposium proceedings will be published by UNECA. Follow-up work on the implementation of the resolutions has already begun.

A.M. Wassef
*Chairman, Commission on Recent
Crustal Movements (Africa)*

and the geochemistry of sedimentary rocks may help to clarify and explain this change.

R. Newton emphasized that Precambrian high-grade metamorphic episodes were discrete events not necessarily directly related to crustal accretion processes. The picture that emerges is of a "wave" of CO₂ driving before it H₂O and K purged from the lower crust. These processes may be due to large-scale underthrusting of continental crust.

Other topics discussed at the meeting were the evolution of life, atmosphere and oceans in the Proterozoic, as well as Proterozoic glaciations and ore deposits. The discussion on the Proterozoic geology of the Lake Superior region centred on the question of whether the Penokean foldbelt in Wisconsin originated in an intracratonic setting (P.K. Sims, Z.E. Peterman) or whether it evolved through subduction and plate collision (D.K. Larue, F.W. Camboray, W.R. Van Schmus). Since no unquestionable Archaean rocks have so far been identified in the volcanic-plutonic terrane of northern Wisconsin, the matter is still unresolved.

The symposium ended with several papers on the Late Proterozoic Keweenaw sequence and related rocks, following which a party of some 30 delegates embarked on a field trip led by C. Craddock, M.G. Mudrey, Jr., and H. Ali, to study these rocks in the Lake Superior region.

The meeting was superbly organized by L.G. Medaris and his staff and a Special Volume of the Geological Society of America on the contributions of symposium delegates is now in preparation. Of the 150 earth scientists attending the symposium, overseas delegates included participants from Europe, Australia and China. There is no doubt that well organized meetings of this nature have certain advantages over some of the more gigantic and costly international sessions that tend to inhibit productive discussion of specific scientific problems.

A. Kröner
Secretary, IUGS Commission on Tectonics

Preston Cloud addresses participants gathered in the new lecture theatre of Lewis G. Weeks Hall for Geological Sciences, University of Wisconsin.