CONFERENCE REPORTS

Coral Reefs and Sea Levels

Tahiti, 27 May – 9 June 1985

Coral reefs are rich oases of life in an otherwise impoverished tropical ocean. They constitute a fragile ecosystem, struggling to keep in tune with the rhythm of relative sea-level change produced by eustatic and tectonic movement, while continually repairing the ravages wrought by slight chemical imbalances, periodic infestations, and major climatic catastrophes. Reefs are also increasingly being subjected to anthropogenic stress, as island societies harvest their products more intensively and urban development discharges pollutants into shallow lagoons. Not surprisingly then, reefs are now the subject of intense study by scientists of a wide range of disciplines seeking to understand better the complex interactions that determine their existence. Much research is being prompted by the desire to help nature recover from the ravages of storm, starfish, and society.

The latest in a series of congresses devoted to the topic was held near Papeete, Tahiti, in the Society Islands archipelago. Sponsored by the International Association of Biological Oceanography (IABO), it followed earlier meetings in India (1969), Australia (1973), U.S.A. (1977) and Philippines (1981). With Unesco providing travel grants to young scientists from developing countries, the congress was generally supported by agencies of the French government, notably the Natural History Museum. It attracted some 550 participants from 48 countries who presented more than 400 papers and posters in six days of five concurrent sessions. Six field trips were offered, two largely geological. The general excellence of the entire program, the outstanding cultural offerings, the warm welcome by civic and senior government officials, and especially the hard work of a dedicated and innovative organizing team made this congress an unqualified success. Even a postage stamp was issued to mark the occasion!

The non-geological part of the Congress dealt primarily with ecological aspects, ranging from details of the physiology and biochemistry of organisms (coral, sponges, molluscs, fish and algae), to biotic relationships, environmental stress such as maturing lagoon hydrology, the processes of bioerosion, poisoning, calcification, diagenesis, sedimentation, and storm damage. Several sessions addressed practical aspects such as management and conservation, assessment of fishery stocks, and the socio-economics of reefs. The great variety of knowledge now being gathered from most of the world's major reef areas is providing the basis essential for the efficient protection of this unique habitat and the sound management of its resources.

But what does a polyp have to do with plate tectonics? The answer is, of course, that because reef-summit coral species strive to grow to within a few decimetres of low tide, the reef is intimately connected to global sea level. The age and elevation of fossil reefs can thus reveal crustal flexure and gravitational change. From reefs on oceanic volcanic islands, whose age and distance from spreading centres and/or hot spots is known, plate movements in mid-ocean areas can be gauged.

For this reason, P.A. Pirazzoli, leader of IGCP Project 200 on Sea Level: Correlations and Applications, organized the project's annual meeting in conjunction with the Congress.

Jointly sponsored by the Indo-Pacific group of the INQUA Shorelines Commission, this meeting consisted of two days of paper sessions and a three-day field trip to a variety of reef, atoll, and volcanic island settings. The meeting also provided a forum to review the progress of sea-level research in other parts of the world since the 1984 meeting in Argentina, and to demonstrate the practice of reef analysis for geodynamic purposes.

New determinations of crustal thickness and mantle viscosity, particularly for southern hemisphere areas, were presented by K. Lambeck (F.R.G.), using global models based on postglacial sea-level change. He stressed that Late Holocene records are the best expression of crust and mantle properties; more and better determinations should be made to define the exact trends at well-chosen localities. Lambeck's refinement of these models is now beginning to influence our understanding of the melting of ice sheets. For a 2,000-km

Figure 1: Emergent Holocene reef (foreground). Dissected remnant of a 3.2 Ma-old volcano (Ote Manu) in background. Bora Bora.

Conference Reports

segment of the Brazilian coast, L. Martin (Brazil) displayed a series of curves to 5,000 yBP that appear to show a large-scale geoidal shift. J.T. Hollin (U.S.A.) discussed the hypothesis that rising levels in the last interglacial undercut the marine-based West Antarctic Ice Sheet, caused a surge and a resulting sea-level rise of 7 m. If true, a similar event might occur in the present interglacial.

In a session devoted to reef evolution in relation to sea-level change, M. Hendry (Jamaica) touched on climatic impact by describing the 120,000 yBP raised reef in Jamaica and its cover of aeolianite formed during the subsequent glacio-eustatic regression. Papers by P.J. Davies (Australia) and by C. Neumann (U.S.A.) demonstrated that reefs do not always keep pace with rising sea level ("keep up" reefs), but may lag for a while ("catch up") or may fail ("give up"). Therefore, unless coral facies are well understood in relation to paleo-depth, sea-level curves based on reef dates may depart greatly from the actual trend. However, Pirazzoli and A. Strasser (Switzerland) showed that petrological analysis of cements and textures can reveal the former phreatic/ vadose contact ($\approx$ mean low water), so that samples for dating can be taken from the most definitive level. Superposition of reefs of present and past interglacial high sea levels on the Great Barrier Reef was described by D. Hopley (Australia) and P.A. Symonds (Australia).

Figure 2: Huahine - a dissected Quaternary volcano fringed by a modern reef flat from which a few remnants of an emergent Holocene reef project.

Sea-level changes in other areas were documented: by P.J. Hearty (U.S.A.) for the Mediterranean, based on amino-acid correlation of raised beaches; by J. Lecolle (New Caledonia) for the New Hebrides where two prominent reef levels enable tilting and faulting to be quantified, by G. Gvirtzman (Israel) for the Sinai where deposits of four transgressive maxima successively cross-cut one another, and by C.D. Woodroffe (Australia) who evaluated differential uplift of crustal blocks adjacent to a submarine trench (Cayman Island vs. Tonga Island). Of particular interest was an analysis of seismic uplift of Ryukyu Island by T. Kawana (Japan) who interpreted a delevelled sequence of marine notches to infer earthquake magnitudes of more than 7 and recurrence intervals of 8,000 years (using the equations of Iida and Matsuda). A seismic gap was also confirmed by absence of young raised shorelines.

A special seminar on "Reef Growth and Sea Level," backed by an exhaustive discussion paper by Davies and L.F. Montaggioni (Italy), presented a global review of the various criteria in reefs, which can be used to infer specific sea-level positions. They concluded that the upper 2 m of a reef body are crowded with sea-level fingerprints, and that there are

several well-preserved and easily recognized coral species and sedimentary textures, equally valid throughout the geological record, which serve to fix sea level at the time of their formation. Reefs thus contain the best record of relative sea-level changes of a few metres, over intervals ranging from 10^2 - 10^5 years. They provide the most practical approach to measuring vertical movements associated with regional tectonics, subduction, rifting and warping over the broad equatorial belt. In addition, they record the broad climatically induced swings of sea level largely attributable to northern hemisphere ice sheets, and thus mark the tempo of glaciation.

On the field excursion to the Leeward Islands, P.A. Pirazzoli demonstrated recently dated emerged reefs. He chose a group of four volcanic islands with reefs (Maupiti, Tupai, Bora Bora, and Huahine) lying to the west of Tahiti, and representing various stages of maturity. The series offers elegant testimony to the general validity of Darwin's original hypothesis of successive creation and subsidence. Together with Tahiti itself, the string of successively younger volcanoes formed as the Pacific Plate drifted westward over a hot spot at about 11 cm per year. Gradual subsidence as the plate thickened and cooled is shown by the progressively greater dissection and submergence of the more western islands, and by the absence of last interglacial raised reef that occur elsewhere in the Pacific.

Mehetia, the youngest island, located east of Tahiti, has no reef yet; Tupai, the oldest (5Ma), is an atoll with the central core completely submerged. Superimposed on the long term submergence of several hundred metres is a very recent (1-5,000 yBP) emergence of the outer islands of about 1 m. This emergence uplift is thought to be due to a slight drop in sea level, evidence of which is widespread in French Polynesia, possibly enhanced in some islands by uplift associated with a crustal bulge peripheral to an isostatically down-warping flexure under the more massive cone of Tahiti. Differences in the height and age of the emergence maximum suggest that the upwarp continues, though at a very slow rate.

From the lessons learned in field and lecture hall, the participants came away with a fund of carefully documented evidence that Late Quaternary reef stratigraphy reveals fundamental changes in hydrosphere and lithosphere over a considerable range of space and time. Together with other records, they contribute to a deeper understanding of past and present global change.

D.R. Grant
Geological Survey of Canada
401 Lebreton St.
Ottawa, Canada K1A 0E8

Macro Meso Micro

Utrecht, The Netherlands, 10-12 April 1985

Growing numbers of structural geologists feel the need - and occasionally even manage - to make the step from the specific, *ad hoc* explanations to a mechanistic understanding of their observations. They move on from faults to faulting, from Himalayas to Himalaying and from fabrics to fabrication. To keep informed of these wonderings about the kinematic and dynamics of the deforming Earth, a series of international conferences has been spontaneously organized by universities in Western Europe. A symposium in Leiden (1976) formed the first in this series, followed by Barcelona

Conference Reports

(1978), Göttingen (1980) and Zürich (1982). In April, Utrecht University welcomed 220 participants from all over the world for a three-day conference on tectonic and structural processes.

The working title "Macro Meso Micro" succeeded in bringing together specialists working on all scales, from the continental to the submicroscopic. The whole gamut of current structural interests was displayed in 35 papers and in more than 60 posters for which ample viewing time was set aside. The proceedings of the meeting will be published in a special issue of *Tectonophysics*.

The Macro-session contained a wide variety of investigations of larger structures, from the Himalayan framework (V.C. Thakur) to the role played by salt levels in reducing the cross-sectional taper of thin-skinned fold-and-thrust belts (D.M. Davis and T. Engelder). The reactivation of interest in mantled gneiss domes resulted in an interesting set of contributions. J.P. Brun and J. van den Driessche attributed the anomalous thermal regime responsible for gravitational instability of domes to large-scale crustal thrusting. The thermo-mechanical modelling of deformation histories is clearly becoming more sophisticated, to an extent that die-hard field geologists are occasionally left behind. However, the danger of theoretical models ignoring the

Figure 1: Mesoscopic folds in pegmatite dike, northeastern Newfoundland.

observations they aim to explain is not entirely imaginary. The clear-cut contributions of J. Jackson and G. Mandl on faulting were therefore well received. One may expect that structural geologists concerned with large-scale structures will increasingly come together with geophysicists in meetings like this to explore common ground.

The build-up of strain was the focus on the Meso-scale day. Well illustrated case-studies, such as that of J.M. Casas and J.A. Munoz on the Eastern Pyrenean Alpine Thrust, alternated with numerical modelling of M. Casey and a study by W. Bosworth on the seamy, anastomosing nature of deformed clayey rocks in olistostromes and at the base of superficial nappes. The Argille Scagliose of the Apennines provides a well-known European example. Small-scale brittle micro-shearing plays an important role in this type of high strain rate deformation. P. Cobbold illustrated the use of Mohr diagrams to explain kinematically the generation of fold

structures in particular types of strain regimes. The work of Lister, de Paor, Means and others has made familiar the idea that anything tensorial can be readily visualised with Mohr's famous circle. It is a powerful conceptual tool and not exclusively for the representation of stress.

An exemplary investigation of mass transfer and synmetamorphic deformation in a small fold was presented by J.M. Caron. French geologists are clearly making rapid advances in integrating geochemical, structural and experimental studies. This was a central theme on the final day of the conference devoted to microscopic processes. M.A. Etheridge ably introduced the subject, emphasizing the role of fluid in crustal deformation and metamorphism. J.-P. Gratier's review of the scales of mass-transfer circuits and their attendant rate-limiting factors admirably completed our present picture of the problem.

Fabrics and recrystallisation were the last topics. To outsiders it must be an odd sight to find, at the end of a tiring three-day conference an entire hall of geologists becoming excited by video clips, movies and sequential slides of deforming mothballs and toilet blocks (camphor and paradichlorobenzene - J. Urai, R. Knipe, J.M. Jessell), common ice (C.J.L. Wilson) and mixtures of salt and calcite (M. Jordan). However, the opportunity to see under the microscope and record on film, these analogue studies of the growth and destruction of grains and the development of fabrics during deformation has given an enormous boost to students of microstructures in metamorphic tectonites. The knowledge of the microscale mechanisms of deformation is rapidly changing because of it. The obvious relevance to the study of ceramics and metals also gives geologists an opportunity to contribute significantly to the general field of materials science.

The next conference in this series will be held in Uppsala, Sweden in 1987 on the occasion of the 70th birthday of Hans Ramberg.

I.v.d. Molen, and
H.J. Zwart
Instituut v. Aardwetenschappen
Postbus 80-021
3508 TA Utrecht, The Netherlands

Sixth Gondwana Symposium

Ohio, U.S.A., 19-23 August 1985

Meeting for the first time outside Gondwanaland, the Sixth International Gondwana Symposium recently attracted about 150 scientists, largely from Europe and North America. This was held at the Institute of Polar Studies at Ohio State University, an important center for Antarctic research. Not surprisingly nearly 40% of the papers presented were on Antarctica, indeed there was one special session on North Victoria Land, the site of recent international co-operative studies. One highlight was the first report of thrust tectonics in this region.

Contemporary Gondwana research is focused on two main research areas. The first comprises the traditional comparative studies of the Devonian to Jurassic Gondwana sequences, including their paleontology, sedimentary basins, glacial beds and volcanics, and most of the lectures were on these topics. Paleontologists still see a number of problems in correlating

Conference Reports

faunal and floral development between the different regions of Gondwana and with the adjacent continental terranes, Laurentia and Angara. Faunal links with Asia Minor and across the Himalayas into Angara raise doubts as to whether the Tethys in this area really was a deep ocean thousands of kilometres wide.

The mobile Gondwana margins constituted the other main theme. Interest in these active margins was high because of their seemingly persistent tectonic setting through several orogenic cycles, extending backwards in time from the present position in the Circum-Pacific mobile belt to about the beginning of the Phanerozoic. Gondwana break-up, reconstructions, and the intervening marine areas were the subject of another group of lectures, but the Gondwana Shield attracted comparatively little interest during the meeting. A discussion of suspected microplates in West Antarctica brought out wide differences in interpretations based on marine and continental data. Plate tectonic models for the Paleozoic Ross Orogen, the Mesozoic Gondwanide Orogen, and the Cenozoic Andean Orogen of Antarctica with its counterparts in other Gondwana areas, especially South America, were also presented.

The first discovery by T.S. Laudon (U.S.A.) of *Glossopteris* at the base of the Antarctic Peninsula (English Coast) was reported. The British Antarctic Survey displayed a copy of its new 1:3 million tectonic map of the Scotia Arc, and a set of draft maps by M. DeWit (South Africa) (1:10 million) of Gondwanaland was also exhibited.

In his address at the closing banquet, C. Craddock (U.S.A.) reviewed the "R" development of American views on Gondwanaland starting in the 1920s with Rejection of the concept followed by Reconsideration in the 1950s, Reversal in the 1960s and Refinement (1970s-). The next Gondwana meeting will be held during 1988 in Brazil.

after reports by F. Tessensohn and C. Craddock

Paleozoic Terranes in Circum-Atlantic Orogens

Nova Scotia, Canada, 27 May-2 June 1985

Tectonic analysis of orogens has evolved from theories based upon geosynclines through simple plate concepts to much more complex and realistic models that attempt to resolve the interactions of microplates and discrete terranes. Applications of these new concepts to the Appalachian-Caledonide-Hercynide-Mauritanide Orogen are in the initial stages, with many differing definitions of basic tectonic units and a spectrum of conflicting terrane maps. A Penrose Conference on these circum-Atlantic orogens was held recently and served as the organizational meeting for IGCP Project 233.

The conference had several objectives. First was the need to clarify the basic tectonic units (terrane) that comprise the orogen, and to prepare a preliminary terrane map at 1:5,000,000. Constraints are required on the times of terrane amalgamation and their accretion to the various continental units that border the orogen, and pre-, syn-, and post-

accretionary movements that have complicated terrane boundaries both from a geologic and paleomagnetic perspective must be defined. The conference also aimed to interpret tectonic elements within each terrane and their overstepping sequences in the context of the genesis of the entire orogen. A final goal was to outline those areas where data are presently lacking and which should be targeted for future collaborative research projects.

With 71 participants from 13 countries the Conference began with a session dealing with terranes as defined in the Cordillera and circum-Pacific region. By the end of the meeting it was generally agreed that the "western" terminology was also appropriate for the circum-Atlantic but that there had been varying degrees of misuse of the term "terrane" in the circum-Atlantic area. Several fundamental questions were posed. What rock properties distinguish normal, oblique and transcurrent accretion? What controls the presence or absence and nature of the plutonism and metamorphism? Is it possible to estimate the thickness of continental crust being accreted and if so how? What criteria can be used to distinguish pre- and post-accretionary movements?

Figure 1: Conference participants holding down a suspicious terrane.

B. Black (U.K.) demonstrated the application of sedimentology and stratigraphy to terrane analysis, with examples from the Himalayas and the Midland Valley of Scotland. Upward-coarsening sediments characterize the Siwalik trough in the Himalayas and record the emplacement of thrust sheets. On the other hand, alluvial fans in Scotland all fine upwards, and the Midland Valley basins contain fragments whose source cannot be found in either the Southern Uplands or the Dalradians. An accretionary mechanism involving both strike-slip faulting and thrusting was thus proposed.

D.L. Bruton (Norway) showed that faunal province boundaries record paleotemperature boundaries that do not necessarily coincide with terrane boundaries. He provided a classic example in one continuous stratigraphic section where the fauna alternated between Baltic and North American as the water depth changed from deep to shallow respectively. P.G. Andreasson (Sweden) illustrated how contrasting P/T ratios deviating P-T time paths and different tectono-metamorphic histories may be used to distinguish terranes and to identify pre- and post-accretionary metamorphism.

The use of paleomagnetism was outlined by P. Lapointe (Canada) who showed how initial paleogeography could be constructed and the time of accretion identified. However, paleomagnetic distinctions are presently limited to latitudinal movements greater than 5° or ~500 km, and there are no longitudinal constraints. Complications may arise where rock and magnetization ages are not synchronous, for example where magnetic overprints occur or where minerals with

Conference Reports

different Curie Points contribute to the observed magnetization. J.P. Lefort (France) demonstrated that characteristic magnetic and gravity signatures associated with some terranes could be used in tracing them beneath cover rocks or beneath the sea. However, caution was advised because of several examples where changes within a single terrane in magnetic or gravity patterns result from metamorphic gradients or plutonic activity. Moreover, long wavelength anomalies relate to deep structures and may not mark terrane boundaries at the surface.

In a review of terranes in circum-Atlantic Paleozoic Orogens, R. A. Gayer (U.K.) showed that Greenland consists of a single terrane with overthrust miogeocline (and basement) in the East and North Greenland fold belts. In contrast, D.G. Gee (Sweden) graphically illustrated that Svalbard consists of at least three units brought together by both thrusting during accretion and subsequent dispersal by strike-slip faulting. A masterful overview of the complexity of the terranes in the upper allochthons of the Scandinavian Caledonides by M. Stephens (Sweden) demonstrated that most terrane boundaries are thrusts of both Finnmarkian and Scandian origin.

For Colombia, H. Duque-Caro presented a terrane map made in collaboration with the U.S.G.S. While most of the terranes here are Tertiary, those of Paleozoic age sparked considerable interest, and T. Anderson (U.S.A.) highlighted the Paleozoic fragments isolated in the Mesozoic terrane complex of north-west South America and Central America. Finally R.G. Hatcher (U.S.A.) unveiled a new terrane map for the Appalachians based upon structural and geophysical maps. The discussion that followed showed that many differences in interpretation result from inconsistent use of the terrane terminology.

The conference provided an exciting start to IGCP Project 233, and plans call for international collaboration in joint research and sharing of laboratory facilities, with the project serving as both an umbrella and a central clearing house. The participation from many countries has helped to launch the new project quickly, thanks to support from IGCP, IUGS, the U.S. National Science Foundation, the Canadian Geological Foundation and the Geological Society of America.

J.D. Keppie
Dept. of Mines and Energy
P.O. Box 1087
Halifax, N.S., Canada B3J 2X1

Brines and Gases in Crystalline Rocks

Fredericton, Canada, 13-15 May 1985

Saline groundwaters, brines and gases are ubiquitous throughout the Canadian Shield and are usually found at depth exceeding 1,000 m. This has long been known to the mining community and was first recorded in the Annual Report of the Geological and Natural History Survey of Canada for 1887/88. "... A curious feature of the vein is the combustible gas which has been met with in large quantities in the workings. This ... is accompanied by water containing

calcium chloride solution Two gallons of this water furnished, roughly measured, nearly a pint of very acrid and deliquescent salt, chloride of calcium.

"The gas and water are principally confined in large vugs or cavities in the vein under great pressure It (the gas) is met with in the deepest workings ... above the 8th level all water infiltrating into the mine is pure lake water, whilst below that level is a zone of highly mineralized water" Since this report, these fluids and gases have received little scientific attention, and it is only in recent years that comprehensive geochemical and isotopic studies have been initiated in response to deep waste disposal programs.

During a recent survey of mines in the Canadian Shield, the general observations quoted above were confirmed, and it was determined that at depths exceeding 500 m the occurrence of Ca-Na-Cl brines has to be expected. They are not confined to ore deposits or to any specific rock type, although host rock chemistry influences minor element abundances in these brines. Total dissolved solids can exceed 300 g/l. Brines and methane discharge from exploration boreholes that encounter open fractures, shear zones or simple vugs ("mega fluid-inclusions"), with flows ranging from a mere trickle to more than 350 l/h and lasting from a few hours to many years or even decades. Gas occurs in varying amounts.

So as to open a broader discussion on the genesis of these fluids and gases and to consider their relationship to the host rocks in which they occur today, an international workshop

Figure 1: A 3-inch borehole at 1,600 m depth in the Lockerby Mine, Sudbury Basin, Canada, discharging methane gas (1), and Ca-Na-Cl brine (2). Gypsum (3) is precipitating from the brine.

Episodes

EARTH SCIENTISTS must contend with rapidly advancing knowledge and an ever-growing information base. They must also keep pace with a bewildering diversity of organizations and a wide array of conferences, seminars and workshops.

EPISODES helps solve these problems by covering both scientific and organizational news. Feature articles - contributed by leading geoscientists from all parts of the world - review developments at the frontiers of the earth sciences.

EVERY issue includes reports on new developments in international earth science. Other standard features (e.g. book reviews, training courses, coming events) emphasize the resource aspect of the newsmagazine.

IF you are interested in what is going on globally in the geosciences, you should be reading Episodes on a regular basis.

Global Standards in Stratigraphy
Forecasting Volcanic Eruptions

The Kola Super-Deep Drill Hole
Chinese Loess & Paleoclimates

Precambrian Paleosols
The Origin of Tethys

Please enter my subscription to Episodes
\$18 (US)/year
(1986 subscription rate)

Name: _____

Address: _____

City: _____ Province: _____

Country: _____ Postal Code: _____

Mastercard _____ & Visa _____

Expiry Date _____

Signature _____

Note: Cheques or money orders must be drawn on a North American bank and made payable to Episodes, Room 177, 601 Booth St., Ottawa, Ontario, CANADA K1A 0E8.

Your Window to the World

Kail's Geophysical PROJECTOR

Enlarge or Reduce opaque images by automatic focusing

The Model K-A5 Reflecting Projector, ideal for mapping, aerial photography, drawings, layouts and seismic studies. Projector enclosed to form a table top for ease in large material handling. Images are transposed from one glass plane to the other without head or hand interference.

- Image in focus in entire scale range
- Scale from 5x to 1/5x
- Reflects opaque materials to translucent papers
- Single automatic focus remote control
- Fine focus control knob
- Dimensions 80" long, 30" wide, 36" high

Philip B. Kail Associates, Inc.
1010 West 1st Avenue
Denver, Colorado 80223
(303) 623-5987

Conference Reports

and symposium were held recently during the 1985 annual meeting of the Geological and Mineralogical Associations of Canada. Known occurrences and their geochemical and isotopic characteristics were summarized. The participation of researchers from seven European countries, the U.S.S.R., the U.S.A. and Canada was sponsored by the International Association of Hydrological Sciences, the International Association of Hydrogeology, and Atomic Energy of Canada Limited (AECL).

The largest body of information on deep brines is available from the Canadian Shield. An overview of their occurrence was presented by S.K. Frape (Canada), followed by more detailed, site-specific information on the underground research laboratory of AECL in Manitoba (M. Gascoyne) and the Chibougamau area in Quebec (J. Guha). The chemistry of these fluids is unique and distinguishes crystalline rock brines clearly from most formation waters in sedimentary basins. It was noted (I. Vouk, U.S.S.R.) that the super-deep boreholes presently being drilled in the U.S.S.R. encountered similar brines to depth of at least 12 km.

More characteristic is, however, their isotopic composition, even though this was only available for the Canadian samples. Participants were, however, told that isotope analyses will be forthcoming from the U.S.S.R. program; the preliminary data appear to agree with the Canadian results. These show ^{18}O depletion and/or deuterium enrichment with respect to normal meteoric waters, they too are very different from sedimentary basin brines, which typically are characterized by significant ^{18}O enrichments (E. Roedder, U.S.A.).

The genesis of saline groundwater and low temperature rock-water interactions was addressed in a number of presentations. Of special interest was the Swiss deep drilling program (W. Balderer, Switzerland; A. Pekdeger, F.R.G.) and a presentation by M. Edmunds (U.K.) on the genesis of water containing about 12 g/l chloride in the Carnmenellis granite of southwestern England. This initiated a discussion on the importance of salt released from fluid inclusions that are believed to contain metamorphic/ magmatic brines. The general feeling, based on the data available, was that such release was insignificant, though K. Nordstrom (U.S.A.) made a strong case that saline waters in the Stripa granite (Sweden) may well have been generated in this manner. However, P. Lampen (Finland) emphasized that fossil and modern "Baltic" salt water intrusions have to be considered throughout Fennoscandia, even though salt waters have been found in Finland that are unrelated to fossil sea water and must be explained by rock-water interactions.

Further substantiating evidence for the importance of rock-water interaction was provided by $^{37}\text{Cl}/^{35}\text{Cl}$ data on Canadian Shield brines (R. Kaufman, U.S.A.), $^{87}\text{Sr}/^{86}\text{Sr}$ comparison of rock and waters (R. McNutt, Canada), the generation of ^{36}Cl in the subsurface in crystalline rocks (M. Fontes, France; E. Murphy, U.S.A.), rare gas analyses from Sweden, Switzerland and Canada (J. Andrews, U.K.) and consideration of the carbonate system (J. Kramer, Canada). Attempts to date minerals in fractures in the crystalline rocks at Stripa where salt waters are found are now in progress (B. Fritz, France), and efforts are being made to relate water chemistry to those of fracture minerals. Model calculations (J.F. Pearson, U.S.A.) and the BRINEQ brine evolution model (B.F. Jones, U.S.A.) led to the conclusion that rock-water interactions at relatively low temperature would have to be called upon to explain most of these saline fluid occurrences.

In all cases surface evaporation and resulting salt enrichments were excluded as mechanisms for generating brines and salt water. However, fossil salt deposits such as those

generated by playa lakes can be locally important. This was convincingly demonstrated by T. Paces (CSSR) who presented data on Na-Cl- SO_4 water for the Bohemian massif.

Not fully explained were the peculiar ^{18}O contents of the Canadian brines, although a consensus existed that long-term, low temperature exchange with the host rock minerals could be the cause. Vouk attracted exceptional interest with his argument that radiolysis in closed systems over geological time spans could decompose enough water molecules to bring about not only the observed salt but also the deuterium enrichments seen in these brines.

The discussion of gases in crystalline rocks focussed primarily on the occurrence of methane across the Canadian Shield in many operating mines (P. Fritz, Canada). Isotope data (^2H and ^{13}C) show that each sampling area (usually a distinct mining district) has a unique composition. Thus, a relationship between rock evolution and gas composition is suspected. However, T. Gold (U.S.A.) eloquently defended the hypothesis that many natural gas accumulations have a mantle origin. If the Shield gases are considered in this context, observed isotopic differences would have to be explained by secondary mechanisms such as biological consumption, oxidation or diffusion. Supporting evidence for the mantle hypothesis was seen by some in the data presented by J. Welhan (Canada) on the occurrence of methane in Pacific rift areas; there is also the reported presence of methane even at great depth in the super deep boreholes of the U.S.S.R. Sampling procedures for downhole sampling (P. Wickberg, Sweden) concluded the formal presentations.

The proceedings of this conference will be published in a special volume of the Geological Association of Canada in 1986.

P. Fritz and S.K. Frape
Department of Earth Sciences
University of Waterloo
Waterloo, Ontario, Canada N2L 3G1

Conodonts: Their Paleobiology and Applications

Nottingham, U.K., 25-29 July 1985

The Fourth European Conodont Symposium (ECOS IV) attracted over one hundred scientists from thirty-one countries who came to discuss conodonts, the microscopic hard parts of a group of organisms of uncertain biologic affinity. Conodonts are composed of calcium phosphate (apatite) and are thus readily extracted from rocks using acids; because of their mineralogy and structure they are resistant and can be recovered from strata of low to medium grade metamorphism. Each conodont animal possessed several individual conodont elements of varying morphology, thus those conodonts obtained by acid dissolution are disarticulated. Two main themes were addressed at the conference: the paleobiology of conodonts, and investigative techniques used in their study and applications of the results.

Consideration of their paleobiology has recently been stimulated by the discovery of conodont elements associated with the remains of soft parts (Briggs et al., 1983; Lethaia, v. 16,

Conference Reports

p. 1-14). The organism, generally accepted as a true conodont animal, was discovered among museum collections from the Lower Carboniferous of the Edinburgh district, Scotland. The animal is small and elongate (40 mm long, 1.8 mm wide) with conodonts preserved at the anterior end. R.J. Aldridge (U.K.), the ECOS IV convener and one of the authors of the original paper reporting on the new conodont animal, opened the session on paleobiology with an announcement of the discovery of two additional specimens that contributed further to the story. He and his coauthors described the new material and placed it in the context of known Carboniferous natural assemblages. They concluded that the conodont animal (in the Carboniferous at least) was laterally flattened in life and that the features displayed by the specimens suggest a closer relationship to chordates than to chaetognaths (an extant group known as arrow worms.).

Figure 1: Searching for the elusive conodont!

Researchers who have been working with fused clusters of conodonts from the Lower Paleozoic, notably H. Szaniawski and J. Dzik from Poland, continued to suggest that conodonts compare well, at least functionally, with the grasping structure of chaetognaths. A lively discussion on the affinities of conodonts took place inside the lecture hall and in the laboratory where the recently discovered conodont animals were on display. A large model of a Carboniferous conodont apparatus constructed by workers at the University of Nottingham was also on view in the laboratory and provided a useful focus for the discussion.

Although there was no specific agreement on the nature of the conodont animal, there was broad consensus on some issues. Early conodont apparatuses, particularly proto-conodonts, paraconodonts and some simple cone euconodonts bear a functional analogy, although not necessarily a homology, to chaetognaths. Carboniferous conodonts, and probably those of other ages, may have been related to the chordates but represent a separate phylum. Early conodonts and other simple cone apparatuses probably had a grasping function; later conodonts with more complex features probably had a grasping and chewing function. In other words, conodont apparatuses were food processors in a macrophagous, probably predatory animal.

One of the highlights of the session on conodont paleobiology was the report by L.E. Fahraeus (Canada) of the isolation of soft tissue from Silurian conodonts. The tissue is characterized by cells, fibrous material and lumina that bear no similarity to known tissue types. This decalcified tissue may

represent the oldest tissue obtained from the geological record, and future studies may provide some clues to the biological affinity of conodonts.

The second thematic session focussed on investigative techniques and applications. Traditionally, conodonts have been obtained by the dissolution of carbonate rocks using organic acids, but recently hydrofluoric acid has been used to dissolve cherts. M.J. Orchard (Canada) presented extensive new results obtained from the Cordilleran orogen by routine dissolution of cherts from previously uninvestigated terranes. Other processing techniques were presented in the quest for a safer alternative to the use of toxic halogen-based heavy liquids. A demonstration of an interfacial technique, based on principles similar to those used in ore flotation, was presented by G. Merrill (U.S.A.) and E. Freeman (U.K.), but many workers remain sceptical about its effectiveness. Some discussion of the newly developed, reputedly non-toxic heavy liquid called sodium polytungstate concluded that tests of its efficiency were still required before it could be considered as a replacement for existing heavy liquids. Regrettably, no single technique is yet available to replace the traditional heavy liquid method.

Conodonts have recently been the subject of image analysis with a view to the quantification of outline shape so that a large number of elements can be compared objectively. T. Carr (U.S.A.) and G. Klapper (U.S.A.) indicated that such studies have provided insights into the variation of shape of conodonts with respect to size and environmental influences and have aided in the recognition of permanent evolutionary changes. Studies of Devonian and Triassic conodonts show that some taxonomy based only on visual discrimination has resulted in inconsistent identification. Image analysis of conodonts is in its infancy, however, and the cost of the necessary equipment is still very high, but clearly there is a future for techniques that make observations of shape more objective.

The fact that conodonts change colour with temperature has been known for some time and a number of regional studies have been completed. Detailed studies of regional patterns of conodont colour alteration indices (CAI) were presented for eastern Canada by G.S. Nowlan (Canada) and northern Hungary by S. Kovacs (Hungary). In the former study, some thermal maturity patterns were related to the passage of Mesozoic hotspots, while in the latter, valuable comparisons were drawn between carbonate texture, illite crystallinity, vitrinite reflectance and conodont CAI values. The value of conodonts as indicators of thermal history has prompted many new avenues of research for conodont workers.

One of the highlights of the investigative techniques session was the report from J. Wright (U.S.A.) and colleagues on the chemostratigraphy of conodonts across the Cambro-Ordovician boundary. She demonstrated that patterns in the secular variation of trace elements in conodont apatite permit correlation of strata across the boundary from many parts of the world. It was shown that the trace-element composition of conodont apatite records perturbations of seawater chemistry. This obviously has tremendous potential for further research along many different lines.

A talk by J. Kovach (U.S.A.) on strontium, neodymium, and oxygen isotope patterns in conodont apatite spurred great interest. He showed that the concentration of strontium in conodont apatite decreases with time and that strontium isotope ratios fluctuate through time, presumably reflecting isotopic ratios of seawater. These exciting studies will surely mean that conodont workers everywhere will be teaming up with isotope geochemists on what are likely to be significant and productive new projects. The results may permit comparison of the history of seawater chemistry, both within and between ocean basins in order to identify oceanic events of both local and global significance. It may prove possible to identify ancient seaways and trace the interrelationships of ocean basins as well as to assess the relationship between seawater chemistry and conodont distribution.

Conference Reports

Paleontological meetings, particularly those of specialist workers, typically begin with a talk on fossils from the earliest part of the group's history and end with a talk on fossils that existed just prior to the extinction of the group. ECOS IV broke that stratigraphic mould very decisively and revealed the immense diversity of studies being undertaken on conodonts. It was an exciting meeting, full of promise for the future of conodont studies and, indeed, of paleontology as a whole.

Godfrey S. Nowlan
Geological Survey of Canada
601 Booth Street
Ottawa, Ontario, Canada K1A 0E8

Clay Minerals Reviewed

Colorado, U.S.A., 28 July-2 August 1985

More than 500 clay specialists from 37 countries gathered recently in the western U.S.A. for the Eighth International Conference of the Association Internationale pour l'Etude des Argiles (AIPEA), held jointly with the annual meeting of the Clay Minerals Society. Research themes discussed ranged widely from crystalline to amorphous material and from surface chemistry to solid mechanics, reflecting the major objective of AIPEA, to promote international cooperation in clay research and technology.

The technical program was highlighted by the first G.W. Brindley Award lecture, which was presented by J.J. Fripiat (France) on the "Internal surface of clays and constrained chemical reactions." A second plenary lecture by P.W. Low (U.S.A.) on "The clay-water interface" emphasized physical accounts of the clay-water interactions. Recently developed techniques for identifying mixed-layer minerals and their bearing on crystal chemistry and diagenesis were then reviewed by V.A. Drits (U.S.S.R.).

A topic of special interest to many workers was that of the Fe-Al oxides and their relationship to amorphous material. It is now known that the presence of such material can drastically change the physical and chemical nature of soils, sediments and rocks, however, its study has been neglected because of the considerable difficulties involved.

Several papers showed usefulness of Mössbauer spectroscopy for the study of samples containing Fe, provided that effects of Al substituting for Fe and crystallite size are taken into account. Selective chemical dissolution techniques appear to be effective for removal of amorphous material and may provide useful information from which its composition could be inferred. The importance of surface reactivity in the understanding of the specific adsorption-desorption characteristics of a system containing amorphous material was also emphasized.

Four years ago, AIPEA established the W.F. Bradley Award in order to encourage young scientists to attend AIPEA conferences. C.P. Herrero (Spain) was chosen as the first recipient and presented a paper on the analysis of Si-Al distribution in micas, using Solids Magic Angle Spinning -

Nuclear Magnetic Resonance spectroscopy. He demonstrated that the tetrahedral Si-Al distribution is mainly controlled by requirements for homogeneous distribution of charge to minimize electrostatic energy and by local balance of inter-layer charges. He concluded that, except for margarite, no evidence for long-range order in the Si-Al distribution was found. This new approach appears to be a very promising alternative to the X-ray diffraction method.

The proceedings of this very stimulating and informative conference will be published. For information contact: International Clay Conference, c/o Western Experience Ltd., 2450 Central Avenue, Suite P2, Boulder, Colorado 80302, U.S.A. The next international clay conference is now being planned for Strasbourg, France in 1989.

H. Kodama,
Mineralogy Section,
Chemistry and Biology Research Institute,
Agriculture Canada, Ottawa K1A 0C6, Canada

Oceanic Fracture Zones

London, U.K., 1-2 May 1985

Fracture zones are among the most conspicuous structures of the ocean floors. Formed where offsets occur in the oceanic spreading centres, they leave scars that stretch thousands of kilometres across entire ocean basins. In the 1960s and 1970s these zones were mapped extensively by magnetic and bathymetric surveys and were used frequently in global reconstructions, but only in the past five years or so have they been studied intensively in their own right.

A sampling of the huge amount of contemporary work on oceanic fracture zones was presented at a conference held recently at the Geological Society of London, with support from the Royal Society. The meeting made clear that the use of modern high resolution geophysical tools is at last narrowing the gap between observations from remote measurements and those of field geologists. There is now hope that we may soon be able to make geological maps of the ocean floor just as we can on land, even if the closer you look the more complicated things become. Much can be learned about the tectonics of young oceanic features by simply looking at the three-dimensional bathymetry and "grain" of the seafloor.

Speakers from the U.S.A. and the U.K. presented new and quite remarkably detailed surveys across several fracture zones in the Atlantic and Pacific Oceans. Among other techniques used were SEABEAM (a multibeam swath echo sounder) and GLORIA (Geologic Long Range Inclined Asdic),

which is particularly good for detecting seafloor lineations and surface irregularities. It is now possible to locate the exact position of transform faults on the seafloor, often to within a few tens of metres. Using a near bottom seismic profiler, A.M. Bowen (U.K.) obtained high resolution seismic reflection profiles across the Vema transform that show the fault marking the plate boundary between South America and Africa cuts vertically through the sediments like a knife-cut through cake. Bowen demonstrated that over, at least, the past million years the transform fault has been confined to a zone only a few hundred metres wide.

New discoveries in the seismic structure of crust formed in the vicinity of fracture zones were also discussed. Although the seismic structure of normal oceanic crust is remarkably uniform throughout the world, recent studies using ocean bottom seismometers have shown that fracture zone crust is always highly anomalous and usually thin. A series of papers by British and American workers demonstrated that many of the anomalous features are common to all the fracture zones studied. However, nearly all the studies to date have been in the slow-spreading North Atlantic, and similarly good data from faster spreading ridges like those in the Pacific are still not available. Shipboard gravity measurements suggest that the oceanic crust thins systematically along strike towards these zones. The deeper structure beneath the lithosphere was described by C. Craig (U.K.) using SEASAT altimetry data across prominent Pacific fracture zones.

Upper mantle rocks are often found in the vicinity of transform faults. The geology of presumed fossil fracture zones in the ophiolites from Cyprus and Newfoundland was summarized by B.J. Murton and I.G. Gass (U.K.) and by J.A. Karson and D. Elthon (U.S.A.). If these ophiolite complexes are representative of modern oceanic settings, then the geology of fracture zones is indeed much more complex and variable than current models indicate.

The many different models of fracture zones presented covered a wide range of problems starting with why these zones should form at particular locations when the oceans first split. Discussions of the evolution of fracture zones on a large scale were continued by B. Collette (Netherlands) and J.A. Schouten and K.D. Klitgord (U.S.A.), using the pattern and history of seafloor spreading in the North Atlantic derived from hundreds of thousands of kilometres of magnetic and bathymetric profiles. Numerous ideas were discussed to explain more local tectonic and geological features of the crucial spreading ridge - transform intersection. The most artistic was undoubtedly J.A. Karson's highly coloured "art nouveau" mosaic to illustrate his concept of the way in which the crust is "chopped up" and mixed in the region of the transform faults.

The breadth of thinking and the voluminous new high quality data presented demonstrate that the study of fracture zones remains an exciting and developing field. The conference proceedings will be published by the Geological Society of London.

Robert S. White
Dept. of Earth Sciences
Cambridge University
Madingley Road, Cambridge
U.K.

Mississauga, Canada, 4-7 June 1985

Mafic dike swarms, which have invaded the lithosphere for more than 3 billion years, provide spectacular evidence of regional tensional tectonics. They are the most important avenue for the transfer of basaltic magma from mantle to upper crust. On continents they occur with variable ages and trends over areas of hundred of kms². In oceanic areas they play a central role in the evolution of ridge crests, and they are believed to underlie the ocean floor as sheeted dike complexes. A global view of dike patterns and ages was provided at a recent international conference at the University of Toronto, which was attended by more than 130 scientists from 20 countries.

A special feature of the meeting was a display of compilation maps from virtually every major Precambrian Shield. These showed that all are crossed by dike swarms, in some cases comprising up to five or six sets of distinct age and trend. Many of these are known only from aeromagnetic surveys or reconnaissance mapping, and information such as absolute age, geochemistry and paleomagnetic direction is lacking. Nevertheless, several maps showed major swarms whose full extent or even existence had not previously been recognized. These include Proterozoic dikes that cross the Zimbabwean craton (J. Wilson, Zimbabwe), an extensive series of fractures east of and parallel to the Great Dyke filled locally with gabbro (D. Jones, Zimbabwe), and a N-S swarm with at least two ages of dikes that extend for 1500 km through Zambia and Tanzania (H. Halls, Canada). A 1.6 Ga-old swarm several hundreds of km in length that parallels the western coast of Greenland (T. Nielsen, Denmark) was also reported, and another one around 1.1 Ga old that trends NW-SE to WNW-ESE for some 1000 kms in south-central Australia (D. Tucker and J. Parker, Australia). The initial impression from the maps was that there were certain peaks in global dike intrusion centred around 2.9, 2.5, 2.0, 1.6, 1.1, 0.7, and 0.1 Ga ago.

As regards the mechanism of dike emplacement, the consensus, on both theoretical grounds and with reference to specific regional swarms, was that flow of material was subhorizontal rather than predominantly vertical during intrusion. H. Sigurdsson (U.S.A.) showed that modern dikes in Iceland are injected laterally for distances up to 60 km from a central magma chamber located beneath Krafla volcano, and similar observations have been recorded from Hawaii and modelled using gelatin (R. Fiske, U.S.A.). It seems that surface topography plays an important role in both instances. In Iceland the hydraulic head of magma cannot rise to cause an eruption at the volcano summit, because the increased magma pressure required exceeds the tensile strength of fissures along the associated rift. In Hawaii, the magma propagates along the axial plane of a topographic ridge that is not necessarily a zone of active crustal distension.

From centrally located plugs of Tertiary age in Greenland, the U.K. and the Canary Islands, dike swarms extend several tens of kms, suggesting that the plugs are deeply eroded equivalents of the feeder chamber below Krafla. In contrast, Jurassic swarms from eastern North America, Africa and South America are not spatially related to intrusive complexes that may be recognized as feeders but instead extend for much greater distances, typically several hundreds of kms. This contrast, if real, could reflect a difference in the rate of continental separation and/or magma viscosity. Thus, a slow rate of crustal extension may allow magma plugs to rise to high crustal levels before encountering stress conditions favourable to lateral magma flow as dikes, whereas a more rapid crustal separation could extend the depth of

Conference Reports

tensile fractures and allow dikes of greater height to be propagated upwards and outwards from the parent magma chamber.

Many examples of dikes radial to central plugs are known. In the Spanish Peaks of Colorado, lateral magma flow is well documented (R. Smith, U.S.A.). Vesicle elongation, marginal scour marks and aligned phenocrysts show that individual dikes propagated as inclined tongues up and away from a central source, estimated to be at a depth of no more than 10 km., though many dikes here extend for 40-50 kms. The extent to which lateral flow of magma took place in the large continental swarms is more difficult to judge, but systematic geochemical variations along a 500-km dike in northern Ontario may reflect longitudinal variations in depth of emplacement (R. Ernst, Canada).

Figure 1: Participants seeking shelter from mafic dikes (Photo R. Ernst)

It was generally agreed that continental mafic swarms are generated during early stages of rifting related to break up of continents and that most ancient ones reflect a failed arm environment. For example, many swarms in the Canadian Shield represent failed arms above ocean-forming hot spots, and several show a variation in dike size, frequency and compositional diversity as they are traced towards the suspected source (W. Fahrigh, Canada). Spectacular examples of "failed arm" swarms about 2 Ga old are found at irregular intervals along the southern boundary of the Archaean Superior Province of Canada. Comparable or even more extensive ones of Mesozoic age trend NE-SW across the Amazon (A. Gibbs, U.S.A.), or NW-SE in southern Brazil (A. Sial, Brazil), Paraguay (M. Druecker, U.S.A.), Botswana and Zimbabwe (J. Wilson, and D. Hunter, South Africa). The latter swarm extends for at least 1000 kms and may have been injected along the crest of a crustal upwarp owing, by analogy with Fiske's gelatin model, its great length to the lateral injection of magma from a source near the continental margin.

Variations in spacing and dip of dikes in a 2.6 Ga-old swarm in the Canadian Shield can be interpreted in terms of broad-scale warping of the Shield, with dikes becoming thinner and less abundant at deeper crustal levels (R. Ernst and H. Halls). Indeed, high-grade terrains in Canada, Australia and India are relatively free of abundant thick dikes.

Discussions of dike geochemistry focussed on the degree and nature of crustal contamination and the extent to which dike magmas reflect variations in mantle composition. I. Campbell (Australia) argued that these magmas, which in the

lower crust may have more komatiitic compositions and turbulent flow become more tholeiitic and more viscous upon contamination. The resulting laminar flow permits crystallization at dike walls in the upper crust, effectively insulating the dike from further contamination.

As regards dike chemistry, it is clear that incompatible elements Ti, Zr, P and Sr as well as Fe^{+3} and V are enriched, and Fe^{+2} , Mg and Si depleted with decreasing age. Major element abundance may also be partly a function of the particular structural province in which the dikes occur. Variation in chemistry with time cannot be explained by fractional crystallization or by different degrees of melting but requires mantle sources that are chemically distinct.

Although mafic dikes are difficult subjects for radiometric dating, J. Hanes (Canada) reiterated the usefulness of ^{40}Ar - ^{39}Ar methods on deuteric and primary hornblendes. The recent discovery of baddeleyite, a zirconium oxide, in gabbroic rocks (T. Krogh, Canada) opens the exciting new possibility of applying U-Pb techniques.

A preference for magnetization directions lying in the dike plane exists for swarms older than about 1.3 Ga, indicating a preference for meridional emplacement during much of the Precambrian (E. Tanczyk, Canada). Remanent magnetization characteristics have been used to determine paleotemperatures, using heat transfer theory. These temperatures are useful not only in calculating the depth of burial, but also in evaluating the loss of radiogenic argon in the host rock.

The conference was sponsored by the ILP, IUGS and several Canadian organizations. Plans are underway to publish the proceedings in a Special Paper of the Geological Association of Canada, to propose a new IGCP project on dike swarms, and to organize a follow-up meeting in Australia in a few years.

After reports by W. Fahrigh and H. Halls.

CALL FOR PAPERS

SECOND INTERNATIONAL SYMPOSIUM ON THE DEVONIAN SYSTEM AUGUST 17-20, 1987 CALGARY, ALBERTA, CANADA

- **Paleogeographic reconstructions**
regional and local
- **Scientific exhibits**
poster, core and paleontological displays
- **Field excursions**
- **Selected topics**
global reconstructions, basin evolution, event stratigraphy, biostratigraphy, paleontology, clastics, reefs, black carbonates and shales, petroleum and economic geology, others

DEADLINE FOR ABSTRACT SUBMISSION IS FEBRUARY 1, 1987.
REQUEST SYMPOSIUM CIRCULAR FROM:

THE CANADIAN SOCIETY OF PETROLEUM GEOLOGISTS

505, 206 - 7 AVENUE S.W. CALGARY, ALBERTA, CANADA T2P 0W7 (403) 264-5610

CANADIAN SOCIETY OF PETROLEUM GEOLOGISTS

CALGARY, ALBERTA

Memoir 10 **Sedimentology of Gravels and Conglomerates**

Edited by E.H. Koster and R.J. Steel

*Hard Cover, 441 pages, 1984
22 x 28 cm, ISBN 0-920230-27-X*

**Cost: \$35.00 postpaid in Canada
\$38.00 postpaid outside Canada**

*Please send money in Canadian funds
with order or supply Visa card number
and expiry date.*

Proceedings of the 11th Congress on Sedimentology. Contains twenty-nine original papers by authors from thirteen countries on five continents. Age of gravels/conglomerates range from Proterozoic to Recent.

The Memoir's six sections are:

- Background Considerations
- Fluvial Processes
- Modern and Ancient Alluvial Systems
- Ancient Fan - Deltas
- Wave/Tide Dominated Systems
- Ancient Submarine Slope Fans

*Send order to:
Canadian Society of Petroleum Geologists
#505, 206 - 7th Avenue S.W.
Calgary, Alberta
Canada
T2P 0W7*