Global Megageomorphology

Oracle, Arizona, U.S.A., 14 - 16 January 1985

Planetary geomorphologists tend to look at planetary landforms and processes from a global perspective, while geomorphologists studying the Earth emphasize detailed local and regional studies. An international workshop, cosponsored by NASA and the IUGS Commission on Comparative Planetology, last year explored the potential benefits of combining the perspectives and techniques of these two fields, for application to the study of the Earth as a planet.

The specialties represented at the meeting included oceanography, glacial geology, coastal studies, resource exploration, physical geography, remote sensing systems, eolian processes, landscape evolution, fluvial processes, tectonic geomorphology, and planetary geology. This initially overwhelming diversity of interests was soon found to be a complementary range of talents, because of the exciting new techniques, now available, especially remote sensing. These tools have arrived when important research thrusts in geomorphology are being defined or rediscovered. Moreover, geomorphologists are increasingly pursuing broad interdisciplinary concerns with tectonic, paleoclimatologic, and even extraterrestrial ramifications. In essence, we are seeing the beginnings of a "New Global Geomorphology" involving scales expanded in both space and time.

While many might question the meaning of megageomorphology, Clifford Embleton (U.K.) made it clear that the term conveys not so much a particular scale as a particular scale of thinking about landscapes. Robert Sharp (U.S.A.) made a more succinct statement: "Think big!" Rather than considering the details of single hillslopes, first-order drainages, or micro-processes, geomorphologists need to concentrate on large spatial scales and long time scales, comparable to those utilized in studying plate tectonics, biological evolution, and macro-climatic change. Embleton observed that the new tools, including remote sensing systems, mathematical modelling and radiometric dating, can enhance the traditional role of mapping in integrating geomorphology with tectonics and historical geology.

One modern concept that has proved quite useful is that of thresholds - critical conditions that must be achieved for geomorphological systems to undergo significant change. Dale F. Ritter (U.S.A.) noted the difficulties in evaluating process thresholds on a regional basis using remote sensing technology, but suggested that landform thresholds might be evaluated over large areas by continuous monitoring. Regions subject to profound climatic change or tectonic activity would be ideal study areas. Ritter uses the relationship of alluvial fans to climatic change as an example of a key scientific issue arising from such a study.

The antiquity of many landscapes and the inheritance of ancient landform elements in modern landscapes are themes that have, until recently, been neglected in modern geomorphology. A detailed review of the ancient landscapes that dominate the terrestrial planets of the Solar System was given by C.R. Twidale (Australia), who noted three major types of ancient planation surfaces: exhumed, epigene, and etch. He pointed out that it was only recently appreciated that there are major exceptions to the view that all the terrestrial landscape is late Tertiary or younger in age. Such anomalies are central to healthy scientific progress.

Sheldon Judson (U.S.A.) also saw an opportunity for geomorphology in the reconstruction of ancient landscapes and landscape processes. New geophysical techniques are providing novel approaches to the measurement of erosion rates. However, Ian Douglas (U.K.) pointed out that the complexities of climate and tectonism must be confronted in interpreting denudation. Douglas, like Twidale, sees profound megageomorphological similarities in the ancient Gondwana landscapes of the southern hemisphere, and in the Pampeanas of the central Andes Arthur L. Bloom (U.S.A.) has utilized a variety of satellite imagery to find additional Gondwana remnants.

The topic of tectonic geomorphology, especially as regards the relationship between drainage and tectonism, is one of growing interest. Laurence H. Lattman (U.S.A.) proposed a study of regional neotectonic movement and its megageomorphic effects, such as stream "slip off," variations in erosional resistance, coastal effects, and subtle eolian effects. Marie Morisawa (U.S.A.) also sees a fascinating role for imagery from space in the study of drainage patterns and channel networks; changes in time and space can thus be observed to understand drainage evolution.

Larry Mayer (U.S.A.) proposed the use of quantitative theoretical models to relate such megageomorphic features to global tectonism. Ian Douglas also observed that climatic geomorphology is posed for a reawakening, similar to that of tectonic geomorphology. The old schools of morphoclimatic zonation now need reassessment on an Earth

Figure 1: Part of the 3000-mile long Valles Marineris on Mars, a huge complex of canyons with walls modified by great landslides. NASA mosaic.

Conference Reports

that is being globally sensed and modelled, and the tropics provide an appropriate test area. In this connection W.N. Melhorn (U.S.A.) described some applications in the unusual landscapes of the Guiana Shield.

The contribution of planetary to terrestrial geomorphology had not been appreciated by many conference participants. B.K. Lucchitta (U.S.A.) noted that planetary exploration proceeds in a logical fashion: large features are identified and mapped before small ones, whereas on Earth the reverse was true. Wilfred Bryan (U.S.A.) emphasized that remote sensing has long been used to study mega-scale morphology in the 70% of Earth that is covered by water. Interestingly, as the new technology has elucidated smaller ocean-bed landforms, marine geologists have found themselves studying phenomena closely similar to those already documented by land-based geomorphologists.

Figure 2: Arsia Mons, one of the four great volcanoes on Mars. USGS image enhanced by computers and stereophotogrammetry.

Mars is an oceanless planet that has approximately the same surface area as the Earth. Sharp observed that Mars geomorphology can teach us much about Earth geomorphology (Fig. 1). The biggest lesson is the potential of different, unusual, or even outrageous explanations for puzzling landscape features. Are there "mega-scars" on Earth from ancient collapse phenomena, such as the chaotic terrain of Mars? What has been the role of catastrophes in geomorphological history? The role of sapping phenomena on Mars has already led to a reawakening of interest in sapping as an agent of terrestrial morphogenesis.

The opportunity to study morphological processes on several planets has great potential for understanding the fundamental nature of those processes. Ronald Greeley (U.S.A.) showed that aeolian geomorphology can be pursued on four planetary objects with a range of gravity, atmospheric pressure, surface temperature, and other parameters. The variation in aeolian landforms in these different settings can be used as a type of natural experiment to document the complex operation of wind systems. Similarly, Richard S. Williams, Jr. (U.S.A.) finds that many Icelandic and Martian volcanoes have remarkable morphologic similarity, though there are also some striking differences that raise basic questions about volcanic processes.

Perhaps the most obvious puzzle posed by the Martian landscape is that its geomorphic features, otherwise analogous to those on Earth, are so large (Figs. 2 and 3). Lisa A. Rossbacher (U.S.A.) listed the planetary parameters that might account for the size disparity, but no firm conclusions are yet available. The simple question remains unanswered: how large can a certain morphological feature be?

The problem of scale in geomorphology was raised in several other presentations. Robert S. Hayden (U.S.A.) saw major differences between the synoptic view provided by space imagery and the detailed view of landscapes used in conventional geomorphology. He noted that local scales are dominated by specific processes, such as volcanism, fluvial, or coastal, but that regional scales take on a unique character related to landscape diversity. Williams provided a detailed analysis of various satellite image scales that are useful in studying active morphological processes. He predicted a "golden age" for regional and global studies once high-resolution stereophoto coverage of the land areas of the world becomes available.

A final area of megageomorphological concern is that of applications. Floyd Sabins (U.S.A.) noted the fundamental input of geomorphology to natural-resource exploration utilizing satellite imagery is that explorationists and megageomorphologists can benefit from an interchange of ideas. Finally, Robert Blair (U.S.A.) reminded us of the application to education. Using W.D. Thornbury's classic fundamental concepts of geomorphology, he showed how space platform imagery easily illustrates the principles of the science.

The exciting technology that prompted our reassessment of geomorphology was discussed by J.P. Ford (U.S.A.), who dealt with the profound advances made in the use of spaceborne imaging radar. This system produces remarkable images of cloud-covered tropical regions and can highlight distinctive terrain and ice textures and topographic features. Another recent development is the Large Format Camera, described by Alta Walker (U.S.A.) who reviewed the results from the October 1984 Shuttle mission. The geometric fidelity, high resolution, and stereoscopic coverage of the Large Format Camera makes it an ideal geomorphological mapping instrument.

Of course, not all the technology ideal for megageomorphology yet exists. Williams proposed the optimum instrumentation for global quantitative geomorphologic studies. These would include an imaging system capable of acquiring photogrammetrically accurate stereoscopic images with a resolution of at least 10 m (adequate for 1:50,000-scale topographic mapping), a laser altimeter capable of measuring elevation differences of at least 10 cm, and a global positioning system for defining accurate positional accuracy. These systems should be operated for at least a decade in order to study the changing surface of the planet.

Several possible research thrusts in megageomorphology were discussed during the workshop. One of these concerns global geomorphic provinces and mapping. Remote sensing has a phenomenal capability for characterizing features on the Earth's surface. At the regional scale it has been useful to divide features into regions of similarity. The classic concept of physiographic provinces should be resurrected for use with new technology for geomorphological mapping. From a global approach a variety of research questions can be envisaged. What is the global distribution of volcanoes in relation to the Earth's heat budget? What are the details of terrain at convergent plate boundaries? What are the morphological signatures of suspect terrains? Can those signatures be used to correlate to other regions? Can landscape palimpsests be recognized that include relicts of ancient crustal processes?

Continental denudation: Geomorphologists at the turn of the century enjoyed a mutually beneficial collaboration with sedimentary geologists through their studies of erosion surfaces and nearby sedimentary basins. Remarkable recent advances have been made in the sedimentary analysis basin and the related erosion surfaces can now be dated, mapped,

Figure 3: Morphological evidence of past surface-water channels among Martian craters. NASA image.

and measured with great accuracy. The time is ripe to resurrect the global study of planation surfaces as a part of a new quantitative approach to continental denudation.

Geomorphology and tectonics: The "New Global Tectonics" has stimulated various branches of geology since its inception 20 years ago. Geomorphology is now playing an important role in this revolution for the earth sciences. Studies of tectonically deformed landscapes are establishing the rates of plate and intraplate movements, and regional studies are yielding tectonic interpretations from analyses of the structural expression of landforms.

Environmental change related to global processes: Sequential imaging of the planet has phenomenal potential for documenting both landform change and the operation of large-scale processes. Short-lived phenomena, such as various natural hazards, can be studied in remote areas of

the globe. Chronic problems such as desertification and deforestation can be placed in a regional and long-term context. It may prove possible to establish regional landform thresholds that will allow the prediction of future environmental change based on studies of existing and past change. This whole topic relates directly to the emerging international concern for global habitability.

Many suggestions for future work were made at the workshop. Specific case studies were proposed. For example, detailed megageomorphologic studies might be made of two stable cratonic areas, such as a comparison of Australia and the Canadian Shield. Similarly, attention could be focused on an area of active tectonism, such as the Andes or Himalayas, or on the geomorphological expression of the Appalachians versus the Caledonides. Alternatively there could be a global focus on a specific issue, such as the distribution of volcanoes, or an inventory of paleoclimatic indicators in the landscape.

Since both radar and the Large Format Camera were found to be of tremendous use in global geomorphological studies, future space missions, including shuttle flights, should give a high priority to imaging with these systems. Terrestrial Data Centers should be established in manner analogous to NASA's Planetary Data Centers. These would devote their attention to the cataloging and archiving of Earth remote sensing data, particularly the older Landsat data. Laser disc technology should be used to assist in the wide and economical distribution of these data, and the centers should also provide training in data access. Likewise, multiple data sets of various remote sensing products need to be assembled for terrestrial work and made available to the scientific community.

In parallel to the workshop, a book entitled, "Geomorphology from Space: A Global Overview of Regional Landforms," edited by Nicholas M. Short and Robert W. Blair, Jr., is now being prepared by NASA for publication in 1986. This volume will provide an excellent perspective on the full range of Earth and planetary landforms viewed from a variety of perspectives, and will be an additional step in viewing the Earth as a planet. □

James W. Head III
Department of Geological Sciences
Brown University
Providence, Rhode Island 02912
U.S.A.

Cretaceous of Eastern and Southern China

Tian Tai, China, 11-22 October 1985

The Cretaceous of southeastern China is dominated by continental sediments and volcanic sequences deposited in numerous fault-bounded basins (Fig. 1). The basic stratigraphy is understood in outline, but correlations and boundaries are still a matter of considerable dispute. Although these basins do not promise the abundance of the large oilfields of northeast China, they do have sufficient petroleum potential to warrant further studies of the structure and stratigraphy of the region. Against this background, the Paleontological Society of China and the

Chinese Petroleum Society recently sponsored a symposium on the Cretaceous of South China and its Petroleum Potential.

The nature of the Jurassic - Cretaceous boundary in this region was the principal point of discussion, but little was agreed upon. Likewise there was much debate on the biostratigraphy of the Cretaceous. The Jiande Group (Fig. 2) contains fossils indicating an Early Cretaceous age, notably the conchostracans *Yanjiestheria* and *Orthestheria*, and the

Figure 1: Occurrence of Upper Cretaceous basins of Zhejiang Province, China. 1 - Jinhua-Quxian Basin, 2 - Yongkang Basin, 3 - Tian Tai Basin, 4 - North Tandong Mountain.

bivalve *Nakamuranaia*. However, some fossils typical of the so-called Late Jurassic Jehol fauna of northeastern China were also found in these strata, e.g. *Ephemeroptera trisetalis* and *Ferganoconcha*. Further work is certainly required on the age ranges and paleobiogeography of the various fossil groups, but for the present the Jurassic - Cretaceous boundary is placed either at the base of the Laocun Formation or at the top of the Shouchang Formation.

There is an increased volcanic component in the basins in Zhejiang Province towards the east, as seen by the paucity of sediments of Jiande and Fangyan age nearer the coast. These are replaced here by the thick rhyolitic Moshishan and Tangshang Formation respectively. The basins are grouped into three northeast-southwest zones of Shouchang, Guantou, and Lanxi ages. This eastward younging of the start of basinal deposition is presumably related to phases of the Yanshanian Orogeny, thought to be then accreting slivers of material at the consuming margin of the Sino-Korean and Paleo-Pacific plates. Most basins are regarded as controlled by rift-faulting under tensional stress, but little work has been done on the structural aspects of this region.

The distribution and composition of hydrocarbons in these basins varies, and at present no commercial resources have been found. However, the grade and abundance of oil increases to the east with the greatest potential lying off-shore in the East China Sea.

Dinosaur eggs have been found in quite large numbers in the Quxian Formation (near Quzhou, Jinhua Basin) and in the Laijia B Formation at Tian Tai. Groupings of six to nine spherical eggs of 7-15 cm diameter have been found at each locality. No identifiable dinosaur bones were found in association with these, the only dinosaur known from this region being the megalosaur *Chilantaisaurus zhejiangensis* from the Zhongdai Formation near Jinhua City.

Figure 2: Schematic stratigraphy of the Cretaceous of Zhejiang Province, China.

A volume of 72 short papers on the Cretaceous of eastern and southern China (in Chinese with English abstracts) and field guide (in Chinese and English) were released at the symposium, the proceedings of which will be published by China Ocean Press. A forum was held to discuss a proposal for a new IGCP project on the correlation of non-marine Cretaceous sequences, and much interest was expressed. (This project has now been approved by the IGCP Board - see report on IGCP elsewhere in this issue. - Ed). □

Chen Pei-ji
Nanjing Institute of Geology and Paleontology
Academia Sinica, Nanjing
Peoples Republic of China
N.J. Mateer
1467 N. 17th Street
Laramie, Wyoming 82070
U.S.A.

Magmatic Evolution of the Andes — The End of the Beginning

Santiago, Chile, 18–20 November 1985

IGCP Project 120 on the igneous development of the Andes was initiated in Santiago in 1974 to establish an objective chronological base that would permit correlation of the tectonomagmatic events throughout the whole mountain range. The main aim of the project was the determination of the age of volcanic, plutonic, metamorphic and tectonic events in the Andean Cordillera and their correlation throughout the chain. The project emphasized geochronological research into the magmatic and tectonic evolution of different parts of the Andes, the study of the petrology and geochemistry of the rocks, and the relationship between tectonism, magmatism and metallogenesis. Progress was monitored through symposia in Neuquén (Argentina, 1978), Arica (Chile, 1979), Paris (26th IGC, 1980), Buenos Aires (Argentina, 1982) and Bucaramanga (Colombia, 1983).

The final meeting of IGCP Project 120 took place recently in Santiago de Chile and provided a fitting close to a most successful project. More than 200 papers on Andean geology have been published, containing over 2,000 new age determinations. With its outpouring of so much geochronological, geochemical and isotope data, the project has revolutionized our understanding of the Andes.

Precambrian rocks have been clearly identified in Andes de Mérida, Venezuela; Sierra Nevada de Santa Marta and Garzon Massif in Colombia; the Arequipa Massif in Peru; the Belén schists in northernmost Chile; in Santa Rosa de Tastil, southeastern Neuquén and northwestern Rio Negro in Argentina. Clearly, at least part of the Andes was built over old continental crust. Early Palaeozoic (450–400 Ma) volcanic-plutonic complexes in northwestern Argentina and adjacent Chile have been identified more precisely so that their tectonostratigraphy is currently interpreted in terms of accretion of exotic microcontinents to the northwest margin of Gondwana. A major accretionary volcanic-plutonic event in the Late Palaeozoic has been well documented in large areas of Argentina, Chile and Peru.

A broad synchronicity in magmatic activity along the Andes has emerged, with maxima of activity being apparent in the Late Palaeozoic, Cretaceous and Cenozoic. An important Jurassic plutonic event has been recognized in the Central Cordillera of Colombia, the Coast Ranges of central Chile, and in the Patagonian

Andes of Chile and Argentina. A gradual variation in the age of granitic intrusions across strike of the Andean Cordillera has been observed in several profiles across the Northern Andes of Colombia, southern Peru, northern and central regions of Chile and Argentina, and in the Patagonian Andes. A superposition of magmatic arcs of Late Palaeozoic to Late Cenozoic age has been observed between latitudes 38° and 40°S.

Extensive volcanic sequences, thousands of metres thick, which occur in Argentina, Chile, southern Peru and Bolivia, appear now to be younger than previously thought, on the basis of new K–Ar analyses with neo-Tertiary and Quaternary ages. Large batholiths of the same age have also been identified. Pliocene magmatic activity, recently discovered near the Taitao Peninsula of southern Chile, appears to be a major effect of the subduction of segments of the Chile rise under the South American plate in the Antarctica–Nazca–South American triple junction. This appears to be a beautiful example of the "blowtorch" effect that can be expected to accompany ridge subduction, as predicted by DeLong and others (see Forsythe et al., *Geology*, January 1986, p. 23–27).

Isotopic studies of the modern volcanic rocks have established regional variations in $^{87}\text{Sr}/^{86}\text{Sr}$, with values of 0.7050 to 0.7095 in Central Andes, and 0.7035 to 0.7055 in the northern Andes and southern Chile–Argentina. These distinctive isotopic signatures are now attributed to differences in crustal thickness and degree of incorporation

Figure 1: Conference participants in front of the Symposium headquarters.

Figure 2: Examining rocks from a Permo-Triassic intrusion that forms the background cliffs, near Uspallata, Argentina.

of subducted ocean floor sediments. Local variations of the same ratio in the southernmost area have been related to the intersection of oceanic transforms with the continental margin. Beryllium 10, formed by spallation reactions in the upper atmosphere and then concentrated in oceanic sediments, has opened a new field of investigation, serving both as a tracer for recycled sedimentary material and even as a pointer to sedimentation rates on the ocean floor prior to subduction.

The majority of intrusives related to copper mineralization have revealed Tertiary ages: in Argentina and Chile some are Cretaceous and a few definitively Late Palaeozoic. Jurassic intrusives do not develop porphyry copper systems with the exception of some recently discovered in Colombia. In the Altiplano of Bolivia and Peru, the most impor-

tant mineralization epoch is about 14-11 Ma, associated with subvolcanic stocks that produced high grade Sn-Ag-Pb deposits.

Of course, much is still to be learned about Andean geochemistry, its variations in time and space, and its relation to tectonic setting. Palaeomagnetic studies will have to demonstrate if terrane tectonics is a reality in the region. Indeed, the Andes will now start to exhibit more of those neat little holes indicating the passage of *Homo Palaeomagnetus*.

One study presented in Santiago shows that Cretaceous rocks from the Peruvian Cordillera have been rotated counter-clockwise by about 30°-40° relative to rocks of the same age in Chile. This is the same sense as the change in trend of the coastline and various other linear features that constitute the Arica deflection, seen by some workers as an orocline. As always, this preliminary work only points the way ahead, and other well-dated rock units must be studied on the craton and in the orogen before palaeomagnetism can be used effectively.

There can be no doubt that Project 120 has succeeded! Good and valuable science has been produced by a truly international effort, thanks in large measure to the groups of enthusiasts, particularly in Chile, Argentina and Brazil, who formed the core of the program. Project 120 has, however, achieved more than scientific success. Earth scientists from many countries have come together in friendship and cooperation for the ultimate benefit of the peoples of the Andean countries.

Extended abstracts of the 55 papers presented at the Symposium have been published, in English and Spanish, in a special volume containing abundant new geochronological and geochemical data on the Andes. This is available at US\$ 15 from Departamento de Geología, Casilla 13518, Correo 21, Santiago, Chile. □

After reports by Francisco Munizaga, Santiago, Chile, and N.J. Snelling, London, U.K.

Deep Internal Processes and Continental Rifting

Chengdu, China, 3-13 September 1985

Last September, an international symposium was convened in Chengdu by the Chinese Lithosphere Committee on behalf of the Inter-Union Commission on the Lithosphere (ICL) and the Chinese Association for Science and Technology, to discuss deep Earth processes and rifting of continental masses. Some 90 lectures were presented, about evenly divided between Chinese and foreign scientists. Many of the Chinese lectures were delivered in that language, then translated wholly or in part into English.

The papers ranged from quite general and theoretical discussions to detailed descriptions of individual areas. The emphasis was on geophysical data and their interpretation rather than on geological field observations of rifts. Indeed, the term "rift" was interpreted quite broadly to include some belts where reverse faulting predominates over normal. Rifts were described from many parts of the world, including the Rhein graben and other areas in Central Europe, but the main emphasis was on China.

Special attention was given to the great compound rift system that extends from central Manchuria south-southwestwards under the Gulf of Bo Hai and the delta of the Yellow River to the middle course of the Yangtze River. Much of this system is hidden under Quaternary alluvium or the sea, so that geophysical data are essential to understanding it. Parts are still active, and faults in Quaternary deposits in places display the local pattern. Since this rift system contains a considerable proportion of China's oil, a great deal of information has been assembled. Another area discussed by many workers was the Pan-Xi rift or Kang-Dian axis in southwest Sichuan and adjacent Yunnan, a zone containing important mineral deposits.

The meeting was pleasantly interrupted by a field trip to Guanxian and an extraordinary engineering project of the 3rd century B.C. Here the waters of the Min Jiang were diverted through a large and deep rock cut, hewed hundreds of meters through solid conglomerate, into a system of

Conference Reports

Figure 1: Normal fault dropping loess on the Palaeozoic at Kouzhen near X'ian. Photo by R.A.Price.

canals that irrigate the western part of the Sichuan plain, one of the rice baskets of China. The two kings, father and son, who organized and supervised this work are appropriately revered in a temple looking down on the intake and in another beside the rock cut called the Temple of the Tamed Dragon. The eloquence of this name becomes evident when one observes the constantly shifting, "boiling" currents of the silt-laden Chinese rivers. Indeed, part of the engineering work was designed to return as much silt and sand as possible to the main river while the water itself is led off into the canals.

Two field trips were organized after the Symposium, one to the Fen-Wei graben in the original Chin homeland around the old capital of Xi-an, and the other to the Pan-Xi zone around the entirely new city of Du-kou, where the Jin-sha Jiang (the upper Yangtze) comes out of the western mountains. The Pan-Xi zone can only be called a rift by a very broad extension of the term. Grabens are apparently forming now in certain parts of the area, in connection with the eastward escape of Tibet from between India and Central Asia.

During the Permian, a large region extending well beyond the Pan-Xi zone underwent strong extension and accompanying alkali-basalt or bimodal volcanism, but the Pan-Xi zone itself is marked by uplift rather than by subsidence. Where the evidence was sufficient to judge, the faults we were shown were reverse and not normal. The style seemed comparable to that of the Eastern Rockies of Colorado and Wyoming, and uplift occurred at least twice, once in the Triassic and once in either the Cretaceous or the late Cenozoic.

The unfailing courtesy and helpfulness of our Chinese hosts struck all of us who came from outside China, from the moment we were met at the Beijing airport to the time when we were returned to Beijing to go our separate ways. I learned a great deal about the organization of the earth sciences in China, about the geology of that vast and complex land, about the way in which the Chinese are combining their traditional ways and strengths with the influences coming in from abroad, and about the warmth and friendly openness of our Chinese colleagues. It won't be difficult to persuade me to go back for more. □

John Rodgers
Dept. of Geology and Geophysics
Yale University, New Haven CT 06511, U.S.A.

EPISODES

1986 Advertising Rates

Episodes has a total circulation of 3,000 and is distributed in 145 countries. It is published in March, June, September and December. Advertising orders should be received 45 days prior to publication date.

	<u>Size</u>	<u>1X Insertion</u>
Back Cover:	18.5 x 24.5cm (7-1/4"x9-5/8")	\$ 500(US)
Ins. Back Cover:	" "	450
1 page:	" "	400
1/2 page:	18.5x12.1cm (7-1/4"x4-3/4")	220
1/4 page:	9.3x12.1cm (3-5/8"x4-3/4")	130
	<u>Size</u>	<u>Minimum 4X Insertion</u>
Business cards:	3.5x5.8cm (1-3/8"x2-1/4")	\$ 100(US)

Write Episodes for mechanical specifications and rates for multiple insertions.

NEW from SWIFT (UK) MODEL "F" H/I AUTOMATIC POINT COUNTER

is a considerable advance over previous models. Now, with 12 channels -- one isolatable from total: Employs microprocessor for compactness, optimum reliability and extra facilities -- such as percentage key and up to 9x interval multiplier: Separate LED displays for channels, and for total plus combined 4-digit preset -- with audible signal.

Point counting is not limited to polarizing microscopes.

Attachments for all well-known microscopes.

H/I HACKER
INSTRUMENTS
INC.

Box 657, Fairfield, New Jersey 07007
(201) 226-8450