Pre-Variscan and Variscan of the Alpine-Mediterranean Belts

Cagliari, Sardinia, 25 - 31 May 1986

IGCP Project 5, with the above title, held one of its last field meetings earlier this year in Italy, in the beautiful setting of the University of Cagliari. Project 5 has really furthered the aims of the IGCP in establishing a firm basis for collaboration between geoscientists from 16 different countries, an association that will long outlast its own lifetime. The goal of Project 5 has been to shed light on the significant and complex patterns of both 'basement' involvement within a more recent (and well-studied) mountain belt and the development of the tectono-stratigraphic patterns of the older elements of the belt.

Throughout the project there has been a standardization of language, nomenclature and criteria. One of the many achievements has been compilation of bibliographies and review papers on the various member countries, thus facilitating access to the local literature. Research activities have been planned both nationally and internationally to achieve a common goal, and information has been well disseminated via regular newsletters, symposia and other gatherings. Field-meetings have, in fact, been held on 10 occasions in 9 of the member countries.

A major contribution of the project has been its attempts to achieve stratigraphic correlation across national frontiers. To this aim, a valuable data-base has been established, based on the compilation of a well thought-out series of Stratigraphic Correlation Forms. The final results of the project will be presented in two volumes, which are expected to be completed in 1987 and 1988 respectively. These volumes should have a considerable impact, not least for the impetus they will give to future work.

Figure 1: On the beach (L-R): A. Ramovs (Yugoslavia), G. Kleinschmidt (W. Germany), J. Sremac (Yugoslavia), I. Filipovic (Yugoslavia), Ch. Spassov (Bulgaria), F. Sassi (Project Leader) and Mrs. M. Sassi (Italy), E. Demirtasli (Turkey).

It was appropriate that the final field-meeting was held in Sardinia, a part of the mountain chain dominated by Variscan and pre-Variscan elements. Recent work, over the past decade, has shown that much of Sardinia comprises a series of major Variscan allochthonous thrust-sheets, which show a progressively increasing degree of metamorphism and of internal structural complexity upwards through the nappe stack. Sardinia is also the type area of the "Sardic Phase", i.e. the early Caledonian (Cambrian/Ordovician) orogenic stage of Stille. Indeed the possibilities for recognition of pre-Variscan orogenic activity was one of the themes that was given much attention during this field-meeting.

The Sardinia meeting was attended by 104 participants, of whom 60 took part in the field trip. The meeting included a one-day symposium where 6 review lectures, 36 short communications and 14 posters were presented, the very crowded schedule spanning the whole of the project activities.

The field-meeting was organized as a geotraverse that proceeded systematically through the Variscan terrain from SW at Inglesente to NE in the Olbia region. The excursion was efficiently and cheerfully guided by G. Vai of the University of Bologna, and his colleagues, who demonstrated the immense progress that has been made in recent years in the knowledge of the Variscan and pre-Variscan geology of Sardinia. Especially notable were the major advances made in the understanding of the tectono-stratigraphy, litho-stratigraphy and chronostratigraphy that have enabled reasonably sophisticated models to be developed for the evolution of this part of the mountain belt. Contributions detailed studies of the metamorphism to important advances made via geochronological studies. Many of these features are to be found in the well-prepared excursion guide-book published as Newsletter No. 8 by Project 5.

Field meetings such as these produce a natural feeling of 'cameraderie' which together with the Sardinian wines and exposure to the delights of Sardinian folklore, made for an unforgettable atmosphere. The organizers of Project 5, particularly the project leader F.P. Sassi (University of Padova) should be congratulated not only on the scientific success of this meeting, but also for the high level of enthusiasm of the participants. The writer hopes the rumour that yet another "Final Field Meeting" will be arranged in 1987 is well-founded. □

B.A. Sturt
Geological Survey of Norway
P.O. Box 3006, N-7001 Trondheim
Norway

Kaiko Project Conference on Subduction Zones

Tokyo and Shimizu, Japan, 10 - 15 November 1986

The KAIKO Project was the cooperative French-Japanese scientific study in 1984-1985 of deep-sea trenches and troughs (Fig. 1) around Japan. In its first year the survey consisted of detailed Seabeam mapping with precise geophysical measurements by the French vessel *R.V. Jean Charcot*. In 1985, carefully selected sites were investigated in 27 dives of the French submersible *Nautilie*.

Although the project itself was a bilateral one, this conference was more international, so as to permit discussion of structures and geodynamic processes beneath trenches and troughs in other parts of the world. Thus the meeting was attended by 30 overseas participants from U.S.A., U.K., New Zealand, F.R.G., Canada and Australia in addition to 28 French and about 100 Japanese scientists.

Figure 1: Japan Trench, Nankai Trough and associated areas.

The scientific sessions of the conference were opened in Tokyo by three overview lectures introducing the KAIKO results (X. Le Pichon, France), kinematics and dynamics of subduction zone complexes (J. Dewey, U.K.), and the relevance of the KAIKO results to the evolution of Japanese island arc (A. Taira and N. Niitsuma, Japan). It was followed by summary reports of the KAIKO cruises by their chief scientists.

The rest of the sessions at Tokyo were devoted to tectonic, geochemical and the biological aspects of fluids in subduction zones. KAIKO scientists in their dives at the landward lower slopes on Nankai Trough and Japan Trench discovered magnificent colonies of clams and benthic worms

as deep as 5,956 m, close to the limit of *Nautilie's* diving capability (Fig. 2). The temperature anomaly in the bottom sediments at the colonies (0.5°C higher than that of the surrounding floor) and a large content of thermogenic hydrocarbons in the venting water (reported by J. Boulegue and others, France) imply that temperatures at the source are at least 50 to 60°C. From thermal gradient data this source depth seems to be 2 to 4 km. All the results indicate that the ecosystem is maintained by seepage of cold water from deep portion of sediments in the landward prism of the trench.

Results of similar investigations of cold seeps along the Oregon-Washington margin, the Florida escarpment and Sagami Bay were also presented. J.C. Moore and J. Gieskes (U.S.A.) reviewed the results of recent drilling in the northern Barbados accretionary complex (ODP Leg 110), which appears to be hydrologically similar to the Nankai Trough. It was recognized that venting of pore water is controlled by a compressional force possibly associated with subduction tectonics and, therefore, long-term observation of seepage may be useful in monitoring secular changes in tectonic stresses.

In the scientific sessions at Shimizu held during the latter two days, 52 papers were presented on the tectonic interpretation of the KAIKO results and those from other areas. A remarkable contrast has been recognized between the Japan-Kuril Trench and Nankai Trough. In the former region tectonic and gravitational erosion prevails in the landward slope, whereas accretionary processes and intraplate deformation are dominant in the Nankai Trough. On the oceanward slope of the Japan Trench normal faulting of the crust is taking place. The collapse of two seamounts, Daiichi Kashima and Erimo has been carefully investigated, and it was confirmed that the 1500 m vertical offset of the

Figure 2: The world's deepest clam colony (5,956 m) being collected by manipulator of the *Nautilie*.

Conference Reports

east and west seamount bodies of the Daiichi Kashima resulted from repeated fault displacements between 2 Ma and 0.5 Ma ago. The Erimo Seamount is now being faulted to form fresh outcrops of basalts and phosphorites.

A contact of the western slope of the Daiichi Kashima Seamount with the landward toe of the Pacific plate was observed by the KAIKO dives. Several photographs of a converging plate boundary were shown, and modes of collision and subduction of such seamounts under trenches were postulated. Observations at the dissected walls of the Sagai Trough-Boso canyon provided actual profiles of compressional fronts.

Initial reports of the KAIKO Project are being printed in *Earth and Planetary Sciences Letters*, the papers on tectonics presented at the conference will be published in *Tectonophysics*, and the descriptive results will be published in book form. A series of topography maps based upon the KAIKO survey were published by the University of Tokyo Press (1986). □

Kazuo Kobayashi
Ocean Research Institute
University of Tokyo
1-15-1 Minamidai, Nakano-ku
Tokyo 164, Japan

Sediments Down-Under: The 12th International Sedimentological Congress

Canberra, Australia, 24 - 30 August 1986

For a soft-rock geologist, the quadrennial International Sedimentological Congress is an important event. The 12th of these was held recently for the first time in the southern hemisphere in Canberra, Australia, with the motto "Sediments Down-Under." It was sponsored by the International Association of Sedimentology, the geological societies of Australia and New Zealand and the Australian Bureau of Mineral Resources, Geology and Geophysics. This Congress was different from earlier ones both in style and in scientific emphasis, owing to the distinctive geological setting.

Australia is a largely semi-arid continent, lacking significant Cenozoic tectonism. There are vast tracts in Central Australia of what seem to be the world's oldest landscape. The continent lies adjacent to southwest Pacific arcs, and the New Zealand micro-continent and associated back-arc circulation impose a somewhat different regional perspective than that of the 'Atlantic community' of sedimentologists. Australia's passive margins are carbonate-dominated, with tropical carbonates, exemplified by the Great Barrier Reef and West Australian deposits, and the temperate carbonates of the Southern Ocean.

The Congress included 42 symposia, grouped under six themes: marine environments, processes and products; continental settings; earth history, tectonics and climate; sedimentology and fossil fuels; sedimentology and ore deposits; and concepts, techniques and history. The broad wide scope of the sessions, though satisfying to both specialists and generalists, required many hard decisions as to which of the ten concurrent sessions to attend, for example, reef complexes, tidal flats, or sedimentary structures.

There were many countries represented and the Chinese delegation was the third largest, after Australia and the U.S.A. The speakers from non-English-speaking countries braved well the ordeal and did a tremendous job! Despite the current "downturn" in petroleum and metal exploration, 938 people attended the Congress, in contrast to the anticipated 800!

There were 27 field excursions including those to Shark Bay (to see modern stromatolites), Canning Basin (Devonian reefs), McArthur Basin (mid-Proterozoic evaporites/stromatolites including one of the oldest oil accumulations

yet found), and the northern Amadeus Basin with its many excellently preserved depositional features in "unmetamorphosed" carbonate-siliclastic facies. Modern sediments were examined on beaches from Melbourne to Sydney, and the Great Barrier Reef was the site for 5 separate excursions. The Coorong-Spencer Gulf field trip offered carbonate specialists a rare treat - a lagoonal evaporite and carbonate environment under a mediterranean climate. A thunderstorm greeted the visitors as they examined the salt lakes, in sharp contrast to the conceptual models drawn for such sedimentary dolomite-forming settings.

During the Congress the Sorby Medal was presented to Robin C.G. Bathurst, and Gerald M. Friedman and Gerard V. Middleton were made Honorary Members of IAS. The Best Poster Award went to three Italian researchers (Piero Bellotti, Paolo Tortorzi, and Publio Uzleri) for their beautifully hand-coloured map and the "excellent science" of their presentation entitled "Sedimentological and Morphological Features of the Tiber Delta."

Conference Reports

During the Congress the new officers of IAS were elected and are as follows:

President:	H. Fuchtbauer (F.R.G.)
Past President:	H.G. Reading (U.K.)
Vice Presidents:	B. Beaudoin (France)
	R.N. Ginsburg (U.S.A.)
	D.A.V. Stow (U.K.)
General Secretary:	F. Surlyk (Denmark)
Treasurer:	P.L. de Boer (The Netherlands)
Editors:	P. Homewood (Switzerland)
	Ph. Allen (U.K.)
	J.A. McKenzie (U.S.A.)

Publication Secretary: M.E. Tucker (U.K.)

Council:	W.S. Al-Hashimi (Iraq)
	K.A.W. Crook (Australia)
	P. Timofeev (U.S.S.R.)
	J. Lajoie (Canada)
	L.A. Spalletti (Argentina)
	Sun, Shu (P.R. China) □

after report by Karl H. Wolf
Canberra, Australia

Extensional Basins and Paleotemperatures

Røros, Norway, 27 April - 1 May 1986

The Executive Committee of the International Union of Geological Sciences looks to its Advisory Committee on Fossil Fuels to keep it informed of on-going research in petroleum geology and to identify opportunities where initiatives might be taken to further existing projects. Earlier this year, the Committee organized its first workshop on Extensional Basins and Paleotemperatures, with the sponsorship of the Norwegian State Oil Company. The Røros workshop was aimed at increasing collaboration between university and oil company geoscientists in order to increase the effectiveness of geological research.

Petroleum geologists are confronted with targets that are increasingly difficult to discover. For the high-tech oil of tomorrow, the foreseeable increase in costs will allow little room to make mistakes. To increase their chances of discovering new oil at acceptable prices, petroleum geologists require a better understanding of the factors underlying the accumulation of hydrocarbons, phenomena that depend on the geological history of the basin. Subsidence and thermal factors are dominant in the maturation of organic matter, in the movements of the fluids and in the preservation of reservoir qualities. The understanding and modelling of all such phenomena are therefore absolutely necessary.

Much of the workshop focussed on the geodynamics of extensional and foreland basins. Following a general review by A.W. Bally (U.S.A.), B. Célérier (France) proposed that the evolution of subsidence and the thermal history of passive margins is the result of a crustal stretching coupled with a thinning of the lithosphere during the initial rifting phase. By applying a thermal geodynamic model for stretching and thinning to the eastern American margin offshore from South Carolina, he confirmed that there must have been a greater thinning of the lithosphere than of the continental crust. However, the lack of data on thermal properties and on the stratigraphy of the syn-rift deposits makes it impossible to characterize the precise thinning rate of the lithosphere. Célérier's numerical model gives a range of depths at which the organic matter in sedimentary series reaches a state of maturity favourable for the formation of hydrocarbons.

An analog geodynamic model coupled with a thermal and a geochemical one was applied by D. Issler (Canada) to the Cretaceous margin of Labrador. His model resembles that developed in 1982 by L. Royden to describe stretching and lithospheric thinning during rifting, according to the principles defined in 1978 by McKenzie. The description of heat

transfer takes into consideration variations in thermal conductivity during compaction. The geochemical model, designed to forecast the maturity of organic matter, uses vitrinite reflectance and the isomerization kinetics of chemical biomarkers to trace the thermal history of the basin. Both thermal and geochemical considerations are calibrated on experimental data.

For each borehole investigated, the evolution of the thermal flow is deduced from the computation of tectonic subsidence. For 80% of the boreholes examined, forecasts agree with the data observed. In cases of disagreement between observed data and computed values, it was found that the boreholes were situated in structural situations where subsidence is not representative of the region.

From their study of the Suez rift, P.Y. Chénét and I. Moretti (France) have shown that advection movements of the asthenosphere underneath rifts must be invoked to explain their structural evolution. They created a geodynamic and thermo-mechanical model that integrates the rheological behaviour of the lithosphere and asthenosphere with thermal evolution. The originality of this concept lies in its description of the physical phenomenon of lithospheric thinning.

The evolution of the Suez rift is characterized by a generalized subsidence in the Lower Miocene affecting the entire 80 km width of the rift. Chénét and Moretti attribute this initial subsidence to the regional stretching of the lithospheric plate. The subsidence zone then retracted progressively to be limited practically to the present Gulf (about 40 km). At the same time, the shoulders of the rift were strongly upthrust. According to the authors, this phenomenon can be explained by the cessation of regional stretching, followed by asthenospheric advection.

G. Quinlan (Canada) showed that the geological evolution of foreland basins is caused by loading due to successive overthrusts from the orogenic zone. Using a viscoelastic downwarping model, he demonstrated that during overthrusting, the lithosphere responds to the loading by a depression in the vicinity as well as by an uplift of a corresponding external flexural bulge. The depression becomes filled with sediments, while the bulge is subjected to erosion. By correlating the different erosion surfaces to the overthrust-phases in the Appalachians, Quinlan was able to characterize the wavelength of the lithospheric downwarps as Taconic, Acadian or Alleghenian. This model can also

Conference Reports

account for the stratigraphic and structural evolution of other foreland basins, and it can, thus, be used for studying the maturity of source rocks.

An example of thermal reconstruction as applied to the Provençal passive margin (NW Mediterranean) was presented by J. Burrus (France). He discussed non-uniform lithospheric stretching, taking into account the influence of impermeable saliferous horizons on compaction, local erosion, and fluid transfers. The lithospheric stretching model fully covers the evolution of this complex margin, and the accelerated subsidence observed must be explained in terms of its regional Alpine context.

These discussions confirmed that thermal and mechanical phenomena in the lithosphere and asthenosphere must be taken into account when explaining the geodynamic evolution of sedimentary basins. Thermal phenomena may be dominant, as in the case of passive margins for example, or they may be subordinate to mechanical phenomena (foreland basins). In all cases, numerical modelling provides data required to reconstruct the history of temperatures needed to compute the petroleum potential.

Thermal evolution also depends on heat transfer phenomena inside the sediments and is vital in the formation of oil as well as in the diagenesis and migration of fluids, especially hydrocarbons. The second session in the workshop was thus devoted to studying heat transfer in sediments, and it reviewed porosity evolution during compaction and during diagenesis.

G.V. Chilingar (U.S.A.) described the different techniques for revealing and explaining the presence of abnormal fluid pressures in formations. A numerical model of clay compaction that considers the kinetics of smectite-illite equilibria, was given by N.C. Dutta (U.S.A.). These kinetics appear to depend strongly on paleotemperatures, and abnormal pressures encountered in smectite-illite types of micaceous clays are to a large extent of diagenetic origin. The model has been applied successfully to pressure profiles measured in the Gulf of Mexico.

A two-dimensional, numerical compaction model (THEMIS) based on the concept of effective stress was described by P. Ungerer (France). Compaction and fluid-migration phenomena are described by linking Darcy's law to a mechanical equation that empirically relates porosity and effective stress. The presence of abnormally compacted (overpressured) zones is explained quantitatively by the existence of permeability barriers created by rapid sedimentation.

Figure 1: Participants in the Røros Workshop.

The model can introduce lithological variations by varying permeability and mechanical behaviour, and it takes into account fracturing phenomena. Likewise, it is coupled to the equation for heat and can thus simulate the thermal evolution of sediments at the same time.

C. Hermanrud (Norway) classified convective phenomena according to their origin as metamorphic, meteoric or diagenetic. He showed that fluid movements linked to the compaction of sediments always have a negligible effect on temperature evolution. D. Chapman (U.S.A.) emphasized the extreme importance of the quality of the data, in particular for determining heat gradients and flows from borehole data. He stressed the importance of compiling conductivity maps for making useful correlations with geology (e.g. facies variations). He also showed the existence of permeability thresholds separating regions dominated by conduction from those dominated by convection. The precise determination of permeability is a difficult task and is rarely achieved.

In a second paper, Chapman gave an example of the quantitative reconstruction of thermal effects induced by hydrodynamic flows in the Uinta basin in Utah. He emphasized the limitations to this type of modelling, including problems of poorly understood boundary conditions, the failure to consider transitory evolution, and improper understanding of the effects of paleocirculations. Quantitative calculations show that thermal effects linked to regional hydrodynamics cannot be ignored from a petroleum standpoint.

As regards free convection with sediments, M. Combarrous (France) showed that experience gained through theoretical analysis can shed light on problems linked to the possible appearance of this phenomenon in sedimentary basins. The exact modelling of a geological example encounters numerous limitations due to such factors as lack of understanding of permeability fields, presence of heterogeneities, and improper understanding of the boundary conditions.

The great sensitivity to thermal phenomena of mechanisms for the formation and migration of oil suggests that we should be able to reconstruct exactly the history of temperatures. In this regard, R. Pélet (France) gave a critical description of all methods using organic material, and he reviewed in detail actual geochemical methods. The important conclusion was that there is no universal calibration scale for methods for the evolution of maturity; each method reacts differently to different thermal histories. However, approximate calibrations can be of practical use.

Chemical methods based on ratios of concentrations of well-defined compounds (e.g. isomerization of hopanes, epimerization, aromatization) seem to be very accurate at first approach. However, they do not depend solely on thermal history, and other factors might even be more important. It is therefore impossible to choose the best method on the basis of theory. The proper choice depends on an examination of the internal coherence of series of measurements performed in basins of varying geological situations. At our present level of understanding, such an examination leads to the conclusion that the use of T_{max} values determined by Rock-Eval analysis together with vitrinite reflectance gives the most coherent results.

T. Throndsen (Norway) described the principles and gave various examples of measurement of vitrinite reflectance. Going beyond the conventional understanding of the term vitrinite, he showed how a precise definition of organic elements being measured can be used to eliminate much of the apparent dispersion in the measurements. This requires a clearcut preference for coals over organic matter

Conference Reports

dispersed in the sediment. Likewise, the difficulty of defining macerals in immature coals makes the method relatively unreliable for slight degrees of evolution.

Methods using mineral material were described by A. Gleadow (Australia) who concentrated on the use of fission tracks in apatite. The method has the great advantage of providing information about both the most recent time when apatite reached the temperature at which fission tracks are erased, and about the history of temperatures since that event. The combination of these two types of information is of great importance for reconstructing the thermal history of a sedimentary basin.

Each of the three preceding speakers emphasized the limitations of the techniques described, and expressed doubt that further progress could be expected in them in the future. The potential of mineral methods, however, has not yet been exhausted, and new and efficient ones may be discovered in the future, even though our present understanding of organic methods leaves little hope of any "miracle" approaches being found. Progress here will stem from greater understanding of the fundamental processes on which these methods are based.

Finally, P. Ungerer discussed laboratory experiments to determine the parameters governing the evolution of the Sarro-Lorraine Westphalian kerogen underneath the Paris Basin. The use of several scenarios to trace the geological and hence thermal history of the Westphalian formation led

to the evolution curve actually observed in natural samples. This type of study will develop to a much greater extent, especially since evidence from different paleothermal methods will be used to limit the number of scenarios to choose from.

Models such as those discussed at Røros will play an important part in the forecasting of phenomena, and we can expect much from them. However, their reliability will depend on the extent to which they can be controlled by comparison with actual cases. Proper control requires the high quality measurement of meaningful parameters. This will be done more easily as the researcher and the professional working on the site come to understand each other better.

Academic researchers must indicate clearly the physical significance of the parameters to be measured and provide a better understanding of their sensitivity. In exchange, operational explorationists must ensure access to proper data and give a critical view of their accuracy. It is illusory to use sophisticated equations when the parameters have been measured roughly, or only estimated. □

C. Sallé

Chairman, IUGS Advisory Committee on Fossil Fuels
IFP, 1A4 Av. de Bois-Preault
92506 Reuil Malmaison
France

EGT and the Evolution of the Northern Baltic Shield

Espoo, Finland, 5 - 22 November 1986

The Geological Survey of Finland recently hosted a workshop to analyse, integrate and interpret new and existing data pertaining to a polar crustal profile, extending southwest from Varanger Peninsula in northern Norway to northwestern Finland north of the head of Bothnia Bay. New deep seismic refraction and reflection data and electromagnetic depth soundings from the profile carried out in association with the European Geotraverse (EGT), and recent field observations were combined with aeromagnetic, gravity, geological, tectonic, Quaternary, geochemical and metallogenetic maps from the Nordkalott Project. The latter was carried out by the geological surveys of Finland, Norway and Sweden as a joint survey of their combined territories north of latitude 66°.

The workshop brought together 60 university and government scientists from seven countries and was divided into eight parallel working groups responsible for refraction and reflection seismology, gravity, aeromagnetism, electromagnetism, geochemistry, geology and paleomagnetism. These groups had access to the main computers of three universities and the nearby Geological Survey, where the rest of the material was stored.

The meeting was concentrated on the composite structure of the pre-Svecofennian (pre-2.0 Ga) foreland. The Polar Profile encompasses three granitoid-greenstone basement terranes divided by a pair of narrow, southeast-trending, imbricate, volcanic-sedimentary belts of probable Early Proterozoic age. The structural vergence of the northern (Kola) belt, within which the Soviet Union's Kola super-deep hole is located, is northward, whereas that of the southern

(Lapland Granulite) belt is southward. Intense interest revolved around the provocative hypothesis that the Kola Belt marks a southwest-dipping suture zone above which the Lapland Granulite Belt evolved as a back-arc basin.

Figure 1: Discussing the Polar Profile: L-R - G. Gall and H. Korhonen (Finland) N.V. Sharov (U.S.S.R.) and A. Berthelsen (Denmark). Photo by M.V. Knorring.

Conference Reports

Preliminary modelling of the refraction data shows a thin low-velocity layer at about 10 km depth beneath the southern third of the profile, and northward crustal thickening under the Varanger Basin at the north end of the profile. The latter suggests a flexural origin (foreland basin) rather than a rift origin (aulacogen) for the basin. Lateral variation in the inferred density structure of the deep crust is compatible with the Kola "suture" being rooted into the mantle 150-200 km southwest of its outcrop belt. The seismic reflection profile clearly substantiates the shallow northward-dipping structure of the granulite belt but does not resolve conflicting geological interpretations of its steeply dipping northern boundary. Electromagnetic data was modelled as a bifurcating, downward-tapered, resistive body in the upper crust, having a shallow conductive channel situated on the profile beneath the Lapland Granulite Belt.

The workshop was an exemplary experiment in mutual cooperation between university and government geoscientists from different countries and disciplines. The interplay

between geologists and geophysicists in a working environment was rewarding. However, the inconspicuous role of isotopic studies was surprising, given the North American experience that U-Pb geochronology often provides a decisive test for tectonic models and that it is the leading force behind the current surge in understanding the origin and evolution of the Precambrian shield.

A volume of reports prepared by working groups covering virtually all aspects of geophysics and geology, as well as an integrated synthesis of Archean and Early Proterozoic tectonics of the northern Baltic Shield is being prepared for publication in *Tectonophysics*. □

P.F. Hoffman
Geological Survey of Canada
588 Booth Street, Ottawa
Ontario, Canada K1A 0E8

Iberian Terranes and their Regional Correlation

Oviedo, Spain, 1 - 6 September 1986

This international conference, held at the University of Oviedo, reviewed the progress of IGCP Project 233 on Terranes in the Circum-Atlantic Paleozoic orogens. The conference began with a review of basic principles by R.D. Dallmeyer (U.S.A.) who graphically illustrated terrane accretion with photographs of his bathtub toys as the water drained out of the tub! B. Bluck (U.K.) then outlined how these units may be identified by mismatches in uplift and subsidence histories of adjacent areas, differences in provenance, and mismatches in the drainage and source areas. He also documented how the nature of terrane amalgamation produces different stratigraphic records. Thrusting produces coarsening-upward sequences, progradation of coarse over fine sediments and roll-up unconformities, whereas strike-slip amalgamation produces upward-fining sequences and variably oriented overlap sequences.

The need to establish whether faunal province boundaries relate to depth or climatic factors before they can be used to establish terrane margins was stressed by D.L. Bruton (Norway). He showed that a junction between a deep and a shallow-water faunal province probably does not coincide with that between adjacent terranes, rather it will migrate up and down the continental margin during transgressions and regressions. On the other hand, different shallow-water faunal provinces probably do demarcate different terranes.

J.D. Keppie (Canada) divided accretionary structures into three major categories: the main "accretionary" thrusts, large transcurrent "dispersive" faults and shear zones, and listric, normal "rift" faults, often located along earlier thrusts. Only some of these may act as terrane boundaries. The diachronism, up to 100-150 Ma, associated with accretionary structures indicates that the current subdivision into time-bounded orogenies such as Acadian and Variscan should be revised in terms of long-lived spatially distinct orogenies early in the convergence history, followed by widespread deformation during continental collision.

In a review of the problems of identifying the protolith after metamorphism, E-an Zen (U.S.A.) discussed the metamorphic consequences of emplacing nappes of various thick-

nesses, with pressure increases followed by an increase in the temperature and then cooling associated with uplift and erosion. W.E. Stephens (U.K.) also dealt with the consequences of nappe emplacement upon the generation of "S" followed by "I"-type magmas, with the melting of pelitic rocks followed by that of igneous rocks. The magma type, together with Pb-isotope signatures, may indicate the nature of the source. However, considering that most orogens are allochthonous, the junctions between S and I-types may not coincide with those between terranes. M.B. Stephens (U.K.) then reviewed the metallogenic consequences of terrane interaction, showing the different kinds of deposits associated with specific tectonic environments.

The conference then turned to Iberia. E. Martinez-Garcia (Spain) provided the framework by subdividing the Iberian Massif into 26 terranes some of which represent strike-slip

Figure 1: Terrane workers on the outcrop in northern Spain.

Conference Reports

repetitions of one another. These may have come from a deformed continental margin, a strongly deformed oceanic marginal area, a less deformed margin and oceanic area, or they may have accreted after the main amalgamation. Accretion is inferred to have occurred in several stages: prior to the Early Ordovician, in the Early Devonian and during the Carboniferous. A traverse of Iberian terranes from the northeast ("inboard") to the southwest ("outboard") was begun by R.H. Wagner (Spain), who reviewed the repetition of turbiditic-shelf Carboniferous sedimentation interspersed with four phases of deformation in the Palenian area.

In a review of the Late Precambrian-Permian miogeoclinal development of the Asturian terrane, E. Martinez-Garcia, showed that it was deformed by thin-skinned tectonics in the Late Carboniferous. A. Perez-Estuan and his colleagues (Spain) described the West Asturian Leonese terrane as a series of shallow water Paleozoic sediments interrupted by turbidites in the Late Ordovician, and all resting unconformably upon upper Proterozoic clastic sediments. These rocks were affected by thin-skinned and polyphase deformation accompanied by low to medium-grade metamorphism and intruded by syn- to post-kinematic granitoid plutons. A. Ribeiro and E. Pereira (Portugal) characterized the Galician-Castillian terrane as a pile of four allochthonous units

transported from the west, with a Silurian para-autochthon at the base, followed by lower Paleozoic continental rift to oceanic magmatics and high pressure Variscan metamorphics, by ophiolites, and at the top by Precambrian-Cambrian basement.

C. Quesada (Spain) interpreted the Ossa-Morena Zone as two Precambrian terranes (Sierra Albarrana and Valencia de las Torres-Carro Muriano) amalgamated in the late Precambrian (Cadomian-Pan African), overlain by Paleozoic overstep sequences and cut by major sinistral shear zones and late Paleozoic plutons. J.T. Oliveira (Portugal) showed that the South Portuguese terrane is separated from the Ossa-Morena Zone by an inverted ophiolite suite. The former is made up of Devonian clastics, bimodal volcanics with massive sulphides of the Pyrite Belt, and Carboniferous turbidites of northerly provenance, all thrust towards the south during Late Devonian-Carboniferous accretion with the Ossa-Morena Zone. Finally, E. Banda (Spain) demonstrated that the crust beneath the south Portuguese terrane is distinct from that beneath the rest of Iberia, which is generally composed of three layers totalling about 30 km in thickness.

The characteristics of the Iberian terranes were described by several authors in terms of igneous, tectonothermal, stratigraphic and paleontological data. Some papers dealt with the petrology, geometry and ages of plutons and their contact metamorphism and relationship to deformation. Others reviewed the tectonothermal evolution of terranes in the axial zone of the Pyrenees and in ophiolitic complexes in Iberia, and a comparison between the terranes in the South Portuguese Zone and southwest England was made. Stratigraphic analyses of the Late Precambrian and early Paleozoic in the southwestern Spanish Meseta, central Ossa-Morena terrane and in central Portugal were presented.

The paleobiogeography of Late Cambrian trilobites and Ordovician faunas and the Devonian-Carboniferous palynology of the South Portuguese Zone were also applied to terrane analysis. E. Martinez-Garcia invoked closing of the Hesperian Ocean beginning in the Early Ordovician, initial collision during Late Silurian-Early Devonian, and progressive convergence throughout Devonian and Carboniferous to end in the Permian.

Other workers surveyed data relating to the Paleozoic of Europe: the French Hercynides, Armorica, the Silesian-Polish Massif, the Eastern Alps, and the Caledonides in Scotland, Eire, Wales, Poland, Scandinavia and Spitzbergen. P. Matte (France) presented a model for the Variscan Orogen in Western Europe involving the closure of the Rheic and Galician-Massif central oceans. This is believed to have occurred between 420 and 280 Ma, with initial collision as early as 380 Ma ago. Migration of deformation, metamorphism and plutonism away from the sutures through time is a characteristic of the Variscan Orogen.

A comparison between an evolutionary model for the southern Mauritanides and southern Appalachians was presented by R.D. Dallmeyer and M. Villeneuve (France). They recognize three orogenic phases: Pan-African I, produced by the collision of a western block and the West African craton (cooling ages of around 650 Ma), Pan-African II structures (about 560 Ma), and Hercynian structures (around 280 Ma). These are comparable to terranes and events recorded south of the Brunswick-Alamaha Magnetic Anomaly in the southern Appalachians. This session ended with a general paper by T.P. Trurnit on the sequence that terranes pass through during global tectonic megacycles.

The conference also discussed legends for terrane maps. Thematic legends were produced, and the concept was developed of time-slice maps that combine various types of

Figure 2: Waiting for the next terrane to come ashore on the northern coast of Spain.

Conference Reports

thematic data. The projected scale to be used is 1:5,000,000, and the Decade of North American Time Scale will be used. The first to be produced will be a Preliminary Terrane Map of the Circum-Atlantic Paleozoic Orogens using the classification scheme as in the Circum-Pacific Terrane Map, and also showing overstep sequences and stitching plutons.

From the conference and the associated field trips the participants learnt a great deal about Iberian geology. For most foreigners this would have been difficult any other way

because most of the literature is in Spanish or Portuguese. As a result a sharper definition of the terranes in Iberia is now available, and participants should also have a clearer idea of how to define terranes elsewhere. □

J.D. Keppie
Department of Mines and Energy
P.O. Box 1087, Halifax
Nova Scotia, Canada B3J 2X1

First African Conference on Agrogeology

Zomba, Malawi 23 - 27 June 1986

One of the most important factors in a country's national security is the development of food self-sufficiency, and a necessary step in achieving this objective is to ensure a steady supply of fertilizer. This is where agrogeologists come on the scene, to search for and develop the geological raw materials farmers can use to promote the well-being of their crops. For the most part this is a question of fertilizer, but it also includes resources that can be used to correct the acidity or alkalinity of soils, improve soil structure, or help to cut down water losses from the land.

To discuss these matters a conference was held recently at the University of Malawi to look at the geological resources of east and central Africa that could be of use in agriculture. Inevitably, much stress was laid on phosphatic rocks, because phosphate, after nitrogen, tends to be the limiting nutrient in many tropical soils.

The 50 participants included geologists, soil scientists, agronomists and economists who provided information on the types of geological resources available, prospecting techniques, ways of processing and using them, soil problems to be dealt with, and the response of a number of important crops to fertilizers. A notable feature of the conference was a series of papers presented by African geologists on problems and prospects for agrogeology in their own indi-

vidual countries. The target was the small farmer, cultivating 5 hectares or less, for, as Zimbabwe has shown, the small farmer is the major factor in developing food self-sufficiency.

Particularly interesting discussions were held on a number of relatively simple techniques being developed in India, U.S.A. and Canada for making phosphatic rocks more reactive than is normally the case, without energy-intensive industrial processing. This included composting, use of zeolites and other cation acceptors, and partial acidulation using weak nitric acid produced in a kind of mini-hydro station by fixation of atmospheric nitrogen.

A plan emerged at the conference to set up an information network based in East Africa to collect and disseminate agrogeological information for the continent. The conference was supported by the International Development Research Centre, the Commonwealth Science Council, the European Economic Community, and the Government of Malawi. The proceedings will be issued by IDRC. □

W. Chesworth and P. van Straaten
Department of Land Resource Science
University of Guelph
Guelph, Ontario
Canada N1G 2W1

LIST OF ABBREVIATIONS

(Others are identified in the text or in Directory of IUGS Bodies)

BGR	Bundesanstalt für Geowissenschaften und Rohstoffe (F.R.G.)
COCORP	Consortium for Continental Reflection Profiling (U.S.A.)
EGT	European Geotraverse
GARS	Geological Applications in Remote Sensing
GGT	Global Geoscience Transects
IAH	International Association of Hydrogeologists
IAS	International Association of Sedimentology
ICL	Inter-Union Commission on the Lithosphere
ICS	International Commission on Stratigraphy
ICSU	International Council of Scientific Unions
IDRC	International Development Research Centre
IGC	International Geological Congress
IGCP	International Geological Correlation Programme

ILP	International Lithosphere Program
IUGG	International Union of Geodesy and Geophysics
NACG	North American Council of Geostatistics
NATO	North Atlantic Treaty Organization
ODP	Ocean Drilling Program
SPS	IUGS Subcommission on Precambrian Stratigraphy
UNESCO	United Nations Educational, Scientific and Cultural Organization

ERRATA

- The review of *African Geology* published in the last issue of *Episodes* (vol. 9, p. 190) erred in stating a time lag of four years between publication of the original work and its inclusion in this new bibliographic series. The average lapse of time is in fact 6 months. This well-received series is compiled from citations in the BRGM/CNRS data base (which is a source for GEOREF), as well as from information provided by CIEG and the Royal Museum of Central Africa.
- Carboniferous Stratigraphers Meet in Argentina (*Episodes*, vol. 9, p. 180). In line 13 of the left-hand column "Falkland Islands" should have read "Islas Malvinas."