The Earth Sciences in the Service of Africa

Dakar, Senegal, 9 - 11 December 1986

Several years ago, Unesco set up the African Network of Scientific and Technological Institutions to promote cooperation among universities in Africa. Operating from the Unesco Regional Office for Science and Technology in Nairobi, Kenya, ANSTI awards postgraduate fellowships, organizes workshops, meetings and courses, issues a variety of publications, and promotes scientific research at its member institutions. Recently the ANSTI Earth Science Sub-network held its first seminar at the Department of Geology, University of Dakar, attended by 40 participants from 10 African countries.

A session on Mineral Resources Development Strategies reflected the declining trend in the development of new mineral resources in Africa, with examples from Tanzania, Togo, Kenya and Nigeria. Several remedial measures, including proper data bases and specific incentives, were suggested for the revival of interest in the mineral resources of the continent.

The results of several geotechnical studies on materials for construction were presented, including strength characteristics of some Nigerian limestones, and the impact of geology on the performance of a bitumin pavement. Other topics discussed included geotechnical studies on materials for construction, a newly discovered phosphate deposit in

Togo, the economic potential of some low-enthalpy geothermal systems in Kenya, the role of reservoir engineering in geothermal energy development, flow trends and changes in the chemistry of groundwater from the Chad basin, and volcanism associated with various African orogenies. A comparative study of geochemistry of the Pan-African syenites of eastern Zambia and associated phosphate mineralization was presented, and there were several presentations on laterites from Nigeria, Cameroon, and Togo.

From discussions on the palaeogeography of coastal Tanzania during the Middle Jurassic, and the biostratigraphic implication of ammonites from the Malivundo Formation of Tanzania, it seemed clear that comparative studies and correlation are needed in order to improve biostratigraphic studies in Africa.

Plans are underway to publish both the abstracts and the full papers. The seminar will be an annual event and the second one is tentatively scheduled for Yaoundé, Cameroon, in December 1988. For further information contact: Dr. Isaac O. Nyambok, Department of Geology, University of Nairobi, P.O. Box 30197, Nairobi, Kenya, or Prof. S.M. Eno-Beling, Université de Yaoundé, Faculté des Sciences, B.P. 812 - Yaoundé, Cameroun.

after report by I.O. Nyambok, Nairobi

Megafaults of South America

San Juan, Argentina, October 1986

The city of San Juan, Argentina, located east of the Precordillera in a region with active faults was the site recently of a joint meeting of IGCP Projects 202 "Megafaults of South America" and 206 "Comparison of Active Faults," organized by Prof. Bruno Baldi of the Universidad Nacional de San Juan.

Recent right-lateral activity of the long-lived N-S El Tigre fault was well documented in satellite imagery and aerial photographs by M. Bastias (Argentina), and its fault scarp in Pleistocene sediments was examined in the field by the participants. The displacement along this structure since the Late Triassic is estimated at up to 100 km. The El Tigre fault parallels the continental margin, and in this region, 300 km from the trench, it is underlain by a shallow-dipping subduction zone. Structural and geomorphological evidence for recent thrusting in the Sub-Andean belt of Peru was presented by J.F. Dumont (Peru), who identified several phases of movement.

In contrast, no recent movement has apparently occurred along the Atacama fault zone between Taltal and Antofagasta. However, M. Hervé and J.A. Naranjo (Chile) produced structural, morphologic and radiometric evidence for normal movements during Miocene time along the zone. Linear structures on Quaternary infill that had been interpreted as evidence of recent movement are now explained by suffosion, the eruption of mobile or water-saturated material. The Atacama zone was also active in Early Cretaceous time, during the emplacement of various

plutons, which have regionally folded basic dykes (F. Uribe, Chile). There was at that time at least 30 km of left-lateral displacement, as indicated by disrupted plutons.

The relation of the megafaults in Colombia with the Caribbean-Nazca-South America plate interaction was analyzed by J.F. Toussaint (Colombia), who presented a tectonic interpretation of this northwestern corner of the continent. A triple junction system does not seem to apply, for the

Figure 1: Pleistocene clastic units cut by the La Rinconada reverse fault at the eastern margin of the Precordillera.

Conference Reports

plates appear to be separated by the Andean and Panama-Bando blocks along thrust and wrench faults and subject to contemporary internal shortening.

G. Sadowski (Brazil) grouped 50 well-defined major continental megafaults into 3 classes: structures affecting the Precambrian basement and of probable Precambrian or lower Paleozoic age with mylonitic belts; thrusts around the cratons and inside the fold belts; and faults that affected the platform during the Phanerozoic.

Evidence from the Pre-cordillera of San Juan and the Pampean Ranges of tectonic controls on sedimentation and volcanism since the early Paleozoic was presented by D. Salda and B. Baldi (Argentina). In the North Patagonian massif the activity was mainly of Late Tertiary age, with a presumed en echelon structure of the Neogene volcanic arc, in contrast to the more classical view of a simple alignment parallel to the trench.

The case for strike-slip faulting related to oblique subduction of the Chile rise was argued by the writer for the 1,000 km long Liquine-Ofqui fault zone, marked by what appears to be a mainly late Tertiary fault. More recent movements, however, have not been well documented.

Project 206 continues, but this was the final meeting of Project 202 "Megafaults of South America," which started in 1982. It is expected that a new project will be submitted to the IGCP Board, probably focusing on processes that take place along megafaults. In the meantime, the papers presented at San Juan will be published in a special issue of *Revista Geologica de Chile*.

Dr. F. Hervé

Depto. de Geología y Geofísica
Universidad de Chile
Casilla 13518, Correo 21
Santiago, Chile

Carbonatites Reviewed

Ottawa, Canada, 20 May 1986

A symposium held at Carleton University in Ottawa last year was the first in many years devoted solely to carbonatites. As Keith Bell (Canada) pointed out in his opening remarks, in the 20 years since publication of the Tuttle and Gittins and the Heinrich books on carbonatites, many new localities have been found, and the interest in these exceptional rocks is much wider, as attested by the audience of over 200.

In his opening discussion of the world wide distribution of carbonatites, the writer pointed out their well-known association with rifts and their commonly linear distribution, in some areas parallel to transform faults in the adjacent oceanic areas. He also noted their spatial and temporal association with orogenic belts. Geochronological studies indicate that in some areas there has been a repetition of carbonatite activity over long periods of time, suggesting that the lithosphere plays an important role in carbonatite petrogenesis.

The evidence for a magmatic origin was discussed by D.S. Barker (U.S.A.) in his review of emplacement mechanisms. He pointed out that late metasomatic replacement and hydrothermal processes are also important. After indicating that most carbonatites were emplaced later than associated silicate rocks, he argued that this implied separation of carbonatite melt from silicate melt by liquid immiscibility at a high level in the crust.

The mineralogy of carbonatites was reviewed by D.D. Hogarth (Canada), who confined himself to the important mineral groups pyrochlore, apatite and amphibole, for which compositional trends in carbonatites can be defined. The economic importance of pyrochlore minerals was discussed in some detail by A.J. Mariano (U.S.A.), with examples from Brazil and Paraguay. He pointed out that Nb is also localized in other minerals, and that some pyrochlores are important potential sources of tantalum.

A potentially powerful tool in the exploration for carbonatites was described by M.J. Kingston and L.C. Rowan (U.S.A.), who pointed out that reflectance spectra of REE-rich carbonatites give intense and diagnostic absorption bands between 500 and 900 nm due to the electronic processes of Nd. Airborne measurements over Iron Hill,

Colorado and Mountain Pass, California using a portable spectrometer enabled rapid discrimination between the REE, CO₂ and Fe⁺⁺ absorption bands.

A.H. Treiman (U.S.A.) stressed that carbonatite magmas differ from silicate ones in the absence of polymerized silica, which leads to exceptionally low viscosities, while the very low heat of fusion will cause rapid crystallization. In a magma chamber, crystallization will probably proceed in a dendritic manner from the walls inward, with dendrites readily breaking away and accumulating on the floor of the chamber.

A review by R.P. Sage and D.M. Watkinson (Canada) of carbonatite complexes in Ontario showed that there appears to be no correlation of mineral resources with age, but that those associated with ultramafic silicate rocks tend to be potentially more valuable. J. Pell (Canada) reported that in British Columbia unusual carbonatites occur in three belts, as both intrusive and extrusive varieties. Many form layers up to several metres thick, which may extend laterally for many kilometres, although the deformation and metamorphism to which they have often been subjected makes determination of primary emplacement mechanisms difficult.

Figure 1: S.E. Haggerty sampling a mica peridotite at Ile Cadieux, Quebec, Canada. Photo by K. Bell.

Conference Reports

M.D. Druecker (U.S.A.) described fenites associated with two recently discovered carbonatites and an alkaline complex in Paraguay. To identify fenitization, cathodoluminescence of feldspar was used to distinguish incipient, massive, and internal types of potassic fenites for one complex.

Stable isotope variations in carbonatites were covered by P. Deines (U.S.A.) who presented a plethora of data for C, O and S. The carbon isotopic composition defines a remarkably narrow range, more restricted than that of O or S, but observed values are considered to be consistent with current ideas on carbonatite mantle compositions. The isotopic composition reflects both variation in the source regions and crystallization and emplacement processes. Primary variations relate to tectonic setting, and may also suggest reflect depleted mantle.

Sm-Nd and Rb-Sr systematics were utilized by K. Bell and J. Blenkinsop (Canada), the symposium organizers, to demonstrate that many Canadian carbonatites were derived from a mantle source depleted in LIL-elements and with a mean age of about 3000 Ma; this region was coupled to the continental crust until at least 110 Ma ago. Data from East Africa is interpreted as reflecting mixing of two mantle sources, one with a depleted signature and the other with Nd and Sr isotopic characteristics similar to bulk Earth.

Work on Pb isotopes by G.R. Tilton and S.T. Kwon (U.S.A.) supports the Sm-Nd and Rb-Sr data in so far as a depleted mantle is also needed to explain the data for both the carbonatites and syenites studied. In isotope correlation diagrams, the slopes of regression lines increase with the age of the complex, suggesting an affinity with the source of modern oceanic volcanics. Crustal contamination can be recognized from some of the isotopic data.

Isotopic evidence was also cited by H.L. Allsopp (South Africa) and S.C. Eriksson (U.S.A.) for ancient lithosphere enrichment in their study of the Palabora complex. Sr, Nd, O and C isotopic signatures of carbonatites and pyroxenites here characterize a source region enriched in LIL elements and $\delta^{18}\text{O}$. The isotopic data indicate mixing of two isotopically distinct magma sources, one similar to Group I kimberlites and the other more enriched than Group II kimberlites. This enrichment, not detected beneath eastern Canada or East Africa, suggests a fundamental difference in the subcontinental mantle of southern Africa.

The concentrations of Zr and P_2O_5 required to stabilize Zr and P-bearing minerals in carbonated alkaline melts coexisting with peridotite at 14 kb were described by J.K. Meen, J.C. Ayers and E.J. Fregeau (U.S.A.). They suggested that three types of carbonatite source regions may be produced through metasomatism by carbonated hyperalkaline magmas, namely an olivine-carbonate-pyroxenite, wittlockite-olivine-carbonate-pyroxenite, and amphibole-carbonate-pyroxenite. These source regions would have distinct trace element patterns and radiogenic isotopic signatures.

Continuing the experimental theme D.H. Eggler (U.S.A.) discussed the melting relations of peridotite with amphibole, dolomite, magnesite and fluid. His new PT diagram, which was consistent with experimental data but which had not yet been tested experimentally, was presented as an alternative to Wyllie's arrangement, which involves a crossover in $\text{CO}_2\text{-H}_2\text{O}$ fluid-melt partitioning and a "blip" on the solidus. The two models were contrasted and constraints placed on lithosphere mantle metasomatism by the experimental evidence discussed.

"Kimberlite-carbonatite Relations - Brethren or Distant Cousins?" was the title of a most lively paper by S.E. Haggerty (U.S.A.). He pointed out that although carbonate

segregations are widespread in kimberlite, they are readily discriminated from carbonatite by their content of silicate-incompatible elements. However, the issue has been clouded by comparisons of the xenocrystic oxides in kimberlites with late-stage groundmass oxides in carbonatites. Neither kimberlites nor carbonatites can be derived directly from fertile garnet lherzolite of the asthenosphere. Although differing significantly in their trace element chemistry, carbonatites and the carbonate of kimberlites are derived ultimately from similar sources, the distinctions being primarily the result of time-integrated metasomatism.

In their review of the latest experimental work on carbonatites and liquid immiscibility, D.L. Hamilton and P. Bedson (U.K.) showed that phonolite and nephelinite melts are immiscible with alkali-rich carbonate melts, the gap widening with increasing pressure and falling temperature. However, no immiscibility exists between CaO-rich carbonate and silicate melts. The partitioning of a range of trace elements between immiscible carbonate and silicate melts was found to be generally similar to that in the carbonatite and phonolite lavas from Oldoinyo Lengai. M.J. Le Bas (U.K.) invoked partial melting, fractionation and contamination to explain the chemical and mineralogical diversity of carbonatites. However, liquid immiscibility played the dominant role in his system of producing a broad spectrum of carbonatite magmas from nephelinitic and phonolitic melts.

Figure 2: Distant view of Pande Hill, southern Tanzania, a Cretaceous carbonatite complex. Photo by K. Bell.

A different scenario for carbonatite evolution was offered by J. Gittins (Canada). He suggested that liquid immiscibility has been much exaggerated and that it is a late, low-pressure phenomenon that plays only a minor role in carbonatite petrogenesis. He favoured the melting of carbonate-bearing peridotite at pressures greater than 27 kb, with nephelinitic magma generated separately and not parental to carbonatite. Both magmas pursue independent evolutionary paths characterized by extensive fractionation in the crust. Alkali-rich carbonatitic magma, such as the natrocarbonatite of Oldoinyo Lengai, is not parental, but rather a late residual magma.

Gittins felt that too much emphasis had been placed on the exotic minerals of carbonatites and not enough effort devoted to the rock formers; the petrography of carbonatites is still poorly understood, partly because journals nowadays do not like to publish papers on this topic. Despite

outstanding recent advances in experimental studies, the work on immiscibility is still incomplete. Gittins noted that the Russians do not now emphasize metasomatism so strongly, and that there is still some confusion on kimberlite-carbonatite relationships. The identification and appreciation of extrusive carbonatites has advanced considerably in recent years, but more accurate field and mineralogical work and opportunities for discussion, like the Ottawa meeting were needed.

Although there is still controversy over the role of liquid immiscibility in carbonatite genesis, there was a remarkable consensus at the Ottawa symposium on the important role of the lithosphere, the evidence coming from field, trace element and isotopic studies. The idea of carbonatites being

characteristic solely of stable intra-plate environments is also changing, with occurrences being found close to destructive as well as constructive plate margins, in oceanic islands, and spatially and temporally related to orogenic belts.

The repetition of carbonatite magmatism in relatively restricted areas must be taken into account in any petrogenetic scheme. Isotope and trace element geochemistry seem to be offering some solutions. One hopes that the next carbonatite meeting will take place in the not too distant future, and that it will be equally as fruitful.

A. Woolley
London, England

Quaternary Sediments of the Arabian Gulf and Mesopotamia

Kuwait, 7 - 12 February 1987

Those who attend international conferences have come to expect that most of the technical sessions will be tedious obstacles to the main function of the meeting, namely the opportunity to meet colleagues informally and to catch up on recent advances in the subject. The Conference on Quaternary Sediments in the Arabian Gulf and Mesopotamian Region held recently in Kuwait confounded this expectation by devoting three days to meetings and three to field excursions, and by including in the lecture program many items that the audience actually relished hearing. In addition there was ample opportunity for personal discussion, fuelled by lavish refreshments, in the elegant premises of the Kuwait Foundation for the Advancement of Sciences (KFAS), the sponsors of the conference with Kuwait University.

For many participants this was a rare opportunity to meet and hear some of the legendary figures of Gulf sedimentology, including B.H. Purser (France) editor of the carbonate geologist's vademecum *The Persian Gulf*, published in 1973, and D.J. Shearman (U.K.), in many minds indissolubly linked to the sabkha cycle. We were also able at long last to discover at first hand what had been achieved in recent years in Mesopotamian geology, a topic on which most international journals have been unduly reticent. And we heard some fine papers on the Gulf itself, notably from some of the younger geologists of Kuwait.

Keynote papers reviewed the sedimentological implications of the contrasting tectonic regimes of the Red Sea and the Gulf, sea-level control of sedimentation, the potential applications of isotope dating to Gulf research, and a comparative analysis of depositional environments in the Gulf, the Bahama Bank, Shark Bay and Florida-Yucatan. The first day of talks was devoted to regional sedimentation in the northwestern Gulf and the influence of neotectonics and sea-level changes upon it; the second to carbonate and evaporite sedimentology in the Gulf and to sedimentation in Mesopotamia; and the third to applied and comparative sedimentology (see Gunatilaka, *Episodes*, December 1986 for a regional review - Ed.). The organizers emphasized poster displays firmly linked to the lectures. The scheme permitted effective development of controversial or difficult points especially where discussion following the paper at issue had been thwarted by garrulous delivery.

Despite a cold, biting wind and the nearness to a bloody battlefield, the field excursions were undoubtedly the highlight of the meeting. A southern trip, led by A. Gunatilaka (Kuwait), focused on the Holocene carbonate-evaporite complex of the Al-Khiran region, while A.Z. Al-Zamel (Kuwait) led two groups to the clastic intertidal and supratidal environments of Kuwait Bay. The excursions gained much from the combined presence of those who had made the original observations and those still engaged in wresting data from the mud and sand (Fig. 1).

Since the classic work of the Shell and Imperial College groups, the main task among carbonate geologists working in the area has been to consolidate the pioneering studies and extend the guiding principles to other parts of the Gulf. Despite considerable advances there are some familiar gaps in our understanding. For example, the evolution of the head of the Gulf is destined to remain controversial until oil company data are released or exploration boreholes are drilled to supplement those studied by Macfadyen in 1936. Likewise if the fluvial history of the Tigris and the Euphrates remains puzzling, it is partly because it has received little detailed attention since the 1960s.

Figure 1: Examining coastal exposures of Kuwait. D.J. Shearman second from right.

Conference Reports

The tectonic development of the Gulf during the Quaternary is still poorly understood: more work is needed on dating deformed coastal deposits, and the seismic data that should complement this work are lacking. One of the few bright spots in this grey panorama is provided by Kuwait, where some of these gaps are being energetically filled by a new

generation of enthusiastic geologists. Their published accounts and the volume planned by the conference organizers at the Geology Department, Kuwait University are eagerly awaited.

C. Vita-Finzi
London, U.K.

Remote Sensing and Neotectonics

Bogota, Colombia

27 October - 3 December 1986

The need for neotectonic studies as a background to seismic hazard assessment has been underlined tragically by the recent events at Popayan, Mexico City, San Salvador, and now Ecuador. In view of the urgent requirement for appropriate training of earth scientists in seismically active parts of the Andes, Central America and the Caribbean, 25 scientists from many parts of Latin America took part recently in a successful training course on the application of remote sensing techniques for the identification and mapping of neotectonic phenomena.

The course was held in Bogota at the Centro Interamericano de Fotointerpretación (CIAF) and organized jointly by CIAF and the International Institute for Aero-space Survey and Earth Sciences (ITC) of Enschede, the Netherlands. Vital support was received from the Fundación Venezolana de Investigaciones Sismológicas (FUNVISIS) of Caracas, Venezuela, ORSTOM (France) and the University of San Luis, Argentina. The Commission of European Communities provided for generous fellowships including air passages, acquisition of materials (almost 5,000 copies of aerial photos and Landsat and SLAR imagery) and the participation of foreign lecturers.

The course concentrated on the practical use of aerial photos and other remote sensing imagery, and there was a strong emphasis on morphotectonics as a means of detecting active tectonism. Lectures on geomorphological and tectonic principles, morphotectonics, geotectonics and regional tectonics were complemented by practical exercises and workshops with the stereoscope on examples chosen mainly from the Andean terrain of Venezuela and Colombia. Use of Landsat imagery was demonstrated by a series of images from China. This was further supplemented by field excursions near the city of Bogota to examine detailed trench mapping techniques and to demonstrate indirect evidence of neotectonism.

A five-day excursion was made to Central and South Colombia with visits to the Ibagué and Algeciras faults. The former traverses the capital of the Tolima province and is an active tectonic feature crossing a huge alluvial fan and giving rise to extraordinary well-developed pressure ridges, sagponds and associated phenomena. The Algeciras Fault, by some considered to be the South American-Caribbean plate boundary, has well-developed pull-apart basins, drainage anomalies, and other active geological features.

The course ended with a three-day International Symposium on Neotectonics and Volcanic Hazards, organized jointly by the Instituto de Investigaciones Geológico Minero

(INGEOMINAS) and CIAF and co-sponsored by AGID and several Colombian Institutes and Organizations. A second course is being planned for the beginning of 1989. Further information can be obtained from: H. Diederix, ITC, P.O. Box 6, 7500 AA Enschede, The Netherlands or from CIAF, Apartado Aéreo 53754, Bogota, Colombia.

After report by Hans Diederix

WELLBORE SAMPLING WORKSHOP

MAY 27 - 28, 1987

(Houston, Texas - near Intercontinental Airport)

WORKSHOP ON SCIENTIFIC WELLBORE SAMPLING TECHNIQUES to identify and prioritize the equipment needs to obtain scientific solid, fluid, and gas samples from drilled boreholes both in ocean basins and on deep continental drilling projects.

SPONSOR: Joint Oceanographic Institutions, Inc.; U.S. Science Advisory Committee; Department of Energy/Basic Energy Science. CONTACT FOR INFORMATION: R.K. Traeger - Sandia National Laboratories, Albuquerque, New Mexico (505) 844-2155; or B.W. Harding - Ocean Drilling Program, Texas A&M University, College Station, Texas (409) 845-2024.

THE WORKSHOP WILL (1) summarize the current state of need for borehole samples to be used in geoscience research, (2) summarize the state-of-the-art in sampling technologies, and (3) use the interactions of geo-scientists and engineers to prioritize hardware needs for effective borehole sampling.

APPLICATIONS ARE INVITED from potential workshop participants. FUNDS from a JOI-USSAC contract will be distributed to defray travel and subsistence expenses on the basis of applicant's eligibility and statements of interest and need. Applications for funding must be received by May 15, 1987 - addressed to B.W. Harding, TAMU Research Park, 1000 Discovery Drive, College Station, TX 77840.

GEOMORPHOLOGY

New!

The journal will publish review articles, research papers, book reviews and letters related to pure and applied geomorphology.

Editor: M. Morisawa, Binghamton, NY, USA

GEODERMA

An International Journal of Soil Science

Editor-in-Chief: R. W. Simonson, College Park, MD, USA

PALAEOGEOGRAPHY, PALAEOCLIMATOLOGY, PALAEOECOLOGY

An International Journal for the Geo-Sciences

Editors-in-Chief: D. V. Ager, Swansea, Wales, UK,
E. J. Barron, Miami, FL, USA

REVIEW OF PALAEOBOTANY AND PALYNOLOGY

An International journal stimulating interdisciplinary cooperation and understanding among workers in the fields of palaeobotany and palynology.

Editor-in-Chief: W. Punt, Utrecht, The Netherlands

Quaternary Dating Methods

edited by W. C. Mahaney, York University, Ontario, Canada

(Developments in Palaeontology and Stratigraphy, 7)

"This is proceedings of a symposium made accessible at its best." (Chemical Geology)

Pleistocene Vertebrate Faunas of Hungary

by D. Janossy, Natural History Museum, Budapest, Hungary

(Developments in Palaeontology and Stratigraphy, 8)

This handbook is outstanding in Europe for its quality and completeness.

Proceedings of the International Symposium on Red Soils

edited by Li Ching-kwei, Institute of Soil Science, Academia Sinica, Nanjing, P.R. China

The book contains valuable reports dealing with the red soils of Africa, Asia, Latin America and Oceania.

Groundwater Modelling

An Introduction with Sample Programs in BASIC

by W. Kinzelbach, Institute of Hydraulic Engineering, Stuttgart University, FRG

(Developments in Water Science, 25)

This book covers material taught in a graduate-level soil physics course at Washington State University.

These and many more will be on display at the Elsevier booth during the INQUA meeting in Ottawa, July 31st-August 9th, 1987.

ELSEVIER SCIENCE PUBLISHERS
P.O. Box 330, 1000 AH Amsterdam, The Netherlands

87001