Book Reviews

TECTONIC SIGNIFICANCE OF THE HUNGARIAN MOUNTAINS IN THE ALPINE EDIFICE

Acta Geologica Hungarica, Akadémiai Kiadó, (Hungarian Academy of Sciences), Budapest, v. 27, no. 3-4, 1984, and v. 28, no. 1-2, 1985.

(Editor's Note: Although *Episodes* does not normally publish reviews of regular journals, nor of publications more than one year old, an exception is being made for the volumes here, because they report important new work on Hungarian geology, which complements the article in this issue by Halmi and Hámor.)

Hungarian geology is extremely important in unravelling the history of the Alpine belts in Europe. The connection between the western and Austrian alps with the Dinarides and Carpathians must be made through Hungary. The external parts of these Alpine mountain systems can be connected by surface geology to the north and south of Hungary, but the critical internal parts of the system must be pieced together from very limited outcrop and subsurface information from below the extensive late Cenozoic cover of the Pannonian and adjacent basins. The Pannonian Basin complex also is worthy of detailed study because it is a young basin system that has not been overprinted by post-basin tectonic activity. Recently, sedimentary basins have become a major focus of research, and the Pannonian Basin may well become one of the best known basins in the world because of the wealth of geological data available on this relatively simple basin system.

These two recently published volumes deal with both the largely convergent tectonics of Mesozoic and early Cenozoic time and the extensional regime that has been superposed on these older structures during late Cenozoic time. During all of these tectonic events, strike-slip faults played an important role. The papers in these volumes resulted from a joint research effort that was part of the Franco-Hungarian Technical Scientific Cooperation Program, conducted by researchers from the Eötvös L. University (Budapest) and Université Pierre et Marie Curie (Paris). The results of this four-year cooperative effort, which also involved other non-academic research workers from several

other Hungarian institutions as well as from other countries, were presented in a workshop in Veszprém, Hungary, in 1983, and many of the papers presented there have been published in these volumes.

The first paper is by J. Dercourt and others and reviews the Hungarian Mountains within the Alpine Framework, using many of the new concepts developed during the project. Dercourt et al. separate European units, such as the Gemericides, from Dinaric units in the Bükk Mountains by the Neotethyan oceanic Darno unit. Subduction is interpreted to have been northward within the Darno unit in Late Jurassic-Early Cretaceous time. Early movements along the Trans-Hungarian line were of a transform character during Mesozoic time, and it was later reactivated as a south-verging thrust zone. Some units in the Bükk mountains are interpreted as tectonic windows through the Darno Unit. The late Cenozoic Pannonian Basin was developed on top of these older tectonic units as Neotethys was subducted southward within the external Carpathians to the north.

The concept of southward thrusting in the Bükk is not new, but the emphasis on large-scale south-vergent thrusting is new, and data are presented in other papers to support the concept. For example, J. Geyssant and C. Lepvrier present an analysis of the microtectonics in the Bükk region to support the interpretation of major south directed thrusting. Z. Balla presents data that the ultramafic rocks of the Darno unit may have developed in a marginal sea environment.

A. Galácz, F. Horváth and A. Vörös present an interpretation of the sedimentary evolution of the Bakony Mountains proposing that they developed as a passive margin facing south from Early Jurassic to Cretaceous time. Their paleogeographic reconstruction shows that the Bakony Mountains belonged to the Periadriatic region of Tethys and developed within the South Alpine paleotectonic realm.

T. Báldi and M. Báldi-Beke interpret the Paleogene basins, developed before the younger Pannonian basin, as a product of transtension along major early Cenozoic strike-slip faults. Their work is based on new biostratigraphic data, and shows that the basins were deep, elongate and characterized by rapid subsidence. Basin development was coeval with significant crustal shortening in the external Flysch Carpathians.

F. Horváth and J. Rumpler use new multi-channel seismic data to interpret the development of the Pannonian Basin as the result of extension and subsidence superposed on the older alpine nappe structure. Some seismic lines are presented as examples of the quality of the available data (see also a collection of papers focussed largely on the geophysics of the Pannonian Basin in Kilényi, E. and Teleki, P., 1985. Cooperative Research in the Geosciences between the U.S. Geological Survey and the Central Office of Geology, Hungary, during 1978-1984. U.S.G.S. Open file Rept. 85-291, 336p. - Ed.). Extension began in Middle to Late Miocene time and continued into Pliocene time. Basin development was dominated by extensional normal faults and related strike-slip faults. Locally compressional structures are present that can be related to transpression.

These two volumes contain many more papers on various aspects of the stratigraphy, paleontology, paleogeography, geochemistry and bauxite development from the Pannonian region. Many papers are presented that concern the pre-Cenozoic rocks of the Pannonian region and that must be considered by those interested in the Alpine mountain chains or the Pannonian Basin of eastern Europe.

B.C. Burchfiel
Cambridge, Massachusetts

Attention Mineral Explorationists

One operator, using the EM-4 coil with an IGS-2 Console, magnetic and VLF sensors can efficiently record up to 12 different geophysical parameters in solid-state memory.

The new Scintrex EM-4 is here to improve your ground electromagnetic surveys

This unique EM receive coil, 25 cm in diameter and weighing only 1.8 kg, is the third sensor now available for the Scintrex IGS Integrated Portable Geophysical System. It complements magnetic and VLF sensors which, with the IGS-2 System Control Console, are already proving that multiple parameter surveys save time and money in ground geophysical surveying.

The advantages of the programmable, microprocessor-based IGS instrumentation are numerous. They include automatic recording of coordinates, time and data in solid-state memory and outputs to printers, modems or microcomputers to list, plot, transmit, store or process data.

An IGS-2/EM-4 Receiver can be used with either moving or fixed source Scintrex transmitters for rapid or deeply penetrating surveys. Either GENIE amplitude ratio or standard Slingram measurements can be made. The wide frequency range, signal/noise enhancement, simplicity of operation, complete interpretation procedures and numerous other features place the EM-4 in a class by itself.

To find out more about how our new EM-4 coil can improve your EM surveys, contact Scintrex at 222 Snidercroft Road, Concord, Ontario, Canada L4K 1B5, telephone (416) 669-2280, telex 06-964570.

SCINTREX

THE MEASURE OF EXCELLENCE

BOOK REVIEWS

MICROCOMPUTER APPLICATIONS IN GEOLOGY

J.T. Hanley and D.F. Merriam (eds.)

Pergamon Press, Oxford, New York, Toronto, Sydney, Frankfurt, Tokyo, Sao Paulo, Beijing, 1986, 274p., £25/\$37.50 (US)

During the past decade relatively low-cost, desktop microcomputers and their peripherals have become integral tools in the geosciences. A few journals and some microcomputer user newsletters have addressed the interests of geoscientists, yet surprisingly few books have been published describing and extolling microcomputers and their applications in the geosciences. This volume of 19 papers written by geoscientists begins to fill this void.

The contributions range from general topical overviews to descriptions of specific hardware configurations and software. Several papers include useful lists of references, and others include the source codes for computer programs described in the text. Programs described in five of the papers are available at minimal expense from the Computer Oriented Geological Society, and the editors state that "other software mentioned in the book are generally available from the authors."

The volume begins with two review papers, the first on the state of computing in the geosciences, and the second on available microcomputer hardware and software, including a brief survey of the trends in microcomputer use in geology. Although written with the explorationist in mind, the second paper would be useful for any new or potential microcomputer user.

In view of the extensive use of microcomputers in the exploration industry, it is appropriate that examples of microcomputer applications in geophysics are described in five papers. These describe the use of microcomputers in the reduction, interpretation, and display of various types of geophysical data including gravity, magnetic, electrical, radiometric, seismic refraction and reflection, and drill-hole log-plotting examples.

Artificial intelligence (AI) programs typically use multiple inputs to arrive at a single output or decision. Expert systems are programs in which decisions are made by following certain rules based on the answers to specific questions, where the rules are based on the opinions of experts in a particular area. One interesting and well-written paper describes two such systems (EXPLOR and PROSPECTOR) in which AI techniques are used for oil, gas, and mineral exploration. American examples quoted include the Mt. Tolman molybdenum deposit in eastern Washington and oil and gas leads in the Denver-Julesburg Basin.

Geological databases of various types and database management systems (DBMS) are widely used by geoscientists. Three papers in this compendium collectively describe programs for building and accessing databases, the manipulation of the data held in such files, and some of the pitfalls to be avoided in designing geological databases. One of

these papers describes the use of Micro-GRASP, which is a scaled down BASIC version of a mainframe DBMS used by the U.S. Geological Survey.

Three papers describe computer programs with specific geological applications. Program ROCALC contains a series of subroutines for the analysis and manipulation of geochemical data from igneous rocks. In addition to CIPW (and other) norms, the program calculates various ratios, ternary projections, and common chemical indices. The paper includes the Applesoft BASIC course code and some examples of the interactive output.

Program TRAVERSE allows a user to construct geologic cross-sections of folded terrains. After the user inputs the elevation and bedding-plane orientations along a traverse, the program calculates the vertical exaggeration and apparent dips. It then plots (on a monitor screen) the topographic profile and the interpolated positions of the folded geologic contacts. The Pascal source code is included in the text. The third paper describes a new computational method for constructing Eh-pH and activity-activity mineral stability diagrams and for plotting them on a monitor screen. The authors use an example of the geochemical behaviour of manganese during the early diagenesis of some marine sediments.

Another paper offers a good review of the use of microcomputers in geological image analysis. The discussion of new developments in image analysis hardware, software, and applications, such as remotely sensed images and images of microscopic textures, is augmented by a list of 64 references.

The last three papers describe various aspects of map and spatial analysis by microcomputer. One describes the general features and hardware requirements of mapping application software, while the other two discuss the use and limitations of a digitizer (graphics tablet) in map analysis and other spatial studies such as thin section analysis. One of the papers includes the Applesoft BASIC source code for the program DIGITS, which includes a simple routine for creating spatial data files as well as doing some simple spatial analysis such as distance and area measurements.

As might be expected from any group of 19 papers, the quality of writing varies considerably, as does the level of expertise required to appreciate the papers. Most of the figures used in this volume are clear and useful, but a few of the papers seem overburdened with unnecessary figures or with figures that are not adequately described in the text. It is clear that a wide variety of microcomputers and computer languages is being used by geoscientists today. However, the breadth of material in this book is such that it is likely that any geoscientist who has an interest in microcomputers will find at least several rewarding papers.

This book will not teach a novice how to use microcomputers, but it does clearly demonstrate how microcomputers are currently being used by geoscientists. For many readers the most valuable aspect of the book will be the source code listings and information about how to acquire the programs described in the text. However, given the rapid rate at which microcomputer hardware technology has been advancing and new microcomputer programs are being written to take advantage of this technology, it is not surprising that some of the papers in this volume are now out-of-date.

R.G. Clark, Jr.
Albion, Michigan, U.S.A.

Book Reviews

THE CONTINENTAL CRUST: A Geophysical Approach

Rolf Meissner

Academic Press Inc., Ontario, Canada, 1986, 426p., (P) \$50.75 (Cdn), (H) \$101.50 (Cdn)

This book is deceptively small. It contains over 400 pages of densely packed information on the physical properties, composition, structure and evolution of the continental crust. It is not to be read for a leisurely guided tour through the present state of knowledge of the continental crust but instead provides a series of intense, forthright 'executive summaries' of just about every aspect of its subject.

In his geophysical approach, Meissner provides definitions and a short comparison of planetary structures and then plunges into a blow by blow account of how the physical properties and state of the crust are measured. The methods and results of gravity, seismology, seismicity, seismic refraction and reflection, electrical sounding, stress measurements, geomagnetism and geothermics are all encapsulated in short reviews. The information on samples examined in the laboratory is then similarly reviewed, with special attention paid to creep data and absolute age dating. Each of these summaries presents methods, basic formulae, data, examples and the current understanding of the subject.

Exactly halfway through the book, the author turns to a brief summary of petrology, metamorphism and sedimentation and the basic principles of geology. This is followed by a series of accounts of continental structure in various geologic provinces. In these, the author provides concise reviews of geophysical investigations in the Precambrian Shield; Paleozoic and Mesozoic areas; Cenozoic Mountain Belts; Continental Grabens, Rifts and Margins, and Active Margins. The treatment of these presents very valuable regional data summaries, with line drawings of recent deep seismic reflection results (up to 1984) and predominantly North American and European examples.

Uncharacteristically, active margins receive a very perfunctory and inadequate treatment (5 pages), considering their importance in continental evolution. The latter is the subject of the last chapter in which the author presents his personal view of the crustal evolution of the earth's continents including plate tectonic type models and processes. The book finishes with a discussion of the mechanisms of continental growth and crustal accretion (vertical and horizontal) and the crust mantle recycling process.

The book as a whole is a powerful and intense attack on the understanding of the continental crust. It is ambitious in scope and only infrequently falls below the high standards that the author sets for himself. However, because its

spread is so wide, it is one of those books that leave the reader feeling overwhelmed. Its information density is so high that one doubts the ability of most specialist readers to follow and assimilate the author's progression from geophysical data and argument to geologic evolution. I judge its principal value as that of a geophysical reference and source book on the continental crust. As such, it would be a valuable addition to the graduate student bookshelf.

R. Riddihough
Ottawa, Canada

QUATERNARY GLACIATIONS IN THE NORTHERN HEMISPHERE

V. Šibřava, D.Q. Bowen and G.M. Richmond (eds.)

Science Reviews, Vol. 5, Pergamon Press, Oxford, New York, Toronto, Sidney, Frankfurt, 1986, 514p., charts and tables, \$99 (US)

This eagerly awaited collection of papers is the final report of Project 24 of the International Geological Correlation Programme, "Glaciations in the Northern Hemisphere," which started in 1971. The project involved active collaboration between "scientists from most of Europe, the United States, Canada, the U.S.S.R., the People's Republic of China and Japan."

The 51 papers are grouped into six parts. Over one-third of the volume (18 papers) is devoted to the "Quaternary Glaciations of the United States of America"; one brief paper covers Mexico. "Quaternary Glaciations in Canada" are discussed in four papers; 19 address various aspects of "Quaternary glaciations in Europe"; current knowledge on "Quaternary Glaciations in Eurasia" are summarized in seven papers, and the volume ends with an 1 1/2-page essay on "Correlation of Quaternary Glaciations in the Northern Hemisphere." In addition 21 loose correlation charts are included.

The hardcover volume and the charts fit in a black cardboard box, which results in a convenient, handsome presentation. The typescript is clear and easy to read; the illustrations, line drawings and charts are very clearly presented and well laid out.

It should be pointed out that although only 36 pages are devoted to Canada, which has more territory affected by Quaternary glaciations than any other country, the Canadian IGCP Working Group chose to publish its results separately (R.J. Fulton, ed., 1984, Geological Survey of Canada Paper 84-10, 210p). The present volume thus includes only a general introduction and three regional syntheses. In choosing this "fast track method" of getting their results out, the Canadian Group saved international readers the onus of buying a volume thick with papers of local and regional interest, but with limited appeal, certainly to non-North American Quaternary scientists. Had the Americans and Europeans chosen the same approach the present volume would have been significantly reduced in size and, presumably, in cost as well.

GEOLOGICAL SOCIETY OF MALAYSIA

GEOSEA V PROCEEDINGS

VOLUMES I & II
(Bulletin Geological Society of Malaysia Nos. 19 & 20)

Some of the articles appearing include:-

Massive sulphide deposits and their possible significance to other ores
R.W. Hutchinson: Palaeogeographic development of west Sarawak
Denis N.K. Tan: Geological evolution of the Southern Philippines C.K.
Burton: Southeast Asia as a part of an early Palaeozoic Australian
Gondwanaland C. Burrell & B. Sturt: Tertiary basins of S.E. Asia
their disparate tectonic origins and eustatic stratigraphical similarities
C.S. Hutchison: Late Palaeozoic palaeogeography of Southeast Asia:
some stratigraphical, palaeontological and palaeomagnetic constraints
J. Metcalfe: The REE geochemistry of Lingshan W-Sn-bearing granites
and their applications to petrogenesis of the granites Yuan Zhongxing
et al.: Chromite deposits of Papua New Guinea P.M. Alenya: Recent
advances in exploration modelling for tin deposits and their application to
the SE Asian environment R.G. Taylor & P.J. Pollard: Some thoughts
on the development of the alluvial tinfields of the Malay-Thai Peninsula
D. Taylor: Base metal exploration in Sabah David T.C. Lee & H.S.
Weber: The nature and potential of gold mineralisation in Kelantan
L.H. Chu & D. Santokh Singh: Quaternary deposits of Thailand P.
Dheeradolok & W. Kaewyana: Soil landscapes in Peninsular Malaysia
S. Paramanathan & S. Zauyah: Aspects of the geochemistry of
Malaysian cassiterites W. Fuad Hassan: Geological evolution of the
Indonesian Archipelago H.M.S. Hartono & S. Tjokrosapetro: The
nature, distribution and genesis of certain authigenic minerals in the
stanniferous alluvial deposits of S.E. Asia K.F.G. Hosking: Global
tectonics and resources W.S. Fyfe: Tin-tungsten-bearing granites in S.
China and their metallogenetic relations Xu Keqin & Zhu Jinchi:
Hydrogeological activities in Peninsular Malaysia and Sarawak F.S.
Chong & Denis N.K. Tan: Status of uranium exploration in Peninsular
Malaysia L.H. Chu & F. Chand: Directions of geologic transport in
Peninsular Malaysia H.D. Tjia: Cathaysia, Gondwanaland and the
Palaeotethys in the evolution of Continental S.E. Asia Y.G. Gatinsky
& C.S. Hutchison: Marginal sea formation by rifting of the Chinese and
Australian Continental Margins and implications for Borneo C.S.
Hutchison: Mesozoic and Cenozoic regional tectonics and metallogenesis
in Mainland S.E. Asia A.H.G. Mitchell: Coal potential and exploration
in Sarawak S.P. Chen: The succession of vertebrate faunas in the
continental Mesozoic of Thailand E. Buffetut & R. Ingavat: Regional
controls of hydrothermal ore localization in northern Thailand P.
Asnachinda & S. Chantaramee: Late Palaeozoic glacial marine facies in
S.E. Asia and its implications P.H. Stauffer & C.P. Lee: Cretaceous
melange in West Kalimantan and its tectonic implications P.R.
Williams *et al.*: Recent advances in the knowledge of geology, mineral and
energy resources of Singapore since 1981 Ansafur Rahman & P.P.
Wong: The integration of remote sensing, terrain evaluation and
engineering geology in Southeast Asia Beaumont, T.F. & Hunt, T:
Recent advances in the knowledge of geology and mineral resources of
Vietnam since 1981 Le Thai Xinh & Nguyen Xuan An.

This 2-volume GEOSEA V PROCEEDINGS of about 500 pages each contains 95 articles presented at the Fifth Regional Congress on Geology, Mineral and Energy Resources of Southeast Asia held in Kuala Lumpur, April 1984.

To: Hon. Assist. Secretary
Geological Society of Malaysia,
c/o Department of Geology,
University of Malaya,
59100 Kuala Lumpur, MALAYSIA.

Date:

Order for GEOSEA V PROCEEDINGS

I wish to place an order for set(s) of the GEOSEA V PROCEEDINGS which will be in 2 volumes of about 500 pages each. Volume I will be available in April/May 1986 and Volume II in July/August 1986.

Both Volumes
Members
Non-members

Price
M\$50.00 (US\$21.90)
M\$125.00 (US\$53.20)

Name/Company/Institution

Address

Enclosed cheque/money order/bank draft for M\$/US\$

Prices include surface mail. For airmail please call 03-7577036 or telex UNIMAL MA37453.

Book Reviews

One essential component of science is the free flow of ideas among scientists conducting research in a given discipline or field. The fundamental vehicle for transmitting an idea is a clearly defined concept. A major synthesis such as this book, resulting from a ten-year international study sponsored by Unesco, should provide us with a set of well-defined concepts for future work. However, this is not done effectively here, and this volume will certainly not be remembered as a monument to stratigraphic nomenclature.

The same name is used for a time division (i.e. Sangamon, page 7), for a paleosol unit (i.e. Sangamon soil, page 20) and for a biostratigraphic zone (i.e. Sangamon pollen zone, page 20). In addition, the following are given equal rank as "time divisions": Late Wisconsin, Middle Wisconsin, Early Wisconsin, "Eowisconsin" and Sangamon. A more strict adherence to stratigraphic nomenclature would require that the Wisconsinan and Sangamonian stages be given a different rank from the Late, Middle and Early Wisconsinan sub-stages. This somewhat relaxed approach leads to confusion, which is not necessarily clarified by a note such as: "The 'restricted' Sangamon time division used in these reports corresponds to marine ^{18}O substage 5e and the 'Eowisconsin' time division corresponds to ^{18}O substages 5d and 5a. The Sangamon time interval recorded by the pollen encompasses the 'restricted' Sangamon and the 'Eowisconsin' time divisions in these reports" (page 20). This reviewer does not believe that the use of "restricted" and "Eo" are significant contributions to chronostratigraphic terminology.

As recommended by the North American Stratigraphic Code (Article 77b) and following earlier workers, it would seem simpler and certainly clearer to use the adjectival form for designating chronostratigraphic divisions (Wisconsinan Stage, Sangamonian Stage, etc.) and the noun form for the time-transgressive events (Wisconsin Glaciation, Sangamon Interglaciation, etc.). Perhaps the urgent need for an INQUA working group to recommend stratigraphic terminology based on clearly defined principles is illustrated by the following: "assignment of numerical ages to the late middle Pleistocene - late Pleistocene boundary and to the boundaries of the subdivisions of the late middle Pleistocene and late Pleistocene time divisions is required to ensure uniformity of time division terminology from one region to another" (page 8). One can only agree that clarification is needed.

In spite of very obvious shortcomings in the clear and correct usage of stratigraphic nomenclature, this volume will certainly be a standard reference for some time to come. The synthesis of the Quaternary geology literature of major regions with extensive bibliographies undoubtedly will be a valuable asset to most scientific libraries and to all those who can afford to buy the volume. For the first time we have a major summary in English of the state of knowledge on the Quaternary for all regions in the Northern Hemisphere. This alone makes the book valuable. The numerous charts, in particular the correlation charts, will be much appreciated by professors and certainly by students who should consider them as a challenge, stressing the need for the great amount of work that needs to be done before we have an adequate knowledge of this most important period for mankind, the Quaternary.

D.A. St-Onge
Ottawa, Canada

ACTIVE TECTONICS

Geophysics Study Committee; Geophysics Research Forum; Commission on Physical Sciences, Mathematics, and Resources; National Research Council

National Academy Press, Washington, D.C., 1986, 266p., U.S.A./Canada/Mexico \$26.95 (US); elsewhere \$32.50 (US)

The stated goal of this book is "to provide assessments from the scientific community to aid policymakers." Such an aim requires the most responsible analysis of the highest quality. The book, therefore, deserves careful scrutiny. With certain reservations discussed here, it can be recommended on the scientific and educational levels, as well as for policymakers.

J.C. Crowell begins with a simple and informative description of the different active tectonic realms along the western continental margin. The statement that "faults known to be recently active are most likely to rupture again" is, however, surprising in view of a quite contradictory statement given in the introductory overview to the book.

On the topic of "epeirogenetic and intraplate movements," L.D. Brown and R.E. Reilinger state that "two of the most destructive earthquakes" in the U.S.A. occurred in the "stable" eastern United States (New Madrid 1811-1812, Charleston 1886). For these intraplate earthquakes "movements of several centimeters are common, and several meters are possible." The authors point out that "Mitigation hazards from earthquakes in the stable interior remains one of the least tractable yet most important problems in contemporary tectonics." They conclude that "the question remains whether there exist heretofore unrecognized mechanisms for intraplate deformations." "Intraplate motions provide clues for our understanding of the state and evolution of stress in plate interiors," but there are still "considerable gaps in understanding intraplate motions."

Active faulting and associated hazards are well reviewed by D.B. Slemmons and C.M. Depolo. Usually there is a good correlation between earthquake size and fault rupture parameters (length, displacement, area). The difficulties in evaluating these parameters generally lead to underestimates. Active fault parameters can be used to estimate the magnitude of paleoseismic events. The segmentation of fault systems is shown to be important. Of fundamental importance for hazard mitigation are recurrence rates, which can be established from studies of paleoseismicity and seismic gaps. Intraplate faulting apparently causes problems. Slemmons and Depolo suggest that the "unexpected young fault" along the old Meers Fault may represent "a possible extensive Quaternary breakup of the midcontinental region along reactivated faults." They speak about conservative design of constructions, and seem pleased that the Auburn Dam project was abandoned for theoretical seismotectonical reasons.

R.S. Yeats discusses flexural slip and bending-movement faults, with worldwide examples. He proposes that active faults are much more widespread than realized today, though his evaluation of the "impact on society" seems rather meagre.

Book Reviews

The fluvial network is one of the most sensitive means of registering active tectonics, so that it is surprising to read in S.A. Schumm's review of "alluvial river response" that "only a few investigators have considered its effects on alluvial river," because these are "difficult to detect." He gives many examples of slow, aseismic deformations ignored by earlier studies.

An ambitious discussion of "coastal tectonics" by K.R. Lajoie is more theoretical than descriptive, and the author seems to be unaware of most of the work done in Europe. It is a rigid and out-of-date analysis stressing principles rather than facts. Data on his Figure 6.10, for example, on Fennoscandia are taken from a 1965 textbook, in which the data are redrawn from another textbook of 1936, going back in turn to primary sources in 1919. Though Lajoie realizes the impact on society, little is said about this. His examples from Alaska, California, Japan, New Zealand, New Guinea and Chile are very limited both for a national and a global survey.

Lajoie claims a "long-term tectonic stability along most passive-margin coastlines." This is certainly most relative, as can be seen on the heavily discussed coast of Brazil. Figure 6.7 is said to represent "a stable coastline" and is "used as an approximate tectonic datum." If this is true, the Netherlands and the North Sea, a well-known subsiding region, would be slowly rising. Holocene sea level records are strongly differentiated over the globe as a result of geoid changes, dynamic variations and crustal movements: simplifications are, therefore, misleading.

"The geometric relationship" is regarded by Lajoie as "a useful means of correlating." This method was proposed in 1929 by L. von Post (Sweden) and in 1930 by V. Tanner (Finland), but subsequent studies have shown that it does not work even within limited regions, except for rough approximations. We read that "where dated, faulted strandlines also yield rates of fault movements." This is definitely not true, for average figures have nothing to do with actual fault movements and are only minimum values. In discussing Alaskan data the author arrives at the conclusion that a new event is to be expected "in the near future" on Middleton Island and that it is "overdue" at Icy Cape. A close examination of the data, however, suggests that these events are to be expected at 800 and 1100 years from now, respectively. In reality, we are often faced with surprises, as P. Pirazzoli has shown in the eastern Mediterranean where a long-term subsidence trend (a "time-predictive trend" in Lajoie's terminology) was suddenly reversed by a major uplift event.

E.A. Keller gives an informative overview of the "use of surficial processes." Geomorphic indices are said to have been successful in providing "basic data necessary for long-term (tens to hundreds of years) earthquake prediction." It is good to read that fault-scarp degradation curves need to be derived locally before estimates should be undertaken, that is, data should not be transferred from region to region.

A short report on seismological techniques by C.R. Allen argues that paleoseismology shows "the most spectacular progress in studies of active tectonic processes," although "one must be aware that the chances of immediate success are not high." Modern seismology has revealed that earthquakes "are far more different from one another" than earlier believed, but Allen claims that "comparable magnitudes tend to repeat one another faithfully and periodically." This statement is both questionable and dangerous, as far as it concerns intraplate seismicity.

W. Thatcher's presentation of "geodetic measurements" is informative. The importance of the "seismic deformation cycle" is demonstrated, although it is stressed that "not all crustal deformation is elastic and recoverable." The agreement between changes in gravity, elevation and strain in southern California is impressive. The characteristics of the interseismic period lead the author to claim that "caution is thus required in extrapolating a few years of measurements, however precise, to the long term." Cases of good agreement between geodetic and geological results are demonstrated. The chapter ends with the sad truth that not until "large earthquakes will occur in these closely studied regions," that is, in a densely populated area, will we know "whether diagnostic crustal movement anomalies preceded their occurrence" - knowledge through disaster.

"Near-field tectonic geodesy" or the instrumental documentation of fault movements is reviewed by A.G. Sylvester, who gives good examples, together with figures of rates and instrumental precision. The list of "problems" is long, and the need of long-term recording is stressed. Great progress within the last 15 years is claimed by D.B. Nash in a study of the "degradation of fault scarps." Various hillslopes processes are described, and each is supposed to operate in fault-scarp degradation. The so-called "dating" of scarps seems very relative. What is most important is to identify young and recent fault scarps in areas where faulting is supposed to be absent or to occur very infrequently.

A good summary of some 26 different dating methods is given by K.L. Pierce. Biological methods, however, are not mentioned, despite the power and widespread use of palynology. The author stresses the dependence of "time window" and claims that "dating control is commonly the limiting factor" in active tectonics. We need to define not only the rates but also "the constancy or variation of these rates" with time.

D.P. Schwartz and K.J. Coppersmith discuss seismic hazard analyses in an instructive and well-written chapter. Fault segmentation is thoroughly discussed, and the authors point out that the exponential distribution of earthquake events per magnitude for large regions "is not appropriate" for individual faults; "the non-linear slope is real." The three main hazard models are discussed; the Poisson-exponential (with doubtful assumptions), the time predictable, and the renewal (usually providing "the most reliable estimates") models.

Volcanoes and their impact on, and relationship to, society are reviewed by D.W. Peterson. This quite different paper ranges from a solid description of volcanoes and monitoring techniques to a more philosophical discourse on public response. Peterson reminds us that many large eruptions (e.g. Krakatau, Pelée, El Chichon) followed quiet intervals, and that "intervals between eruptions may be so long that the hazards are overlooked or forgotten."

The book ends with a chapter by B.M. Crowe on volcanic hazard assessment for nuclear waste disposal. This is a regrettable end to a nice book. As soon as nuclear waste handling enters the picture, common geological sense seems to be lost, and scientists who usually are so critical, conservative and rigid in accepting new ideas, suddenly drop all caution in order to assist the politicians. First of all, it is impossible to isolate the volcanic question, for the entire geodynamic picture must be evaluated. The area under consideration by Crowe is an active volcanic one, which would hardly be chosen by a geoscientist. The reason why this area is being considered is that the Nevada Test Site is already contaminated (a "dead valley" in new sense). The so-called probability calculation is based on very uncertain volcanic activity and on unfounded assumptions about an

Book Reviews

exponential and a Poisson distribution. This is not good enough for a serious decision about high-level nuclear waste deposition. The absence of a complimentary discussion on tectonic-geodynamic effects is a major short-coming of the book.

A well-written and concise summary to the whole book is provided by the introductory "Overview and Recommendations" by R.E. Wallace. He states that "the historically most inactive sections of great faults, indeed, may be candidates for generating the largest future earthquakes." Both "tragedy of ignorance" and "benefits from understanding" are cited. The author poses the question "how many other unidentified active faults exist in seismically quiet regions of intracontinental regions"? From the Fennoscandian Shield, I can report the recent discovery of numerous faults and fractures and other paleoseismic indices of postglacial age. Wallace recommends a "fundamental research focusing especially on Quaternary geology and geomorphology, paleoseismicity, neotectonics and geodesy." The international work of the Neotectonics Commission of INQUA is one relevant activity.

N-A. Möner
Stockholm, Sweden

REEF DIAGENESIS

J.H. Schroeder and B.H. Purser (eds.)

Springer-Verlag, Berlin, Heidelberg, New York, London, Paris, Tokyo, 1986, 455p., \$59 (US)

This valuable and stimulating collection of nineteen case studies is the product of the 5th International Coral Reef Congress in Tahiti in 1985. It embraces the diagenesis of reefs and mud mounds of Cenozoic, Mesozoic and Palaeozoic ages. In all, the papers give a good idea of the state of the art, and they illustrate the application of a range of approaches involving the use of mapping, core logging, thin sections, strains, cathodoluminescence, stable isotope ratios, trace elements, X-ray diffraction, scanning electron microscopy, EDAX, fluid inclusions and ultraviolet light fluorescence. A brief introduction by the editors provides a clear and up-to-date background to the studies that follow.

These begin with Cenozoic reefs. J.F. Marshall (Australia), using core from the Great Barrier Reef, shows a trend for cementation to be greater in framework-dominated reefs than in sediment-dominated ones. D.M. Aissaoui, D. Buigues and B.H. Purser (France) report a study of cores from Mururoa Atoll, French Polynesia. These reveal more cementation by magnesian calcite towards the periphery with reduced precipitation, mainly as meteoric calcite spar, in the middle. The authors also give their (long awaited) data on the regional pattern of dolomitization, which includes a centripetal trend from dolomite cement to dolomite replacement. B.R. Constantz (U.S.A.) deals with the

primary surface area of corals as a control on diagenesis in Pleistocene reefs of Barbados. He provides acute observations on neomorphic changes and secondary porosity and on dissolution-precipitation fronts.

W.-C. Dullo (F.R.G.) writes on the various patterns of leaching and calcite cementation in different coral taxa in Pleistocene reefs of the Red Sea. Pore water velocity and residence time are related by R.W. Buddemeier and J.A. Oberdorfer (U.S.A.) to chemical environments in the Holocene Davies Reef, Great Barrier Reef and Quaternary reefs on Eniwetok, Marshall Islands. They emphasize the importance of substantial flow of water through the reefs combined with its chemical modification (pH, alkalinity) by bacterial activity. D.M. Aissaoui, B.H. Purser and M. Coniglio and N.P. James (Canada) record, in platform edge and talus lithofacies of Miocene age in the Gulf of Suez, Egypt, syn-sedimentary cements of fibrous and micritic calcite (now dolomitized). In the talus there are pisoids overlain by hardgrounds, and meteoric dissolution has given rise to karst. A model of dolomitization is suggested. The diagenetic diversity in Palaeocene reefs in southern Egypt is set out by J.H. Schroeder (F.R.G.). Diagenetic events include replacement with ferroan calcite and a manganese-bearing ankerite.

On the Mesozoic, P. Enos (U.S.A.) examines Mid-Cretaceous rudist reefs in northern Mexico, noting a variety of lime mudstone fabrics, some sedimentary, some cements, and provides evidence of alternation between marine and vadose meteoric diagenesis. J. Reitner (F.R.G.) makes a comparative study of diagenesis in Albian reefs in northern Spain controlled by diapirism and block faulting. A. M'Rabet, M.H. Negra, S. Sassi, N. Ben Ayed (Tunisia) and B.H. Purser have studied micrite diagenesis in Senonian rudist buildups in Tunisia. They have a useful section on the diagenesis of micrite and the problem of microsparitization. R. Koch and M. Schorr (F.R.G.) describe Jurassic sponge reefs in southwest Germany in which marine cements have been modified by meteoric diagenesis, dolomitization in mixed water and meteoric dedolomitization. Finally, Upper Triassic reefs in the Bavarian Alps southwest of Salzburg, are discussed by R. Henrich and H. Zankl (F.R.G.). These reefs contain a range of fabrics that include algal cement crusts, radial fibrous calcite and dolomites (cement and replacement).

On Palaeozoic reefs, M.E. Tucker and N.T.J. Hollingworth (U.K.) give valuable observations from a rare occurrence of undolomitized Permian reef in northeast England. They have discerned marine cements (initially aragonite and magnesian calcite), and they discuss the roles of a range of precursor fabrics on subsequent dolomitization. M. Scherer (U.K.) deals with the diagenesis of aragonite sponges in Permian patch reefs of Tunisia. He constructs a petrogenetogram on the basis of cements and internal sediments and analyses of isotopic ratios and strontium levels. A most thoughtful analysis of facies relationships and diagenesis in Waulsortian mudmounds in Ireland and England is provided by J. Miller (U.K.). He relates lateral facies changes, across flank, toe and bank, to cement sequences (calcite and dolomite) and phases of dissolution. Using isotopic ratios, H.-G. Machel (U.S.A.) traces diagenesis during burial of Upper Devonian buildups in Alberta. The changes included dolomitization of lime muds and microspar, dissolution of calcite and late burial anhydritization.

C. Kerans, N.F. Hurley (U.S.A.) and P.E. Playford (Australia) survey the diagenetic history of Devonian reef complexes in the Canning Basin, Western Australia. This is related to position in the complex and shows lateral variation in marine cementation, neptunian dikes, mechanical compaction and pressure dissolution. The distinction made

by the authors between syntaxial and epitaxial in describing cements is surely unwise since these words have previously been recognized as synonyms. K.R. Cercone and K.C. Lohmann (U.S.A.) examine the diagenetic development of some Middle Silurian pinnacle reefs in Michigan. A sequence of dolomitization and leaching is followed by recrystallization of the dolomite, some cementation with calcite and halite, then fracturing, more dissolution and dolomitization and a final calcite cementation. From a study of the cements in Silurian bioherms on Gotland, Sweden, P. Frykman (Denmark) recognizes a sequence of marine and meteoric cements related to burial and a change of pore water from marine to oxydizing meteoric to reducing meteoric.

Finally, the editors give a substantial and interesting analysis of the field in the light of the preceding papers. This reviewer was interested to see how the role of radiaxial fibrous calcite, first described in 1959, has come increasingly to the fore in studies of reef diagenesis. It is sad that workers familiar with Cullis's researches on Funafuti still neglect the comparable study by E.W. Skeats (Bulletin Museum of Comparative Zoology, Harvard College, vol. 42, 53-126, 1902). Regrettably, the precise nomenclature of Folk has become blurred so that now sparry, sparitic and spar are all synonymous adjectives and the easy noun "spar" appears to be obsolete.

This book is well organized, attractively set up and well illustrated with a reasonable index. The overall quality is high, as is to be expected from the many internationally respected workers among its contributors. A review cannot adequately express the richness of fascinating and valuable detail in these studies. For those engaged in research work or advanced university teaching the book is an essential companion.

R.G.C. Bathurst
Clwyd, North Wales, U.K.

PALAEOBIOLOGY OF CONODONTS

Richard J. Aldridge (ed.)

John Wiley & Sons, New York, Chichester, Brisbane, Toronto, 1987, 180p., \$79.95 (Cdn.)

CONODONTS: INVESTIGATIVE TECHNIQUES AND APPLICATIONS

Ronald L. Austin (ed.)

John Wiley & Sons, New York, Chichester, Brisbane, Toronto, 1987, 422p., \$89.95 (US)

Conodonts are small tooth-like phosphatic microfossils from marine strata of Cambrian to Triassic age. These microfossils are very useful in studies of biostratigraphy, and lately, thermal maturation and trace-element geochemistry. A major debate has centered on the function of the conodont and the nature of the conodont animal - the conodont has been an orphan in search of a Phylum.

Until recently, models of conodont function and the conodont animal have suffered somewhat from the lack of extant or fossil conodont-bearing animals. In 1983, a paper published in *Lethaia* illustrated a single, small (about 4 cm in length) worm-like animal from Lower Carboniferous strata near Edinburgh (Episodes, v. 1983, no. 2, p. 28). At one end

of this body are a cluster of conodonts. The initial, cautious assignment of the conodont animal to its own, extinct Phylum has been replaced by a more detailed study of the original and three new specimens published in late 1986, which suggested that the conodont animal was a chordate.

The two multi-authored volumes under review here are largely made up of papers presented at the Fourth European Conodont Symposium (ECOS IV) held at the University of Nottingham in 1985. The first seven of eleven papers of *Palaeobiology of Conodonts* are concerned with the conodont animal and its mineralized elements, the Conodonts. In spite of the discovery in Scotland, these papers show a continuation of the interesting and stimulating debate on the nature of the conodont animal. In this volume, various authors argue that conodonts are related to: (1) chaetognaths - "arrow worms," (2) cephalo-chordates - "amphioxus," or (3) myxinoids - "hagfish." The elements function is suggested to be a bilaterally operative lingual apparatus or, perhaps as a more passive, tissue-covered food gathering device.

The lead paper by R.J. Aldridge (U.K.) presents a good historical view of conodont palaeobiology including the composition, structure and function of the conodont elements, the nature of the conodont animal, and the architecture of the conodont apparatus. Interpretations of these apparatuses are based in part on natural clusters of elements, whether found on bedding planes, as in the Scottish animal, or as a fused cluster. The latter form is used in a study by R.S. Nicoll (Australia) and C.B. Rexroad (U.S.A.) of a Silurian conodont genus from northern Indiana to show that its bilaterally symmetrical apparatus contained fifteen elements comprising seven distinct forms.

R.J. Aldridge, M.P. Smith (U.K.), R.D. Norby (U.S.A.) and D.E.G. Briggs (U.K.) noted that bedding plane assemblages occur in a limited number of recurrent patterns, and that all of these can be formed by collapsing and flattening one three-dimensional structure. Their method was to photograph clay models of a conodont cluster to produce a two-dimensional view of a three-dimensional subject. Their interpretation of the functional apparatus is that the anterior elements grasped food, which was then processed by a shearing and grinding action of other elements.

Soft tissue of Silurian conodonts from Sweden was examined in a preliminary report by L.E. Fahraeus and G.E. Fahraeus-van Ree (both of Canada) on the decalcification of the conodont mineral. The histological staining of the freed tissue revealed variously shaped cells, collagen and lumina. The last four papers follow the evolution of conodonts from Early Ordovician time through to their "final fifty million years" in the Late Triassic.

The first seven of twenty-three chapters of *Conodonts: Investigative Techniques and Applications* deal with laboratory methods of conodont processing. To the specialist, the conodont is "ore" and the surrounding rock is "gangue," and these chapters are worth reading by anyone interested in techniques of rock disintegration and heavy mineral separation. For example, J. Stone (U.K.) reviews thoroughly the basic methods of sampling, rock digestion, sieving and concentration of the residues, and picking, mounting and photographing conodonts. The next two chapters are collections of suggestions by a number of contributors for improving or simplifying the task of conodont preparation. The first of these is on rock disintegration methods, the second on conodont concentration methods. To avoid the use of toxic heavy liquids, other techniques are discussed such as the interfacial method (which exploits the lipophilic properties of the conodont mineral apatite), cryogenic density, electromagnetic, and magnetohydrostatic separation.

Book Reviews

M.J. Orchard (Canada) and J.E. Barrick (U.S.A.), in separate papers give their recipes for recovering conodonts from chert using hydrofluoric acid. The former author describes conodonts from his 100 samples of deep-water chert from western Canada, which range from Carboniferous to Triassic in age and are differentiated into eight succeeding faunas. This is the first step to form a chert-based conodont biostratigraphic scheme for the tectonostratigraphic terranes of this region. Other biostratigraphic studies in this volume involve conodonts from the Ordovician of Wales and the Welsh Borderland, and from the Carboniferous-Permian of Bolivia.

Two chapters review techniques of "photography": R.D. Burnett (U.S.A.) and M.P. Smith (U.K.) concentrate on scanning electron microscopy (secondary and back-scatter electron imaging), and R.D. Norby and M.J. Avcin (U.S.A.) illustrate contact microradiography of conodont assemblages in shale. N. MacLeod and T.R. Carr (U.S.A.) co-author a chapter on quantitative methods of morphological analysis that can potentially contribute to conodont taxonomy, biostratigraphy, palaeoecology and phylogenetics.

Organic metamorphism is the theme of three papers. G.S. Nowlan and C.R. Barnes (Canada) discuss the application of conodont CAI (colour alteration index) to hydrocarbon and mineral exploration, and regional tectonic interpretations. They also use the CAI to trace hotspots. The apatite of conodonts seems to preserve original marine isotopic and trace-element signatures. J.E. Geitgey and T.R. Carr (U.S.A.) compare the diversity and distribution of conodonts with the isotopically inferred palaeotemperatures and find some correlation. For example, diversity and/or distribution of conodonts was widespread when temperature was increasing or relatively high. J. Wright, J.F. Miller, and W.T. Holser (U.S.A.) use variation in trace element signatures from apatite of conodonts and inarticulate brachiopods to correlate strata of the Cambrian-Ordovician boundary interval. The volume ends with four papers on a palaeoecological theme.

Both volumes have taxonomic and general indexes, and the plate quality is quite acceptable. There are, however, typographical errors, and in the techniques volume there is at least one incomplete sentence and missing reference. These errors are annoying when considering the price of the volumes. However, both these books and their companion volume published in 1985, *A Stratigraphical Index of Conodonts* (which should be more accurately titled "Conodonts of Britain and Ireland") are recommended to conodont workers. New students of conodonts will find the summary chapters especially useful. *Palaeobiology of Conodonts* also may be of interest to vertebrate and invertebrate biologists, and the second volume with its descriptions of techniques should find a broad readership in other fields of geology. It is unfortunate that some potential buyers of these two volumes (and the earlier one, which has a comparable price) may find the cost exorbitant.

A.D. McCracken
Ottawa, Canada

GEOLOGY AND OFFSHORE RESOURCES OF THE SOUTHWEST PACIFIC

The Circum-Pacific Council for Energy and Mineral Resources (CPCEMR) recently published three volumes on the Southwest Pacific, which are here reviewed together. The books report the results of multinational studies sponsored by the Governments of U.S.A., Australia and New Zealand, coordinated by CCOP/ SOPAC (Committee for the Coordination of Joint Prospecting for Mineral Resources in South Pacific Offshore Areas).

VOL. 2 GEOLOGY AND OFFSHORE RESOURCES OF PACIFIC ISLAND ARCS - TONGA REGION

David W. Scholl and Tracy L. Vallier (eds.)

CPCEMR, Earth Science Series, Vol. 2, AAPG, Oklahoma, 1985, 488p., \$28 (US)

This collection of 35 papers reports the results of the 1982 cruise of the USGS Research Vessel S.P. Lee from Eastern Samoa along the Tonga Ridge to Tongatapu and a transit across the Lau Basin and Ridge to Fiji, together with data from subsequent laboratory studies. The volume also includes a number of invited or commissioned papers discussing new or largely unpublished information covering other aspects of the geology of the region.

Part I contains an introduction summarizing the geological and tectonic setting and providing maps of the submarine topography and block drawings of the perspective physiography. Five short papers make up Part II with descriptions of the shipboard data-collection operations, which comprised 4000 line km of single-channel seismic profiles plus gravity and magnetic data; 2400 km of multichannel seismic (MCS) profiles, 54 wide-angle sonobuoy refraction measurements and 9 dredge hauls. Gravity coring proved unsuccessful.

Part III constitutes the major part of the volume and comprises three sections on the geology and geophysics of the southern Tonga platform, the remainder of the Tonga Ridge and adjacent areas respectively.

The structure and basin analysis of the southern Tonga forearc based on MCS profiles is discussed by R.H. Herzer and N.F. Exon, who present structure and isopach maps for several horizons and trace the history of the area since the Eocene arc volcanism. Total sediment thicknesses reach over 4 km in several places. Eight papers record studies of samples from eight dredge hauls from between 600 and 3750 m depth on the southern Tonga platform, and one from 7800-8600 m on the western wall of the Tonga Trench. In general, the platform is surfaced by westerly dipping and easterly thinning sediments on an igneous basement that is exposed on 'Eua. Planktonic foraminifera indicate that outcropping materials on the platform are Late Miocene or younger (G.C.H. Chaproniere), while coccoliths indicate generally shallow-water deposition during this period (D. Bukry). Geochemical studies show that these sediments contain organic material of only low maturity, whereas oil

REVISTA BRASILEIRA DE GEOCIÊNCIAS

IF YOU INTEND TO LEARN MORE
ABOUT BRAZILIAN GEOLOGY AND ITS
RELATIONS TO SOUTH AMERICAN
CONTINENTAL EVOLUTION,
YOU SHOULD RECEIVE THE REVISTA
BRASILEIRA DE GEOCIÊNCIAS.

Revista Brasileira de Geociências is the Brazilian journal covering all the fields of the geosciences. New advances in geological research constitute the main body of each quarterly issue. The papers are from Brazilian and/or foreign geoscientists, revealing the amount of institutional cooperation and the level of scientific research that has been made at the present time in the country and adjacent regions.

The issues are complemented by abstracts of academic theses, conference reports, notices, new publications and book reviews, debates and comments, news and lists of coming events. Excursion guides to the most spectacular geological features of different parts of Brazil are to be included in the current volume 17, 1987.

The editors intend to expand the international distribution of the Revista. Subscriptions and exchange agreements are strongly encouraged.

RECENT RESEARCH ARTICLES

Post-paleozoic magmatism in Brazil
Classification system of Brazilian sedimentary basins
Transcurrent tectonics in petroleum exploration
Origin and evolution of a part of Goiás State

FORTHCOMING ISSUES

Evolution of the South Atlantic Ocean
Proceedings (21 articles) of a Symposium on the subject, comprising tectonics, stratigraphy, paleontology, paleogeography, evolution and filling of South Atlantic basins.

Granites and associated mineralizations
Proceedings (about 46 papers) of the International Symposium on the Granites and Associated Mineralizations (ISGAM) held in Salvador, January 1987.

I enclose \$36 US (1988 rates) for one year subscription of the Revista Brasileira de Geociências. The check must be made out to "Sociedade Brasileira de Geologia."

Name: _____

Address: _____

City: _____

Province: _____ PC: _____

Country: _____

The Revista is maintained by the Sociedade Brasileira de Geologia and sponsored by governmental agencies: DNPM - Departamento Nacional da Produção Mineral, CNPQ - Conselho Nacional de Desenvolvimento Científico e Tecnológico, and FINEP - Financiadora de Estudos e Projetos. The total distribution is 5,000.

It is air-speeded to subscribers outside Brazil.

Mail order form to:
Revista Brasileira de Geociências
Caixa Postal 20897
CEP 01000
São Paulo, SP, Brazil

Book Reviews

seep material in the Tonga region is from thermally mature source rocks, presumably from deep on the west part of the platform (M.W. Sandstrom; B. Buchbinder and R.B. Halley).

The volcanic debris on the southern Tonga platform is material of low-K island-arc tholeiite and minor calc-alkaline affinities, and of basic, intermediate and acid types; the post Miocene material was probably derived from the Tofua Ridge (P.A. Cawood; J.W. Hawkins). Ferromanganese crusts on some samples, collected from 600-3000 m water depths, are up to 1.5 cm thick, and mainly hydrogenous in origin, compositionally similar to abyssal Pacific nodules with low Co. The dredge haul from the Tonga Trench contained serpentized harzburgite as well as sediments with a Miocene fauna and silicic volcanic debris (T.L. Vallier et al.).

The final three papers in this section are on geophysical studies using gravity and magnetics (C. Alexander) and MCS and sonobuoy data (R.M. O'Connor) in study of a domal structure, and on problems of near surface velocity inversion on processing MCS data (J.E. Cole and D.M. Mann). Exon, Herzer and J.W. Cole point out that MCS records indicate two widespread unconformities in the sediments, in the Early Oligocene and Early Pliocene, as well as the unconformity of sediments on the presumed Eocene basement.

The "Insular and Subsurface Geology, Tonga Ridge" is dealt with in nine papers. They include a summary of the geology of all the Tonga islands (J.K. Cunningham and K.J. Anscombe). Various studies of the volcanism show that both the young Tofua Arc and the older eastern arc represented by 'Eua are formed of normal intra-ocean island arc tholeiite types, although 'Eua has some spreading-centre affinities. Planktonic forams and coccoliths from island samples indicate that the eastern belt volcanism during Miocene time was in an outer shelf depositional environment, whereas samples dredged from the Tonga Trench imply a subsidence of the forearc since the Middle Miocene of 3 km or more (P.J. Quinterio). The rate of subsidence decreased with time (G.H. Packham) until the onset of Lau Basin spreading led to uplift in Early Pliocene time. Subduction of the Louisville Ridge during the past 3.5 Ma has resulted in a terrain of domed and tilted fault blocks in the northern part of the Tonga Ridge, with some important transverse faults (J. Dupont and R.H. Herzer).

The general geology of the Lau Ridge is described by D. Woodhall who stresses the dominance of NE- and NW-trending fault and volcanic alignments. There is more petrological variation in the larger Fiji Islands than on the Lau Ridge, but their histories are generally comparable, with chemically distinct mature arc, incipient rifting and remnant arc stage volcanics (J.W. Cole et al.; P.M. Whelan et al.). J.L. Morton and N.H. Sleep document seismic reflections from a magma chamber beneath the Valu Fa Ridge in the southern Lau Basin. Some island samples from Fiji do show some gas and some oil generation potential (B. Buchbinder and R.B. Halley), though none of the Tonga samples do - a situation in agreement with Sandstrom's findings on the offshore Tonga samples.

Part IV of this volume is a good synthesis by D.W. Scholl, T.L. Vallier and G. Packham of the geological framework and resource potential of the southern Tonga platform and adjacent areas derived from the earlier papers and previous work. Despite the negative results on the Miocene samples, the authors conclude that there is potential for hydrocarbon accumulation in Lower Tertiary rocks, especially on the west side of the Tonga Ridge, or possibly on the now

separated eastern part of the Lau Ridge. There is also some potential for mineralization associated with the volcanic centres, although at present this is largely unproven.

Over the past few years there have been cruises by research vessels from several countries in the Tonga and Lau region, which provides us with, perhaps, the best and simplest example of back-arc spreading in a young volcanic arc. Nevertheless, the cruise tracks remain few in number and all forms of sampling, geological, geochemical and geophysical, are rather sparse. This volume does not set out to provide a complete statement of the geology and mineral resources of the Tonga-Lau region, because the Lau Basin and the north Tonga Ridge are largely ignored. It does, however, provide us with a great deal of new information and the most complete picture currently available of the southern Tonga platform and the Tongan and Lau islands. It will consequently be an essential reference for future studies of the region, especially when complemented by the studies of the northern part of the Tonga Ridge and Lau Basin arising from the recent cruises of the RV *Natsushima* and RV *Sonne*.

VOL. 3 INVESTIGATIONS OF THE NORTHERN MELANESIAN BORDERLAND

Thomas M. Brocher (ed.)

CPCEMR, Earth Science Series, Vol. 3, AAPG, Oklahoma, 1985, 199p., \$24 (US)

This volume consists of 10 papers summarizing the results of an investigation conducted by scientists aboard the RV *Kana Keoki* (Hawaii Institute of Geophysics) during a 20-day cruise in April 1982 from Western Samoa to Fiji during which 4 piston cores, 4 free-fall cores, 14 rock dredges and 14 sonobuoy profiles were collected. The area concerned is part of the boundary zone between the Pacific and Australasian plates, covering the northeastern part of the North Fiji Basin, the Vitiaz Trench and the Samoan Islands.

A useful introduction by T.M. Brocher and R. Holmes briefly reviews the geological framework of the area, with an overview of the objectives of the cruise and of the results reported in the papers that follow. Brocher presents evidence that the Vitiaz Trench lineament was initiated by "easterly" subduction of the Pacific Plate beneath the North Melanesian Borderland, and that such subduction was abandoned progressively from west to east. The original trench morphology has been somewhat distorted by the collision with the Samoan seamount chain. On the North Fiji Basin, subduction-related volcanism continued into the Pleistocene (J.M. Sinton, K.T.M. Johnson and R.C. Price), long after reversal of subduction had occurred farther west. North of Fiji a new spreading ridge, trending N71°W for a short time, now appears to be inactive (Brocher).

On the basis of 11 new rock dredges, Sinton, Johnson and Price were able to define the volcano-tectonic provinces of the region. These include the low-K tholeiites of the Lau and North Fiji basins, typical of spreading centres, and arc tholeiites of the Horne Island. On the north side of the

Book Reviews

Vitiaz Trench lineament, the Samoan chain shield-building tholeiites and alkalic lavas have been extended westwards. However, the Eocene ocean island tholeiites of Alexa Bank do not fit a fixed hotspot model applicable to other volcanoes of the Samoan chain.

Regional sedimentation patterns (Brocher et al.) show the dominance of deep-sea fans with little deposition on the narrow shelves of the volcanic arc. The young North Fiji Basin is significantly poorer in volcanoclastic sediments than elsewhere. Sedimentation around the Samoan seamount chain is rich in reef-derived detritus, partly redistributed by Antarctic bottom waters. The other parts of the Pacific Plate have received sediments by ash-fall and turbidity currents from remote topographic highs. Microfaunal studies have allowed a refining of the local zonation of the Late Pliocene and Quaternary strata and indicate that water depths and oceanographic conditions have generally remained stable during that time, although Tuscarora Bank has subsided from about 100 m to its present depth of 1060 m since the Early Pliocene (G.C.H. Chaproniere).

Small sedimentary basins south of Viti Levu contain three layers of sediment separated by two unconformities; the latter probably record two separate rifting events, the first occurring with the initial fragmentation of the Vitiaz Arc during the Late Miocene/Early Pliocene, and the second associated with formation of a new spreading ridge just west of Viti Levu (Brocher and R. Holmes). Both events contributed to the anti-clockwise rotation of the island of at least 30°. Several of the basins contain over 1 km of sediments, but their hydrocarbon potential is unknown. Though

the data are still sparse, radiometric ages (J.H. Natland and D.L. Turner; R.A. Duncan), foraminiferal ages (Chaproniere), and geophysical data including palaeomagnetic reversal history (B. Keating) indicate a westerly increase in age of the Samoan chain, consistent with a fixed hotspot origin.

The studies reported in this volume add significantly to our knowledge and understanding of the complex boundary between the Pacific and Australasian plates in the North Melanesian Borderland region, and provide a good starting point for further studies. Few research ships have crossed this area, however, and the number of samples studied is still too small to allow a complete picture of the development of the region. In particular there remain question marks over the affinity of the oceanic islands westward of the Samoan chain and over the precise effects of the collision of this chain with the Australasian Plate and the deformation of the Vitiaz Trench. There are also uncertainties about the development of the northern part of the North Fiji Basin and, in particular, about the reality and nature of the widely quoted, but unproven, Fiji transform fault.

Vol. 4 GEOLOGY AND OFFSHORE RESOURCES OF PACIFIC ISLAND ARCS - CENTRAL AND WESTERN SOLOMON ISLANDS

J.G. Vedder, K.S. Pound, S.Q. Boundy (eds.)

CPCEMR, Earth Science Series, Vol. 4, AAPG, Oklahoma, 1985, 306p., \$28 (US)

This volume reports the results of an offshore survey in 1982 by a multinational research team aboard the S.P. Lee to investigate the structure and stratigraphy of the Central Solomons Trough, a major intra-arc sedimentary basin. The book contains a 25-pages introduction by J.G. Vedder and F.I. Coulson on the geomorphology, geology and tectonics of the region, with a review of previous offshore and onshore research. A section on "Shipboard Operations" comprises a number of short papers describing the geophysical and bottom sampling techniques employed by the survey. Included is a series of bathymetric maps and physiographic diagrams by T.E. Chase, B.E. Seekins and K.E. Lund, which illustrate the main submarine topographic features of the region.

The main body of the volume, on "Topical Studies" describes the onshore and offshore geology of the region, including the presentation and interpretation of a wealth of new information. There are background reviews of the geology of the islands (Coulson and Vedder), correlation of rock units (K.S. Pound), and the regional offshore geology (Vedder and Coulson). Together these provide a good synopsis of the geology of the region, and draw upon information obtained over many years by the Solomon Islands Geological Survey, which has not previously been widely publicized. Succeeding papers cover specific aspects of the geology, and there are noteworthy contributions based largely upon new geophysical data.

NEW from SWIFT (UK) MODEL "F" H/I AUTOMATIC POINT COUNTER

is a considerable advance over previous models. Now, with 12 channels -- one isolatable from total: Employs microprocessor for compactness, optimum reliability and extra facilities -- such as percentage key and up to 9x interval multiplier: Separate LED displays for channels, and for total plus combined 4-digit preset -- with audible signal.

Point counting is not limited to polarizing microscopes.

Attachments for all well-known microscopes.

H/I HACKER
INSTRUMENTS
INC.

Box 657, Fairfield, New Jersey 07007
(201) 226-8450

Book Reviews

L.A. Beyer presents and discusses a new and much needed gravity-anomaly map of the region. P.A. Cooper, L.W. Kroenke and J.M. Resig describe, in two separate papers, the shallow and deep crustal structure as interpreted from seismic and gravity studies. The seismic stratigraphy and structure of major intra-arc sedimentary basins are discussed by T.R. Bruns and others, who present a multichannel seismic-reflection profile across the arc. The following papers by J.B. Colwell and colleagues describe dredged-rock and gravity core samples collected during the cruise, and recent depositional patterns within the Central Solomons Trough. Resig reviews the foraminiferal biostratigraphy and palaeobathymetry of dredged bottom samples, which indicate that the region has been subject to large and rapid vertical movements during the Pliocene and Quaternary, confirming conclusions reached from study of the islands.

Resig and others also discuss the palaeobathymetry of sediments exposed on Malaita and neighbouring islands, as deduced from depth controlled benthonic foraminifera. They report substantial uplift along the northeastern flank of the arc during Pliocene and Quaternary time, with rates of uplift as high as 2,600 m/my.

The final two papers of the section are related to the hydrocarbon potential of the region. Buchbinder and Halley report that preliminary source-rock evaluation of onshore samples gave disappointing results indicating that the little organic material present is immature; however, the collected samples represent only a very limited portion of the sedimentary column of the Solomons. In discussing the offshore petroleum potential, Vedder and Bruns suggest that suitable reservoir rocks and traps may exist within the Central Solomons Trough.

In the final part of the volume, Vedder summarizes the geology of the region and the results of the cruise. Final conclusions are that the major sedimentary basin of the Central Solomons Trough owes its origin to the reversal in the arc polarity during the Miocene Epoch, and that its thick sedimentary fill initially accumulated as a back-arc apron to a northeast-facing Oligocene-Miocene volcanic arc, and later as a back-arc apron related to the currently active southwest-facing arc. Although source rocks for the generation of hydrocarbons were not positively identified, structural and stratigraphic traps may exist within the basin.

The papers in this book on the offshore geology provide a great deal of new and factual information where little previously existed, and the various background papers provide an excellent and balanced introduction to the geology of the Solomons arc. In contrast, the content related directly to offshore resources is minimal, being confined to a brief report on the source-rock evaluation of a small number of onshore samples and one paper outlining potential areas for offshore hydrocarbon accumulation. Despite this, many of the papers will be invaluable to any future hydrocarbon exploration within the region, and in this respect the volume largely fulfills the original aims of the survey. In a wider context the contributions provide a good insight into the geology, evolution and geophysical structure (but not the petrology) of a complex island arc system. The volume will, therefore, be of general relevance to researchers interested in both ancient and modern destructive plate-margin environments.

Because these three volumes are collections of papers rather than planned compilations, there are inevitably repetitions of figures and ideas and the standard of printing, particularly of volumes 2 and 3, could be better. The binding and backing can also be criticized for being too flimsy to

stand up to the widespread consultation that these books deserve and are sure to get. However, they incorporate a wealth of new information that will be of great interest to scientists working in the complex and geologically exciting environment of the Southwest Pacific.

D.I.J. Mallick, P.N. Dunkley,
J. Ridgway and R.T.R. Wingfield
Nottingham, U.K.

SEDIMENTATION IN THE AFRICAN RIFTS

L.E. Frostick, R.W. Renaut, I. Reid and J.J. Tiercelin (eds.)

Geological Society of London Special Publication 25, 382p., 1986, (C) \$88 (US)

This book presents the results of a symposium held by the Geological Society of London in September 1984. An attempt was made to provide a comprehensive coverage of the development of the rifts and, indeed, the editors have been so successful in meeting their objectives

that the title suggests a rather more limited range of contents than in fact the book contains. Tectonism and volcanism received considerable discussion, which is appropriate, for these two controls have a major effect on sedimentation.

The editors state that considerable interest in the symposium was expressed by petroleum geologists, because the African rifts can be used as analogues for several significant petroleum accumulations, in the North Sea, Libya, and elsewhere. Although there is little in this book that directly addresses the interests of the petroleum community, the background documentation of this huge, continent-wide case example will undoubtedly prove invaluable. Another of the major interests in the rifts is the superb record of humanoid evolution preserved in their sediments. The special reasons for this are examined in several papers, although the evolutionary story itself is not dealt with in this book.

There is a useful introduction by H.G. Reading, who discusses the African Rift System in a global context, provides a brief review the variable style of tectonism and volcanism along its different segments, and examines the controversy as to whether the rifts are "active" (driven by mantle upwelling) or "passive" (a consequence of plate kinematics). A.T. Grove summarizes the present surface expression of the rifts, the drainage patterns through and across them, and the huge, broad domes that accompanied the development of the rift system, the largest of which corresponds to the sites of triple points in the oceanic spreading centers.

The next section of the book deals with the tectonism and volcanism of the rift system, and the relationship of these processes to sedimentation. J.D. Fairhead discusses gravity models of the lithosphere based on several transects across the West African Rift System and two transects across the Kenya Rift. Petroleum company seismic data are used by B.R. Rosendahl and coworkers to develop a tectonic model for the Lake Tanganyika Rift. Their surprising discovery is that very little of this rift consists of true two-sided graben. Half graben are the dominant structures, each bounding fault having a curved outline in plan view. The rift basin appears to comprise a chain of these half graben facing in alternate directions, and the authors suggest that this may represent a general pattern for rift faulting.

Book Reviews

B.H. Baker, L.A.J. Williams, G.R. Chapman, and R.T. Watkins examine in three separate papers the details of the relationships between volcanism and sedimentation in different parts of the Kenya Rift. This is far from being a simple story because volcanism is accompanied by faulting and regional doming, which affect the position of lake shorelines and the direction of tilt of paleoslopes. C.J. Swain, in a related study, uses gravity and seismic data to investigate the basin geometry below part of the sediment wedge east of Lake Turkana in Kenya.

The third section describes sedimentary processes in the East African Rifts, mainly the Kenyan segment. L.E. Frostick and I. Reid contribute two papers dealing with fluvial-deltaic processes and resulting coarse clastic sedimentation, the first discussing slope processes, and the second the development of the lake deltas emptying sediment into Lake Turkana. A.S. Cohen and co-workers discuss the effects of faults and volcanic barriers on the distribution of coarse sediments in this lake basin.

The arid nature of the climate of East Africa, and the enclosed nature of the lake basins, leads to unusual water chemistries and consequent chemical sedimentation. This is discussed by R.F. Yuretich (clay minerals), P.L. Abell and J. P. McClory (sedimentary carbonates), R.W. Renaut and co-workers (sedimentary diagenesis) and H.P. Eugster (hydrochemistry of Lake Magadi). Abell and McClory attempt to use stable isotope geochemistry to "type" carbonate sediments for use as stratigraphic marker beds. This could be an extremely useful aid to detailed chronostratigraphic correlation of hominid-bearing beds. Eugster's paper presents no new data, but reviews comparisons between Lake Magadi and ancient Lake Gosiute, in which the Green River Formation of the Rocky Mountains region was deposited. R.

Crossley describes an unusual type of sedimentary process, the production of homogeneous clayey sand sheets by the vertical transfer of sedimentary particles by termites. This section ends with papers on stromatolites by J. Casanova, and peat development by A. Hamilton and D. Taylor.

The stratigraphic history of various parts of the rift system in Ethiopia, Kenya and Tanzania is then reviewed in nine papers that provide a paleogeographic and paleoclimatic framework for the hominid discoveries for which these basins are famous. Much local detail is presented, and the three final papers deal with the preservation of fossil faunas. R.L. Hay picks up on a theme developed by the late W.W. Bishop, that "rift basins are a good place to die," because the carbonatite volcanic ash that is abundant in parts of the East African Rift System both helps bury fossil material and provides a suitable chemical environment for preservation. M. Pickford summarizes the variations in depositional environment in which different faunal assemblages are found, and C. Denys and co-workers examine the paleoecology of rodents.

This is a very thorough and comprehensive review of one of the world's unique geological provinces. There is far more information here than the general reader would wish to digest, but a substantial amount of important information and many useful ideas for those engaged in the study of other ancient or modern rift basins. Anyone with a professional interest in African geology will probably want to purchase a copy. The quality of production is excellent, and the price, though rather high, is not out of line for this specialized type of book.

A.D. Miall
Toronto, Canada

CONTRIBUTIONS TO THE GEOLOGY AND HYDROCARBON POTENTIAL OF THE EROMANGA BASIN

Editors: D. I. Gravestock, P. S. Moore & G. M. Pitt.

Price: Aus\$ 40.00 Surface mail
Aus\$ 54.00 Airmail

Chapters on:

- An Exploration overview of the Eromanga Basin
- Historical Perspective
- Regional Basin Studies
- Structure and Tectonics
- Petroleum Reservoirs
- Source Rock Geochemistry and Organic Petrology
- Hydrocarbon Generation and Migration

1986

SPECIAL PUBLICATION NO. 12

Geological Society of Australia Inc.

10 Martin Place, Sydney N.S.W. 2000 Australia