East Asian Palaeoenvironments

Hong Kong, 9 - 13 January 1987

The first Conference on the Palaeoenvironment of East Asia held in 1983 by the Centre of Asian Studies at the University of Hong Kong was followed recently by a second conference under the same auspices. Specialists from eleven countries delivered some 60 papers in seven sessions, each of which was opened by a leading Chinese scientist who summarized major recent advances in that field in China.

Geology

The Cenozoic development of Hainan Island was reviewed by Huang Yukun (P.R.C.). He concluded from an analysis of stratigraphy, tectonics, and volcanism that initial rifting took place here in the Eocene, followed by extension and subsidence in the Miocene, depression in the Pliocene, break-up and closing during the early mid-Pleistocene and, finally, by differential block-faulting and overall uplift after the mid-Pleistocene. Edward Derbyshire (U.K.) showed how granulometry and fabric of loess and desert lacustrine sediments from eastern Asia reflect processes of grain derivation, transportation and deposition. Superimposition of deformation and diagenetic fabrics on the primary depositional fabrics provides information on changing palaeoenvironments since initial deposition.

Monique Fort (France) noted, from work in the Pokhara basin along the Himalayan front, that distinction between glacially and tectonically induced deposits is crucial before drawing conclusions about climatic variations. Recognition of catastrophic morphogenesis may also be useful for the reconstruction of palaeontological succession. V.J. Gupta (India) then described the Quaternary biostratigraphy of the Nepal Himalaya, where up to 500 m of Plio-Pleistocene lacustrine, fluvial and fluvio-glacial sediments occupy a

synclinal depression in the Kathmandu Valley. He correlated the faunal and vegetational sequence with those in the Kashmir Valley. Another correlation using magnetostratigraphy by M. Yoshida and colleagues (Japan) linked the Churia (Siwalik) Group of Nepal, with the Pakistan Siwaliks (Fig. 1).

Sea Levels and Coastal Changes

The complex history of eustatic changes, climate variations, and tectonic movements along the coasts of north and east China were reviewed by Yang Huai-jen (P.R.C.) on the basis of sedimentological and micropalaeontological data. He recognized five major Cenozoic sea level cycles, with Eocene, Pliocene and Early Pleistocene transgressions reaching far inland, and eight to ten transgressions occurring during the Quaternary. On the basis of palaeogeographic and sedimentary evidence, Duan Weiwu (P.R.C.) described the rapid changes that have taken place in the northern margin of the South China Sea since the Late Oligocene. Microfossils permit correlation between the various basins of the region and testify to three transgression-regression cycles: Late Oligocene to earliest Early Miocene, medial Early Miocene to mid-Miocene, and Late Miocene to Pliocene.

In a related work Huang Jinsen (P.R.C.) classified extensive coral reefs of the South China Sea into four ecological categories. Fringing reefs predominate over three sedimentary facies: reef front, reef core, and back reef. He showed how these reefs are affected by geological structure, sedimentation, biogenic processes, cementation, erosion, marine currents and storm tides, changes in sea level and by human activity.

Palaeoclimatology

Opening this session, Wang Pinxian (P.R.C.) remarked on the multidisciplinary approach which had characterized the 18 symposia on palaeoclimatology held in his country since the first Hong Kong conference. Major attention has focussed on the Quaternary, particularly on sea level changes and transgressions, followed by palynology and palaeobotany, glacial and periglacial studies and regional or areal syntheses. The evolution of physical regionalization of China since the early Tertiary was then analyzed by Zhao Songqiao. According to him, the monotonous, peneplained topography and warm, homogeneous climate with low levels of areal differentiation of the early Tertiary were replaced by complex climatic zones following the Himalayan tectonic movement and the onset of the monsoon system.

Wu Xiangding described climatic fluctuations in Tibet since the Late Pliocene when it was at an altitude of 1,000 m, through four glacial and three interglacial periods during the Quaternary, with an overall trend towards decreasing temperature and precipitation accompanying orogeny. The progressive intensification of aridity in north China since the mid-Pleistocene was then discussed by Zhang Liansheng. Widespread traces of glacier deposits in west China and extensive development of lakes in the Qinghai-Xizang Plateau, Xinjiang and Inner Mongolia testify to the occurrence of a wetter period during the mid-Pleistocene. The southern

Figure 1: Magnetostratigraphic correlation between the Siwaliks of Nepal and Pakistan. From M. Yoshida et al., in press (Conference Proceedings).

Conference Reports

limit of loess reached the lower Yangtze basin during the Late Pleistocene. Finally, S.K. Gupta (India) reviewed climatic changes in northwestern India over the past 700,000 years, noting that while the ocean cores indicate about 18,000 years as a glacial maximum, palynological evidence shows that deglaciation had started here well before this period.

Palaeobotany

Sun Xiangjun (P.R.C.) pointed out that greater attention had been paid in recent years in her country to Quaternary palynology than to Tertiary palaeobotany. However, Guo Shuangxing (P.R.C.) described megafossils from Eocene to Pliocene strata in the Hengduan Mountains, western Sichuan, that reveal a change from a sub-arid, sub-tropical climate at 300-400 m in the Eocene through a humid, sub-tropical climate in the Oligocene and Miocene; the present elevation is 3,984 m. Megafossils collected during the Qinghai-Xizang Scientific Expedition were then used by Tao Jun-rong (P.R.C.) to describe the Eocene flora from the Liuqu Formation along the Yarlung-Zangbo River, indicating a hot, humid climate in or near tropical rainforest.

R.N. Lakhanpal (India) examined the geological evolution and palaeogeography of northern India in order to detect the time at which temperate elements first entered the region. He concluded that the final withdrawal of the Tethys during the Oligocene, followed by further Himalayan orogeny, created conditions favourable to the immigration during the Middle Miocene of temperate elements from mainland Asia. In contrast, Huang Tseng-cheng (Taiwan) pointed out that since the Oligocene, most fossil genera have corresponding modern counterparts, and the six modern vegetation types have changed little over the period. Vegetational and climatic changes at the Pleistocene-Holocene transition in Yunnan were discussed by D. Walker (Australia). Shortly before the last glacial maximum, the climate at Menghai was similar to that of today, though wetter in winter and less inclined to frosts or spring droughts. At Kunming, rainfall was higher than today, with mean annual temperatures only 1-2.5°C lower, the modern pattern becoming established around 7,000 yBP.

▲ "Megamacaca"

○ Middle Pleistocene *Rhinopithecus*

● Late Pleistocene *Rhinopithecus*

Figure 2: Distribution of living *Rhinopithecus* (solid black) and fossil colobines in China. After Jablowski and Pan, in press (Conference Proceedings).

Invertebrate Palaeontology

Wang Pinxian told the conference that the focus on invertebrate palaeontology in China was on microfauna rather than on megafauna, with greater attention being paid to distribution and applications than to taxonomy. For example, Liu Tungsheng and Gao Fuqing described the value of fossil snails from the Luochuan loess as indicators of palaeoclimate. Two assemblages are discerned, the first composed of cold-dry types, reflecting a desert-steppe environment, and the second, from the palaeosol layers, warm-humid types, reflecting a forest-steppe or steppe environment.

Wang Pinxian described work on foraminifera, geochemistry and palaeoceanography based on cores from the northern part of the South China Sea over the last 130,000 years. Five oxygen isotope stages are distinguished on the $\delta^{18}O$ curve of *Globigerinoides sacculifer*. Considerable variations in water surface temperatures are attributed to changes in water circulation during the glacial period and to an estuarine circulation system that may have caused oxygen deficiency in bottom water. Wyss Yim (Hong Kong) then showed that the distribution of Holocene ostracoda from nine boreholes in Hong Kong waters is controlled mainly by water depth and salinity. Likewise, He Xixian (P.R.C.) noted that Holocene foraminifera from the same region differ greatly from upper to lower parts of the sequence. He attributed this to palaeogeographic changes caused by water depth, salinity and temperature variation during the Holocene transgression.

Palaeozoology

R.J.G. Savage (U.K.) began by reporting on his studies of South Asian mammal faunas to detect their sources. Middle Eocene sites in Pakistan and India have yielded a wide range of mammalian families whose affinities are essentially with central Asia or the African shores of the Tethys, while richly fossiliferous Miocene sequences from Africa, Eurasia and India provide a picture of migration between the continents during the later Tertiary. He discussed Afro-Indian links during the early Tertiary in detail. Nina Jablonski (Hong Kong) then reviewed the evolution and palaeobiogeography of monkeys in China (Fig. 2), noting that the fossil record of the Cercopithecidae extends back only to the Pliocene (Pakistan Siwaliks), while the majority in east Asia are Pleistocene. *Macaca* has been the most successful and widespread genus in China. One suggestion was that speciation followed fragmentation of the original range caused by the uplift of the Qinghai/Xizang Plateau during the Pleistocene.

Qiu Zhuding (P.R.C.) described recent discoveries of Neogene micro-mammals in China, including representatives from five orders: Scandentia, Insectivora, Chiroptera, Rodentia and Lagomorpha, with affinities to European, North American and Indo-Pakistan taxa. Differences in fauna in the north and south testify to major environmental differences. His colleague, Xu Qinqi then related the southward migration events of mammals during the Pleistocene to climatic changes. There were at least four such events: at 1.4 Ma, 0.9 Ma, 0.28-0.24 Ma, and most severe at 18,000-10,000 yBP.

The Plio-Pleistocene fauna and flora from the Siwalik sediments of Nepal were reviewed by Gudrun Corvinus (F.R.G.) who showed that the environment was generally waterlogged at the time of their deposition, with a humid tropical, non-monsoonal climate. Robert West (U.S.A.) showed how the stratigraphic and palaeontological record of the Kathmandu Valley, hitherto poorly known, is contributing to a general understanding of the geologic and palaeoclimatic history of the Himalaya. The combination of

Conference Reports

Palaeoanthropology

In the final session, Charles Oxnard (U.S.A.) used theoretical data models to show that hominoid sexual dimorphism can be distinguished and that the approximate adult sex ratio can be estimated. Study of distribution of real dental dimensions in samples of extant hominoid teeth (humans, gorillas, chimpanzees, bonobos and orangutans) confirms these findings. Wu Xinzhi (P.R.C.) agreed with earlier workers that there were exchanges of peoples between China and Australia, but he disputed the division of Australian human population into two types in the Late Pleistocene. The correlation of important Chinese hominid localities with O¹⁸ stages was then reviewed by Jean Aigner (U.S.A.) who was unconvinced that evidence from Zhoukoudian reveals long occupation. Floral and faunal analyses do not demonstrate climatic shifts of glacial- to interglacial magnitude. Here, as at Lantian sites, there are problems with the reliability of methods or consistency in evaluating chronometric dates.

The controversy over the date of Yuanmou Man (*Homo erectus yuanmouensis*) was discussed by Zhou Guoxing (P.R.C.) who provided evidence confirming the original

Early Pleistocene date. Finally, Sohn Pook (South Korea) showed that Palaeolithic hunters of the Yonggul Cave made wide use of bone tools as part of their cultural equipment. SEM photographs clearly distinguish between human and animal agency in the fashioning of bone remains.

A 76-page volume of abstracts of papers to the conference is available from the writer at US\$7.50, including postage, and the full proceedings are now in press. (Editor's Note: The detailed and informative *East Asian Tertiary/Quaternary Newsletter*, begun at the request of participants in the first conference and also published by the Centre for Asian Studies, is now at its seventh issue).

It is hoped that a third conference will be held in China, probably in 1991 in Kunming. □

Pauline Whyte
Centre of Asian Studies
University of Hong Kong
Pokfulam Road, Hong Kong

Correlating the Nonmarine Cretaceous

Urumqi, China, 24 - 30 August 1987

The far west of China was the site last year for the first symposium to be organized by IGCP Project 245 (Nonmarine Cretaceous Correlations). Despite the remote location some 35 foreign participants joined 70 Chinese specialists in the first wide-ranging review of nonmarine Cretaceous deposits around the world, so as to assess the potential for correlation between them.

A useful summary of the nonmarine Cretaceous of the Soviet Union, a topic that has received little attention in non-Russian literature, was provided by V. Krassilov and colleagues (U.S.S.R.). Similarly, D. Grigorescu (Romania) reviewed for his country the important but little studied Upper Cretaceous, which has yielded a rich vertebrate fauna of interesting paleobiogeographic affinities. Terrestrial Cretaceous rocks are found throughout Africa, but no detailed analysis of the correlation of these deposits has yet been achieved. Brief reviews by R.A. Reymont (Sweden) of the Saharan nonmarine Cretaceous and by J.D. Congleton and others (U.S.A.) of preliminary results from northern Cameroon provided an initial attempt in this direction. Other projects are active in Kenya and Malawi, and it is hoped that this activity will spread to other regions in Africa.

Linking nonmarine to marine strata is important in attempting a unified biochronological zonation. M. Matsukawa and I. Obata (Japan) presented a significant example of this approach from southwest Japan where the nonmarine and marine strata can be seen to interfinger. Since the Japanese nonmarine biochronology can, to some degree, be correlated with that of China, their work may have implications for dating the extensive nonmarine Cretaceous throughout China and Central Asia.

The comparatively well-studied nonmarine Lower Cretaceous of western Europe was subjected to biostratigraphic reassessment by E. Buffetaut (France), D.A. Norman (U.K.), and J. Neale (U.K.). Neale correlated ostracod evolution with salinity fluctuations and emphasized the promising

potential of ostracods in correlating nonmarine Cretaceous strata in the northern hemisphere. However, he pointed to considerable problems in using ostracods for correlating the northern and southern hemispheres, and this was corroborated in a paper on ostracod dispersal mechanisms by R.C. Whitley (U.K.). E. Hailwood (U.K.) provided a useful account of the magnetostratigraphy of the Wealden (Lower Cretaceous) of southern England and emphasized the importance of this approach for inter-regional correlation and the need for more such studies around the world.

The contributions by Chinese workers tended to be detailed assessments of specific aspects of the nonmarine Cretaceous of their country. A recurring theme, however, was

Figure 1: Some of the Cretaceous specialists at the Urumqi symposium. The two project leaders, N. Mateer and Chen Peiji are in the front row right side.

Conference Reports

the placement of the Jurassic-Cretaceous boundary, which has achieved little further consensus since 1985 (see Episodes, March 1986, v. 9, p. 41-42). Gu Zhiwei again urged a high stratigraphic placement for the boundary, while others preferred a stratigraphically lower position. Clearly, another approach is required to break the deadlock of these interpretations, for example using paleomagnetostratigraphy and correlation with marine sections. Liu Benpei analyzed the evidence for a temperature drop during the Early Cretaceous and argued that this might be used for event correlation throughout northern Asia. Chen Peiji assessed the age and displacement of the Tangcheng-Lujiand megafault of eastern China, the reconstruction of which is important for correlating beds on either side of this large structure.

Absolute dating methods have indicated in recent years that biostratigraphically dated strata need to be closely reviewed. N.J. Mateer, S.G. Lucas and A. Hunt (U.S.A.) gave an overview of Jurassic-Cretaceous boundary beds in western North America and reported that the boundary appears, in general, not to lie within a previously assumed extensive Lower Cretaceous hiatus, but rather within units previously thought to be Late Jurassic in age (notably, the Morrison Formation and equivalents). The implications of this reassessment have repercussions in other parts of the world, especially where typically Morrison vertebrate faunas have been found.

Two future directions for IGCP 245 were identified in Urumqi. These are the creation of a global biochronological zonation for the nonmarine Cretaceous, including the use of event stratigraphy (e.g., magnetostratigraphy) and radiometric dating, and the compilation of a distribution map of

the nonmarine Cretaceous deposits around the world with indications of their tectonic framework.

After the symposium participants visited the northern foothills of the Tian Shan at Ziniquanzi, south of Manas, west of Urumqi, where, a relatively complete Jurassic - Lower Cretaceous section could be seen, although there was disagreement about the position of the boundary. Lack of sufficient biostratigraphic detail in this remote region meant lithostratigraphic boundaries were used to identify time boundaries. However, the area shows promise for understanding the Cretaceous of Central Asia. A second excursion to the Turpan basin, southeast of Urumqi, examined a Triassic-Neogene continental sequence beautifully exposed in this desert region. Preliminary work in the Houyanshan (Flaming Mountain) uplift indicates another good section for evaluating the nonmarine Cretaceous of western China, including a possible continuously deposited Cretaceous-Tertiary boundary sequence.

One area largely ignored in this symposium was palynology, which is potentially of great significance. This topic will be the subject of the next project symposium to be held in conjunction with the 1988 International Palynological Congress in Brisbane. A volume of abstracts from the Urumqi symposium was published (available from Chen Peiji), and the proceedings will be published by China Ocean Press. □

Chen Peiji & Shen Yanbin
Institute of Geology
Academia Sinica
Nanjing
People's Republic of China

Niall J. Mateer
1467 N. 17th
Laramie, WY 82070
U.S.A.

European Research on Fluid Inclusions

Oporto, Portugal, 4 - 6 May 1987

Every second year, European scientists interested in fluid inclusion meet to discuss this rapidly growing discipline. The meeting has a traditional link with the International Association of the Genesis of Ore Deposits Commission on Ore-forming Fluids in Inclusions, and last year it was also sponsored by the IUGS Commission on Igneous and Metamorphic Petrogenesis. More than 200 geologists attended the meeting, which illustrated the rapidly developing techniques for studying fluid inclusion in practically all branches of geology. These include Micro-Raman Spectroscopy, infrared microscopy and isotope analysis, which in many cases allow a nearly complete, non-destructive analysis of single inclusions. However, it is also evident that fluid inclusion data alone may be ambiguous or inconclusive and that they must be systematically compared with evidence from independent sources, especially in metamorphic and igneous petrology. Much work is being done on diagenetic processes, and fluid inclusion studies are increasingly an essential part of any petrological investigation.

As in preceding meetings, some American inclusionists took part, but more unusual and promising for the future was the presence of leading petrologists from the U.S.S.R., Czechoslovakia, and Hungary, notably an important delegation from the Vernadsky Institute in Moscow. So much work on fluid inclusions has and is still being done in these countries that the west can only benefit from the development of personal contacts and collaboration. The meeting abstracts should appear in *Fluid Inclusions Research*, and a selection of the most important papers will be published as a special issue of the *Bulletin de Minéralogie*. □

J.L.R. Touret
Inst. voor Aardwetenschappen
Vrije Universiteit
Amsterdam, The Netherlands

The Jewel in the Crown of Gondwana

Kandy, Sri Lanka, 14 - 20 August 1987

A landmark conference on the geology of Sri Lanka and its relationship to other parts of Gondwana was held last year at the Institute of Fundamental Studies, in Kandy, organized by IGCP Project 236 (Precambrian Events in the Gondwana Fragments). Much new information on the Proterozoic mobile belts of the country was presented.

According to some participants the oldest rocks of the island (the Highland ones and Southwestern Group) are derived from a 2.0-3.2 Ga-old continental source terrain. An early phase of Proterozoic granulite metamorphism (7-8.5 kb, 760-930°C) at 1.1 Ga has been confirmed, and a second phase of granulite metamorphism (around 5.5 kb, 630°C) may be coeval with spectacular "in situ" charnockitization, dated at 746 Ma. Retrograde shear zones cut across all older structures and record massive fluid transport associated with widespread post-tectonic granite intrusions 550 Ma ago. Amphibolite grade orthogneisses from the Eastern Vijayan Complex were also metamorphosed at 1.1 Ga, but their Nd/Sm model ages of only 1.0-1.8 Ga suggest derivation from a geochemical source distinct from that of the Highland Series. Thus the melting event(s) producing the Eastern Vijayan gneisses are younger than the Highland Series rocks, and the Vijayan granitoids represent juvenile additions to the crust.

Figure 1: Highland Series marble at Digana, Sri Lanka, with metamorphosed and deformed dolerite dikes of pre-1.1 Ga age, similar to those in other parts of the once contiguous Gondwana fragments.

Detailed petrological work has demonstrated that, although of different origins, the Highland and Vijayan Series were coupled and underwent similar styles of metamorphism from 1.1 Ga onwards. Conference participants heard reports of the first occurrence of relict kyanite and staurolite as rare inclusions in garnets from sillimanite-bearing granulites in both the Highland Series and Southwestern Group. This shows that prograde metamorphism changed from earlier, relatively high-pressure and low-temperature conditions to later lower pressure and higher temperature ones. Striking similarities in geology between the Highland Series rocks of

Figure 2: The conversion (in situ charnockitization) of Highland Series hornblende gneiss into orthopyroxene granulite, possibly due to CO₂ streaming from the mantle.

Sri Lanka and those of Lutzow-Holm Bay of the East Antarctic Shield were also noted, as shown by relict kyanite, P-T-time paths of the major granulite facies metamorphism and the sequence of tectonothermal events.

The major amphibolite-granulite facies metamorphism at 1.1 Ga that affected Sri Lanka is also evident in the Proterozoic mobile belts of other Gondwana fragments such as East Antarctica and the Albany region of Western Australia. In some places this event represents a reworking of older crust; in others significant new crustal additions. Metamorphosed and deformed dolerite dikes in Highland Series, recognized for the first time during the conference, bear similarities to the reworked, meta-dolerite in Proterozoic mobile belts of Western Australia and the Rayner Complex of the East Antarctic Shield. The extent to which these mobile belts were the sites of ancient plate margins with new crustal growth as opposed to reworking of existing crust remains a fundamental problem.

It is apparent that adjoining crustal segments with differing isotopic and geochemical histories prior to 1.1 Ga ago evolved similarly during and after this event in Sri Lanka. The 700 Ma and 500 Ma ages (Pan-African) superimposed on these older mobile belts are recognized in East Africa, Antarctica, India-Sri Lanka and southern Australia. However, the manifestation of these "events" differs in character in different parts of eastern Gondwana. Thus, in the East Antarctic Shield granite and pegmatite were emplaced around 700 Ma ago, whereas Sri Lanka at this time was the site for spectacular in-situ charnockitization due to CO₂ streaming, which is absent from the East Antarctic Shield.

Conference Reports

In view of currently popular models for the origin of granulites due to CO₂ streaming, it is most significant that the charnockitization in Sri Lanka was not developed at the time of the main granulite facies metamorphism (1.1 Ga), but some 400 Ma later. The 500 Ma event has produced minor volumes of granitoid intrusions throughout much of the Gondwana Proterozoic mobile belts.

These data and new interpretations, together with the lively discussions at each outcrop on two enjoyable field trips, has provided a stimulus and framework for further work. The most pressing need is for detailed field and structural work to be undertaken in Sri Lanka. Indeed, a workshop on field techniques was run in conjunction with the conference for

geologists from China, India, Kenya, Sri Lanka and Tanzania, under the sponsorship of IGCP 236, AGID and the IUGS Commission on Tectonics. Greater effort should also be made to compare the geological history of India and Sri Lanka with that of East African Gondwana.

The excursion guide and abstracts of thirty-six papers presented at the conference can be obtained from the Geological Society of Sri Lanka, c/o Department of Geology, University of Peradeniya, Sri Lanka. □

David J. Ellis
Department of Geology
The Australian National University
GPO Box 4, Canberra City, ACT 2601
Australia

Andean Volcanism

Tucuman, Argentina, 14 - 18 September 1987

One of the important events held during last year's tenth Argentine Geological Congress was an international symposium on Andean volcanism. Sponsored by IGCP project 249 (Andean Magmatism and its Tectonic Setting), IAVCEI and several Argentine organizations, the meeting and related field trip were attended by some 180 geologists from 13 countries.

Participants in the pre-Symposium field trip studied and discussed outcrops of a wide range of Cenozoic volcanic rocks in the Puna-Altiplano plateau of the provinces of Jujuy and Salta between latitudes 22°30' and 24°S. This is a unique subduction-related region where magmatism has created broad ignimbrite plateaus and volcanic complexes. One highlight was the examination of a several hundred metre-thick section of ignimbrites that form the distal pyroclastic flows of the immense Panizos center (over 80 km in diameter) on the Argentina-Bolivia border. Others included a view of Quaternary monogenic cones of the shoshonitic series in the far eastern Puna, and a traverse of the Coranzuli stratovolcano center, one of the largest and easternmost Miocene Andean volcanic edifices, which appears to have been built on an older center of voluminous ignimbrites.

The majority of the papers were concerned with Tertiary to Cenozoic volcanism between latitudes 18° and 28°S (the Central Volcanic Zone - CVZ) and south of 33°S (the Southern Volcanic Zone - SVZ). Tertiary volcanism between 28° and 33°S, where modern volcanism is absent, was also reviewed, as was Precambrian to Mesozoic volcanism throughout the region.

One area about which new work is appearing is the back-arc region of the SVZ between 37° and 44°S. Paleocene to Oligocene volcanic rocks here (north of 42°S) have been shown to have transitional characteristics between the Tertiary-Recent volcanics of the Patagonian plateaus and the modern SVZ Andean arc volcanics. The arc and non-arc character of these rocks can be correlated in time with the relative obliqueness of convergence between the Farallon and South American plate.

Study of the back-arc region between 37° and 39°S has shown a complex interleaving of Quaternary calc-alkaline and alkaline volcanism resulting from intra-arc and back-arc extension. Similarly complex volcanic patterns may exist as far north as 34°S for the Quaternary and may have existed here and also to the south during the Tertiary. However, more geochemical and tectonic work is needed to understand the relations between subduction and intraplate volcanism over this whole region.

Figure 1: Field trip participants in front of the historical flow at Tuzgle Volcano (4300 m) in the eastern Puna near San Antonio de los Cobres, Salta Province, Argentina.

The genesis of the magmas that produced Quaternary basaltic andesites, the most basic rocks of the Puna, was attributed to varying degrees of partial melting of garnet peridotite and to compositional variations in the mantle below. Although crustal contamination and fractionation have occurred in these rocks, it was argued that their

Conference Reports

mantle signature has not been completely obscured. The spatial association with young faults is consistent with rapid rise from the mantle, little crustal contamination, and a change in tectonic style and volcanism in the eastern Puna in the last several million years. In contrast, crustal contamination is extensive in Mio-Pliocene andesites to dacites from the eastern Puna, and new $^{87}\text{Sr}/^{86}\text{Sr}$ data show that these rocks are among the most radiogenic in the central Andes.

Special attention was focused on the Tertiary evolution of the volcanics between 28° and 33°S , and correlations were reported between the chemistry of this magmatism through time and the shallowing of the subduction zone in this region. Other studies centered on Ordovician volcanism in the Puna, with the objective of identifying and characterizing early Paleozoic arc volcanism in this region. A future priority is to look at other Ordovician sequences here in more detail, particularly those in the region of the Faja Eruptiva of the eastern Puna, which appear to be of back-arc character.

Physical processes of volcanism were also discussed. The abundance of late Cenozoic calderas and related pyroclastic

flows permits detailed study of a great variety of ignimbrites and their structures, the key to mechanisms of formation and emplacement. Calderas of the order of 30-100 km in diameter with eruptive volumes of over 100 km^3 are common in the CVZ and SVZ, and many speakers highlighted their importance. An interesting example is the Cerro Panizos caldera, composed of eight main ignimbritic units. In spite of its large dimensions, no features indicating catastrophic collapse have been observed, although evidence for down-sagging is found. Pyroclastic flows also play an important role in early Paleozoic volcanism.

Conference abstracts are available through the Asociacion Geologica Argentina (Maipu 645, 1° piso, 1600 Buenos Aires, Argentina). Full papers will be published in 1988 in a special edition of the *Revista de la Asociacion Geologica Argentina*. □

Beatriz Coira
Instituto de Geologica Mineria
Universidad Nacional de Jujuy
Azorin 130 (4600)
S.S. de Jujuy, Argentina

Andean Neotectonics and Seismicity

San Miguel de Tucuman, Argentina, 14 - 19 September 1987

The Andean mountain belt constitutes a gigantic laboratory in which to investigate the interaction between earthquakes and surficial processes. It is also a region where a great deal of research is now being carried out by seismologists and experts on active tectonics. The tenth Argentine Geological Congress held last year provided an excellent opportunity to review this work.

One of the highlights of the Regional Symposium on Andean Neotectonics and Seismicity held then, with sponsorship from the IUGS Commission on Tectonics, was the presentation by B. Isacks (U.S.A.) on the origin of the Puna plateau and the related Andean orocline. He regards the subduction geometry of the different longitudinal segments of the Andes as controlled by previous magmatic and thermal histories within a framework of a non-collisional tectonics. The relationships between topography, orogenic shortening and the roll-over of the subducting plate beneath the oceanic trench provide the main geometrical constraints for this concept, which can be tested with the available paleomagnetic information.

New techniques developed in neotectonic studies by a French team led by M. Sebrier (Université de Paris) showed the importance of deviatoric stresses in producing local tensional faulting at the northern end of the Peru Altiplano, though the entire plate is submitted to the high compressive stress that characterizes Chilean-type subduction at these latitudes. Similar mechanisms were proposed for the southern border of the Puna plateau, by R. Allmendinger (U.S.A.) and co-workers.

Several local workers emphasized the importance to tectonic activity of lateral adjustments, not previously recognized. Since the last two big earthquakes of San Juan (1944) and Caucete (1977), the dominant view based on surface evidence has favoured reverse faulting as the main active process in the foothills of the Central and Southern Andes, and this is partially corroborated by earthquake focal mechanisms. New evidence from the Precordillera demonstrates the importance of right lateral strike-slip displacements, as demonstrated by H. Bastias (Argentina; see also article by Baldis et al., *Episodes*, September 1984).

The discussion of large and medium size intraplate earthquakes ranged from southern South America north to Costa Rica. Practical aspects were not forgotten, and special attention was paid to seismic risk and zoning, using as an example the host city of Tucuman, located in an active foothill area of the Andes. A few days before the meeting the city was awakened after many years of quiet by a M4 earthquake that shook "Tucumanos" and provided a precursor for the symposium itself. □

Victor A. Ramos
Servicio Geologico Nacional
Santa Fe 1548
1060 Buenos Aires, Argentina

Gold-bearing Hydrothermal Systems in Volcanic Centers

Santiago, Chile, 9 - 18 November 1987

Since the 1970s there has been a growing awareness that many mineral deposits of this region are located in eruptive centers marked by remnants of stratovolcanoes, calderas, dome complexes, diatremes, and fossil geothermal systems. The spatial and genetic relationships of epithermal gold-silver deposits to eroded volcanic centers in the central Andes are now recognized as important, for example, by the many mining companies now exploring for precious metal deposits in the Neogene-Quaternary volcanic complex of the central Andean region of southern Peru, western Bolivia, and northern Chile.

This young volcanic complex covers about a third of the central Andean mineral province, where most known mineral deposits are Miocene in age or older. New deposits are being discovered within this complex, and many others of Miocene and pre-Miocene age almost certainly exist beneath the younger cover. The El Indio Au-Ag-Cu deposit of northern Chile is by far the most economically important recent discovery in the young volcanic complex; it has produced about 65 metric tons of gold since 1979. Other deposits, some of which have been mined almost continuously since early colonial times, are now recognized as being genetically related to subvolcanic intrusions in former volcanic centers.

Against this background the Deposit Modelling Program (DMP), jointly run by IUGS and Unesco, held its fourth workshop late last year in Chile. Following the pattern of earlier workshops, the 45 participants including those from six Latin American nations attended lectures, took part in discussion sessions, and visited mines that illustrated features discussed in the workshop.

In her opening address to the workshop, Maria Teresa Canas Pinochet, Director of the Chilean geology and mines service (SERNAGEOMIN) recognized the advantages for exploration of using genetic models of hydrothermal systems in volcanic centers in the central Andes. She pointed out the importance of mineral deposits to the Chilean economy and noted that the exchange of ideas in the workshop could be of great benefit to future mineral exploration in Chile. Lectures by J. Davidson and C. Ramirez of SERNAGEOMIN provided information about the regional geological setting of mineral deposits in the central Andes. Davidson emphasized the role of plate tectonics in the magmatic and tectonic evolution of the Andes, and he reported that accretion was an important process along the western margin of ancestral South America during the Paleozoic but insignificant or absent during post-Paleozoic time.

The chemistry and structure of active geothermal systems were discussed by R. Henley (Australia) on the basis of his extensive studies in New Zealand and elsewhere. He explained how the information gained by studying active phenomena could be applied to the search for precious metal deposits in fossil geothermal systems. He discussed metal contents, temperatures, and alteration patterns of active geothermal systems and explained why boiling of hydro-

thermal fluids was an effective process in the deposition of metals in solution. Broadlands, New Zealand, was cited as an example of an active system in which gold is currently being deposited. Finally, Henley stressed the importance of understanding the movement and mixing of fluids in geothermal systems, factors of prime importance in the formation of orebodies.

Figure 1: Looking for anhydrite veinlets in drill-core at the Los Bronces deposit.

One focal point of the workshop was the relationship of geothermal systems and precious metal deposits. C. Cunningham (U.S.A.), who defined deposit models as "the systematic arrangement of the essential characteristics of a group of mineral deposits," discussed features of hot-spring, acid-sulphate and adularia-sericite gold deposits in the western U.S.A. and compared them with similar features in active geothermal systems to demonstrate the genetic relationships. A. Panteleyev (Canada) described epithermal precious metal deposits in volcanic centers of western Canada and presented a model for these deposits that relates features to magmatic-hydrothermal processes at depths to about 3 km.

Conference Reports

Data on mineral deposits and volcanic centers in other areas were also presented. H. Colley (U.K.) described the volcanic terrane of Fiji and reviewed criteria for recognizing deeply eroded volcanic centers of this area. M. Brodtkorb (Argentina) described the precious metal deposits of the Salle area of northwestern Argentina, which occur in an eroded stratovolcano complex. M. Arenas (Peru) described the classic epithermal precious metal deposits in the Neogene-Quaternary volcanic terrane of southern Peru, including deposits that have been exploited almost continuously since the 16th century. The major porphyry copper deposits of Chile, which together contain a large portion of the world's known resources of copper, were reviewed by E. Tidy (Chile), and A. Puig (Chile) related the El Guanaco, Cachinal de la Sierra, and El Soldado Au-Ag deposits of northern Chile to a newly discovered, eroded caldera complex.

Figure 2: F. Camus demonstrating the relationships between vein systems and nearby Cretaceous caldera at El Bronce de Petorca mine.

Interspersed with the formal workshop meetings were visits to several mining districts that illustrate specific characteristics of mineral deposits in volcanic centers. Los Bronces, a porphyry copper deposit, is characterized by at least seven hydrothermal breccia pipes and a diatreme in a terrane dominated by a quartz monzonite pluton. The

breccia fragments in the pipes have selvages of black tourmaline, and cavities between fragments are filled with pyrite, chalcopryite, and specularite. This deposit apparently formed in a hydrothermal system that vented to the surface, though no evidence remains of any former volcanic landform. In contrast, the El Bronce deposit consists of polymetallic base-metal veins bearing gold and genetically related to a deeply eroded, Late Cretaceous caldera.

The El Indio Au-Ag-Cu deposit, located in the high Andes near the Argentine border and discovered during an exploration program initiated in the mid-1970s, consists of two types of epithermal veins (gold-quartz and massive enargite-pyrite) in an intensely altered volcanic complex of Oligocene-Miocene age. Ore grades are variable, but one of the major gold-quartz veins averages about a quarter kilogram of gold per tonne, with local grades up to several kilograms of gold per tonne. The massive enargite-pyrite veins contain 4 to 10 g/t Au. The nearby, more recently discovered El Tambo deposit consists of gold-bearing breccia pipes containing barite and alunite. Typical ores contain 2 to 20 g/t Au and up to 50 g/t Ag.

In a review during the closing session of the workshop, Henley drew comparisons between models of active geothermal systems and features in epithermal gold deposits. He emphasized the need to recognize the upflow and outflow parts of geothermal systems, which influence the location of ore deposition, and showed how fluid inclusions could be used to achieve this end. Cunningham discussed the use of USGS Bulletin 1693 (Mineral Deposit Models, 1986) pointing out that the organization of models of the different types of deposits according to host-rock lithology and the indexing of these types by commodity and mineralogy make this a particularly useful reference. He also demonstrated ways to use fluid inclusions in paleothermal systems to determine position in a vein system relative to the paleowater table and to document processes that caused ore deposition such as boiling or effervescence.

It is anticipated that SERNAGEOMIN will publish the proceedings of the workshop. (Ed's Note: At the 1988 meeting of the DMP Steering Committee, D.C. Findlay (Canada) replaced G.H. Allcott (U.S.A.) as Chairman and plans were laid for future workshops in China (late 1988, on sedimentary copper deposits) and South America (on precious metal deposits) and for an international conference on deposit models to be held in Canada in 1990 in conjunction with the IAGOD congress.) □

- after report by C.G. Cunningham and G.E. Erickson

All-China Conference on Tectonics

Beijing, China, 24 - 28 August 1987

Last August several members of the IUGS family, including its Commission on Tectonics and ICL, joined with the Chinese Academy of Geological Sciences on behalf of the Geological Society of China to sponsor the third in a series of meetings that began in 1965, was then interrupted by the Cultural Revolution, and started again in 1979.

The meeting was opened by Ma Xingyuan, Chairman of the Organizing Committee and the Chinese Commission in Tectonics, who stressed that great progress had been made

in tectonics in China and abroad since 1979. He reviewed the importance of tectonics in the study of China's continental margins, large sedimentary basins, Precambrian geology, Phanerozoic orogenic belts and in deep seismic sounding to unravel the structure of the continental lithosphere. Several important new oil, gas and mineral deposits have been found in recent years and more discoveries are to be expected. Of particular importance for China is the increasing cooperation of earth scientists from various disciplines and the collaboration with colleagues from

Conference Reports

Figure 1: Generalized section across the Longmen Shan range. For explanation see text.

abroad. A. Kröner (F.R.G.), Secretary of the IUGS Commission on Tectonics, also pointed out that the modern tectonician can no longer be a specialist in his own field of research only but requires input from many branches of the earth sciences in solving some of the major unresolved questions in global tectonics and crustal evolution.

Among the scientific highlights were papers on the tectonic evolution of Tibet and the Tien-shan, and there were heated discussions on the favorite topic of Chinese geology, the evolution of the Qinling Belt. The Tarim Basin and other basins in NW China received much attention, particularly in view of the recent oil and gas discoveries there. Discussions on Precambrian geology centered on the evolution of the North China craton, and new data were presented on several circum-Pacific belts including southeast China, Japan and the Tethysides. The concept of exotic or suspect terrains seems to have found enthusiastic supporters in China, though there is a need for hard data. Recent crustal movements were discussed, and a wealth of new seismotectonic data was presented.

One of the post-conference field trips visited the Longmen Shan thrust and nappe belt of the Sichuan Province. This forms part of the 1000 km-long Chuan-Dian thrust belt along the eastern border of the Tibetan high plateau. It is important because it delimits the enormous, largely Triassic flysch-filled Songpan-Ganzi system against the Yangtze platform. The Longmen Shan belt suffered repeated shortening during the closure of the main branch of Palaeo-Tethys (Late Triassic-Jurassic), the closure of the Banggong-Nu Jiang ocean in Tibet (Early Cretaceous) and the Himalayan collision and subsequent intra-continental convergence (Eocene to present).

Until recently, there were only a few reports on this most interesting mountain range. However, Chinese earth scientists have accomplished a huge amount of work, especially during the last several decades, and the range has been mapped at a scale of 1:200,000. Figure 1 shows a generalized and schematic cross-section across the Longmen Shan range, which is divided by Chinese geologists into the frontal belt, the thrust belt, and the rear Longmen Shan belt.

The frontal belt rises abruptly from the Cheng Du plain with a number of flat irons consisting of Palaeocene to Triassic molasse sequences, of which the clasts yield much information about the uplift history of the Longmen Shan. The folding in this belt is clearly Cenozoic and earthquakes take place here at depths of 15 to 18 km (Fig. 2). It was reported that Longmen Shan was now rising by 0.1-0.3 mm/

yr. However, the reliability of the levelling and triangulation results may be questioned because of the shortness of the observed lines.

The thrust belt consists of two fundamentally separate parts. An eastern part is formed by Triassic to Cretaceous folded and thrust molasse, overlain locally by klippen shallow marine carbonates of Devonian through medial Triassic age. It seems clear that the klippen were thrust from the northwest in Cretaceous time, but there was some question whether this thrusting post-dated earlier folding of the

underlying molasse, or whether a refolded flat-ramp-flat structure could satisfy the present observations.

Figure 2: Examining exposures in a trench crossing the Haiyuan fault of NW China, which was active during the disastrous M8.5 earthquake of 1920.

The Gengda fracture zone separates the Pengguan crystalline complex from the rear Longmen Shan belt (Fig. 1). The latter consists of metamorphosed continental rise and slope sedimentary rocks of the Yangtze platform, Silurian to Late Permian in age, followed conformably by Triassic flysch. The source of this flysch was the then developing Songpan-Ganzi accretionary complex, a product of the closure of Palaeo-Tethys.

The major disadvantage of the meeting was the difficult communication between foreign and Chinese scientists as a result of the language barrier and insufficient translation facilities. This is a serious shortcoming, particularly in view of the rapidly increasing number of international research projects in China. □

after report by A. Kröner
and A.M.C. Şengör

Carboniferous Stratigraphy and Geology

Beijing, China, 31 August - 4 September, 1987

In 1927 the first International Congress of Carboniferous Stratigraphy and Geology (ICC) was held in The Netherlands. Sixty years later, 420 specialists from 31 countries gathered in Beijing for the eleventh congress, the first time this has been held outside Europe and North America. Previous ICCs contributed significantly to the search for fossil fuels and mineral resources from Carboniferous rocks worldwide, and the XIth ICC included for the first time as major topics coal geology and Permian stratigraphy. Indeed the Permanent Committee for the Congress decided in Beijing that future meetings would be labelled "International Congress on Carboniferous-Permian Stratigraphy and Geology," with the next session to be held for the first time in the southern hemisphere in Buenos Aires.

A highlight of the Beijing ICC was the presentation of reviews by Chinese scientists on the Carboniferous stratigraphy, geology, and coal resources of China. Detailed descriptions of Carboniferous stratigraphy in China were presented to participants in the form of three books from the Nanjing Institute of Geology and Palaeontology: Carboniferous Stratigraphy of China (ed. Zhang Linxin), Carboniferous Boundaries of China (ed. Wang Chengyuan), and Bibliography of Carboniferous and Permian Stratigraphy and Palaeontology (ed. Jin Yugan et al.). Special issues on Carboniferous-Permian paleontology were also presented in the journals Acta Palaeontologica Sinica and Acta Micropalaeontologica Sinica.

Coal is the major energy resource of China; total resources amount to 40,000 billion tons, with an annual production of about 0.9 billion tons. Carboniferous and Permian deposits now provide the basis for the coal industry of China. In addition to deltaic deposits, Permian coal is characterized by the presence of back-barrier, peat flat deposits.

The question of a global stratotype for the Devonian-Carboniferous boundary attracted much attention during the Congress. Boundary sequences in shallow-water platform, slope, and basin facies in South China were described in detail, and two candidate sections were proposed: the Nambiancun Section in Guilin, Guangxi Province, China and the La Serra Section of France. It was decided to vote on the candidate sections at a meeting to be held in 1988 in Ireland.

For the mid-Carboniferous boundary, the theoretical basis for which had been decided at the last ICC (see Lane and Manger, *Episodes*, vol. 8, No. 1, June 1986 - Ed.), three candidate sections were discussed in Beijing. These were the Jingyuan Section of Gansu and the Ludian Section of Guizhou, China, and the Arrow Canyon Section, U.S.A. As regards the stratigraphical level of the Carboniferous-Permian boundary, there was a wide difference of opinion. In Beijing the emphasis was on sequences close to the boundary in platform and slope facies in South China and the western Urals. The boundary Working Group of the IUGS Subcommission on Permian Stratigraphy was reorganized, with Wu Wangshi (China) and B.I. Chuvashov (U.S.S.R.) as Chairman and Vice-Chairman respectively.

Although opinions differ as to whether the Permian System should be subdivided into two or three parts, most of the participants at the meeting were in favour of taking the base of the Kungurian as the base for the Upper Permian Series, which would then include 5 stages: Kungurian, Ufimian, Kazanian, Dzhulfian and Changhsingian (Dorshamian). Correlation charts prepared by working groups in Japan, U.S.A., U.S.S.R., China, Australia, India and southern Europe will soon be made public.

Figure 1: On the Li River, Guilin, en route to examine a proposed stratotype section.

It was clear at the meeting that Paleozoic miospores are being used more and more to establish worldwide zonal correlations, especially in China, Ireland, Poland and the Donetz Basin of the U.S.S.R., though there remains a shortage of data on Namurian miospores. The International Commission on Paleozoic Microflora, therefore, decided in Beijing to set up a working group to study this subject.

As regards coal, an international classification has been suggested by the International Commission on Coal Petrology after many years discussions, based primarily on the ranks of coalification, types and facies. Three subcommissions are respectively engaged in coal nomenclature and standardization, and the geological and industrial applications of coal petrology. The scheme for bitumen nomenclature has been adopted already, and the new discipline of organic petrology has come into being.

Conference Reports

Another topic of special interest at the ICC was the reconstruction of global and regional paleogeography for the Carboniferous and Permian. New geophysical data has helped to determine the relative position between different tectonic blocks in China. And it is now clear that the minor transgression during the Visean and Namurian was an important period for the widespread formation of syngenetic ore deposits in western Europe and China as well as of important oil and gas in oil fields.

The congress was held under the auspices of the Palaeontological Society of China with the help of the Nanjing Institute of Geology and Palaeontology, Academia Sinica, and several other Chinese organizations. □

Jin Yugan

Nanjing Institute of Geology and Palaeontology
Academia Sinica, 39 Eastern Beijing Road
Nanjing, China

Prospection РАЗВЕДКА
Exploration 87
Exploracion 勘探

Finding Minerals and Groundwater

Toronto, Canada, 27 September - 1 October 1987

Nearly 1000 geoscientists from industry, government and university in 76 countries gathered in Toronto last September in the third in a decennial series of conferences on mineral and groundwater exploration. For a major conference in North America there was a higher than usual participation from developing countries, especially Africa, thanks to major financial support from the Canadian International Development Agency. Despite the last few years of doom and gloom in the minerals industry, the mood at Exploration 87 was markedly buoyant, and the current rage for gold was very much in evidence.

As regards the general outlook for the future of minerals, Peter Fozzard (U.S.A.) spoke of the end to the age of materials and of the trend away from the search for large deposits (especially base-metals) towards smaller scale ones, especially "exotic" minerals. W.S. Fyfe (Canada) echoed this theme by pointing to the increased use of ceramics and other metal substitutes, such as fiber optics for copper wire. D.W. Strangway (Canada) even went so far as to discuss minerals on the Moon and the asteroids, and ways to locate them involving geophysical, geochemical and remote sensing techniques modified from terrestrial programs.

Most attention was devoted to exploration techniques. H.E. Siegel (Canada, the Conference Chairman) emphasized airborne surveys using helicopter-borne multi-sensors such as lasers to measure mineral reflectance. He also pointed to the potential of high-resolution, multi-channel seismics for mineral exploration, even in hard rocks. Remote sensing techniques were, of course, in evidence, especially in the poster displays, and the growing interest in multispectral digital data was obvious, even for vegetation-covered areas. A.R. Barringer (U.S.A.) spoke of the new generation of airborne methods, including advanced, digital time-domain EM systems, multi-component magnetometers and geochemical methods, and of the importance of using Global Position Satellite (GPS) receivers to position ground stations more rapidly and accurately in three dimensions. This would especially facilitate the routine use of the gravimeter in

exploration. Of course, the use of computers, even laptops on field surveys, drew the attention of almost everyone at Exploration 87.

Case histories, especially in poster form, attracted much attention. To quote but one example, M. Lewis (U.S.A.) reviewed the Perkoa volcanogenic massive sulphide deposit of Burkina Faso discovered in 1982 by the UN/DTCD (see article this issue by Guy-Bray and Lewis) in the Birrimian greenstone belt of West Africa. The primary discovery tool was soil geochemistry, which turned up two weak Zn anomalies in a reconnaissance over this flat, unexposed terrain, and further geochemical sampling plus traditional ground geophysics (magnetics, EM, IP, resistivity) and drilling outlined a massive sulphide lens 300 m long x 1.15 m wide x 500 m deep.

Figure 1: Drilling for gold in the Peruvian Andes. Photo courtesy of UN Revolving Fund for Natural Resources Exploration.

Conference Reports

There were many participants and many examples illustrated from Australia, and much praise was given to exploration geologists there. Yet R. Woodall (Australia) in his keynote address to the conference stated that "our image out in the community is not nearly so glamorous. We are, to an extent, like Narcissus, admiring our reflection in a pool. . . . We can maximise the chances of world-class discoveries, and thus the return on the investment in exploration through innovative management." This requires that decision-making be "in the hands of competent scientists who are aware of the world of economics and competition. Exploration requires more than technical excellence; it requires persistence, flexibility, decentralised management, and an informed and sympathetic government."

Woodall stressed the need for the minerals industry to talk the language of the public and to attract the attention of the media so that they can properly publicize the important role of mining in society and its many successes. He suggested the use of the term "innovation" rather than "exploration," which seems to bring with it a "tarnished, diminished, and distorted image."

Exploration '87, sponsored by the Canadian Geoscience Council, itself had an interesting innovation, a series of unstructured evening sessions to discuss informally "hot" topics. These included such themes as the twilight of the mining industry (doubted by most), the need for more, advanced training (but in universities, or on-the-job?), remote sensing ("very good for filling in lineaments after an orebody has been found," according to one sceptic; "excellent material for lectures" - from another), leaking vapours (what are the Swedes doing?), and how to gain more information from drill holes ("can we learn from the petroleum industry?").

In a session devoted to national and international programs C. Oiti Berbert (Brazil) reviewed the long history of mineral exploration in his country, starting with the "Adventurer's Phase" (1500-1811) when gold and gemstones were found. The period 1811-1907 he termed "the Foreigner's Phase"

when gold had declined, the first scientific geological reconnaissance of the country was undertaken, and the first institutions established (e.g. the School of Mines in Ouro Preto, and the Sao Paulo State Geological Commission). The first "National Phase" followed the establishment of the Brazilian Geological Survey in 1907, and included the formation of state enterprises (DNPM in 1934, followed by CVRD and Petrobras), and the discovery of the huge mineral deposits of Carajas.

The latter part of this phase (1970-1985) saw the completion of 1:1,000,000 geological mapping of Brazil (and 42% of the country at 1:250,000), and the discovery of some 100 new mineral deposits, now mostly in production. Since 1978, however, there had been few discoveries (except Serra Pelada!), and there was a slowdown until 1985 when a new national phase began. This would last until 1999 under a plan for 1:50,000 and 1:100,000 mapping of 3,100,000 km² of Brazil and many detailed surveys, with the focus on gold, gems, coal and peat, and construction materials, especially in the Amazon region.

Near the end of the conference, P.G. Cooray (Sri Lanka), asked what was really meant by the term "development?" Stating that "exploitation of minerals for export is no longer acceptable as a major role of the geosciences, or as a measure of development," he urged a re-focussing of efforts on industrial minerals for local use, more efforts on the research for and management of groundwater, studies on agrogeology and the search for better (less polluting) fertilizers, urban and environmental geology. If those participants who listened to Cooray did not agree with his views, they were at least challenged about their role in society.

The proceedings of Exploration 87, when published, will undoubtedly provide a very useful state-of-the-art summary of exploration techniques and issues, as did the final reports of the two earlier Exploration conferences. Further information may be obtained from H.O. Seigel (Scintrex Ltd., 222 Snidercroft Rd., Concord, ON, Canada L4K 1B5). □

- Episodes Staff

Rocks, Fossils and History

Pisa and Padua, Italy, 23 September - 1 October 1987

The 13th symposium of INHIGEO, the IUGS Commission on the History of Geology, was held recently in Italy in order to stimulate interest in the history of the earth sciences in that country, and to bring Italian contributions to the development of geology to the attention of the worldwide scientific community. The spectacular geology, landforms and sites of historical interest were magnets to the historians of geology from 14 countries who gathered in northern peninsular Italy in late September.

The early sessions belonged to those of the non-Italian contributors who dealt with the biostratigraphic matters from a mainly European standpoint. One group of papers was mainly concerned with the contribution of particular geologists to European scholarship. These included Goropius

Becanus (F. Ellenberger; France), Zygmunt Bosniacki (W. Narebski; Poland), Nicholas Steno (G.H. Davies; Republic of Ireland), Charles Lapworth (B.M. Hamilton; U.K.) and Nicolas Demarest (K.L. Taylor; U.S.A.). The second category of papers dealt with broader issues and ideas on a more national basis, such as German stratigraphy (M. Guntau; G.D.R.), palaeontology in Russia (V.V. Tikhomirov; U.S.S.R.), Scandinavian geology in Europe (G. Regnell; Sweden), and Italian/German influences (W. Langer, F.R.G.). There were also papers on the value of geological work on the diagnostic qualities of jade and nephrite in establishing the evolution of the jade cultures of Neolithic China (Wen Guang; China) and of perceptual bias in the interpretation of geological photographs (S. Figueiroa de Mendoça, Brazil).

Conference Reports

The papers by Italian participants discussed Tuscan geology, (A. Riparbelli), volcanicity (L. Ciancio, P.D. Burgazzi, E. Schettino and A. Nazzaro), Leonardo da Vinci (G. Vai), Giuseppe Monti, the founder of palaeontology in eighteenth century Bologna (C. Sarti) and Giuseppe Scarabelli, the creator of the first geological map of the Romagna (S. Marabini). Finally there was a socially based analysis of nineteenth century Sicilian science by L. Montanari in which the role of the aristocracy in the development of fossil collecting and volcanic studies was examined.

The two scientific sessions were separated by a field excursion, which took the party across peninsular Italy from the

N.W. Apennines to the Veneto region, with stops at places significant in both the geological development of Italy and the growth of its history. The rocks, mainly Tertiary, showed spectacular features, both in structure and Quaternary and Holocene dissection. □

B.M. Hamilton
Head of Geography
Liverpool Institute of Higher Education
Woolton Road
Liverpool L16 9ND, U.K.

Evolution of Metamorphic Belts

Dublin, Ireland, 17 - 30 September 1987

Many participants in IGCP Project 235 (Metamorphism and Geodynamics) met last year in Ireland to discuss the pressure-temperature-time (P-T-t) histories of metamorphic belts of different ages and tectonic settings. E.J. Essene (U.S.A.) opened the proceedings with a critical review of the current methods of determining metamorphic pressure and temperature. Relative thermobarometry - the determination of changes in P and T as recorded by the chemical zonation within individual grains in a rock sample - was described by F. Spear (U.S.A.), who advocated its application as a technique for monitoring structural evolution. While many speakers implicitly based their thermobarometry on the assumption that metamorphic rocks obey the rules of equilibrium thermodynamics, others questioned this assumption. As an example, G.T.R. Droop (U.K.) illustrated a case of disequilibrium reaction in aluminous granulites from the Archaean Limpopo belt.

The principles and practice of dating metamorphic processes were reviewed by P.K. Zeitler (Australia) in an elegant case study from the Himalayas. A detailed chronology of events in a very young orogenic belt can be obtained using ion-microprobe U-Pb data from zircon. Isotopic systems in most minerals are generally thought to record the time elapsed since the so-called blocking temperature was reached and are usually regarded as cooling ages. Dating of metamorphic mineral growth as opposed to metamorphic cooling is rarely achieved, but K. Mezger (U.S.A.) showed that the U-Pb method could be applied successfully to date garnet growth at all but extremely high metamorphic temperatures. R.D. Dallmeyer (U.S.A.) emphasized the importance of grain-size and grain boundary effects in the interpretation of $^{40}\text{Ar}/^{39}\text{Ar}$ age studies of very low grade slates and metatuffs.

Several contributions demonstrated the success of a multi-disciplinary approach based on integrating thermobarometric estimates with geochronological data on metamorphic minerals, supplemented by fission track data in younger belts. P. Dymoke (U.K.) presented a detailed study of the Dalradian of the SW Highlands of Scotland, and V. Schenk (F.R.G.) traced the uplift history of a lower crustal section from southern Calabria during the Hercynian and Alpine orogenies.

The relative timing of metamorphic and deformational events was raised by R.H. Vernon (Australia), who discussed

recent developments in the interpretation of microstructural relationships, while J. Ridley (Zimbabwe) addressed the matter of the outcrop and larger scale. On a still larger scale - both temporal and spatial - C. Beaumont (Canada) considered the relationship between the deforming orogen and events in the foreland, and he showed how metamorphic P-T-t data could be related to the sedimentation history of foreland basins.

Until recently thermal modelling of metamorphic belts has concentrated on conductive heat transfer, and several speakers sought to explain thermal anomalies in various metamorphic terrains by convection. C.P. Chamberlain (U.S.A.) and T. Hoisch (U.S.A.) invoked aqueous fluid as the medium responsible, while J. De Joreo (U.S.A.) demonstrated a case of magmatic heat transfer. Convective heat transfer was generally accepted as important, though J.B. Brady (U.S.A.) pointed out that the magnitude of the effect is still unclear.

Figure 1: R.H. Vernon demonstrates metamorphic fabrics in the Ox Mountains of County Mayo to S. Banno and other field trip participants.

Conference Reports

As at previous IGCP 235 meetings, granulites were much discussed. S.L. Harley (U.K.) considered crustal extension as a control on isobaric cooling of certain granulites and emphasized the need for a quite separate event to cause their eventual exhumation. Anticlockwise P-T paths caused by the intrusion of hot magma, especially under continental arcs, was suggested by S.R. Bohlen (U.S.A.) as an alternative to the widely accepted continent-continent collision model for the origin of granulites. J. McLelland (U.S.A.) argued that recrystallization of already anhydrous assemblages played an important role in the development of the classic Adirondack Highlands granulites.

During a workshop on geothermobarometry, Bohlen (U.S.A.) and K.K. Podlesskii (U.S.S.R.) gave the experimentalists' perspective on well-calibrated barometers, and S. Banno (Japan) discussed the treatment of activity composition calculations. The application of the Gibbs method as

applied to P-T-t path calculations was explained by F. Spear (U.S.A.), while S.L. Harley (U.K.) and B.J. Hensen (Australia) discussed applications to actual rocks. Practical sessions included demonstrations and distribution of micro-computer software for thermobarometry.

A majority of the papers presented at the meeting will be published by the Geological Society of London, a co-sponsor of the meeting, through its Metamorphic Studies Group, which is also affiliated to the Mineralogical Society of Great Britain and Ireland. The next meeting of IGCP 235 is to be held in Leningrad in mid-1988. □

J.S. Daly
Department of Geology
University College Dublin
Belfield, Dublin 4
Ireland

Event Stratigraphy and the Pacific Neogene

Berkeley, California, 29 - 31 July 1987

The IUGS Commission on Stratigraphy's Regional Committee on Pacific Neogene Stratigraphy held its 4th International Congress last year in conjunction with IGCP Project 246 on "Pacific Neogene Events in Time and Space." Some 180 scientists from 16 different countries participated in discussions ranging from paleoclimatology, stratigraphy and geological history, to evolutionary studies, modern biofacies and sediment relationships.

Reviewing climatic events detected from late Neogene strata of New Zealand, A.R. Edwards (New Zealand) stated that the extinction of about 25% of New Zealand molluscan genera during the latest Miocene (Kapitean Stage) accompanied the greatly accelerated diversification of planktonic foraminifer lineages; New Zealand events are coeval with the Messinian "salinity crises" in the Mediterranean. A series of events including extinctions of mollusca, appearance of nannofossils and of glacial rock-types, and for-

minifera speciation, in Late Pliocene rocks reflect climatic deterioration. A sub-Antarctic molluscan fauna abruptly appeared in central North Island at 2.4 Ma, coeval with the onset of major glaciation in the northern hemisphere.

The variation in diversity of larger Neogene foraminifera throughout the Indo-West Pacific was discussed by C.G. Adams (U.K.). Contrary to earlier statements by Shackleton that central Pacific climates had much lower gradients than present during the mid-Cenozoic climatic optimum, Adams argued that faunal diversity gradients were then actually similar to those of today, with past latitudinal ranges having expanded and contracted to reflect climatic deterioration and amelioration. Huang Chi-Yue (Taiwan) interpreted a marked faunal change in Neogene benthic foraminifera accompanying the Late Miocene hiatus in Taiwan (12.9-10.2 Ma) as resulting from the marked intensification of circulation of the Kuroshio Current. His colleague, Liew Ping-mei showed that five thick, shallow marine Pleistocene sequences in Taiwan, all have been dated by magnetostratigraphy, record a prominent cool phase, dominated by Pinaceae pollen, that is correlated with oxygen isotope stages 16 and 18.

A special session on the Monterey Formation and related rocks highlighted their importance to Pacific Neogene stratigraphy, paleoceanography and hydrocarbon exploration, because of the wide distribution of this distinctive diatomaceous facies around much of the Pacific rim. Various speakers emphasized the suboxic paleoenvironmental setting of these organic-rich sediments, the paleoceanographic implications of terrigenous, calcareous, and siliceous facies within the Monterey sequences and diagenetic alteration of Monterey sediments to porcelanites and cherts. Considerable discussion centered on analogues in Peru and Japan and their use as indicators of increased upwelling and productivity in the Middle and Late Miocene.

Two Japanese speakers, K. Hirooka and A. Hayashida, demonstrated with ample paleomagnetic data the rapid rotation of the Japanese Islands and the resulting opening of

Conference Reports

the Japan Sea about 15 Ma ago. The paleoclimatic history of West Antarctica, based on the results of ODP Leg 113, was reviewed by J.P. Kennett (U.S.A.), who noted that a major cooling of bottom waters and the permanent establishment of the East Antarctic ice sheet occurred between 16.5 and 13.5 Ma ago.

New techniques for refined correlation and high resolution chronology were represented by T.S. Loutit (U.S.A.), using seismic stratigraphic analyses of deep sea and outcrop sections, by L. Mayer (Canada) on water-gun seismic profiles, by L.D. Keigwin (U.S.A.) on late Neogene isotopic events, and by D.J. DePaolo (U.S.A.) on strontium isotopes applied to the Monterey Formation.

The need for proper analysis of Tethyan and Pacific events in view of paleoceanography, paleobiology, and geodynamics was stressed by F.F. Steininger (Austria), and many summaries of Neogene stratigraphies were presented by speakers from U.S.A., U.S.S.R., China and New Zealand. For IGCP 246, Project Leader R. Tsuchi (Japan) proposed four major Neogene events for joint international study. These are the mid-Neogene climatic optimum at 16 Ma,

Middle Miocene upwelling and productivity events between 14 and 9 Ma, the opening of the Sea of Japan at 15 Ma and related tectonic events, and accelerated evolutionary events occurring in the latest Neogene between 3 and 1 Ma.

IGCP-246 has now entered its second phase of activity, and participants have begun the synthesis of some major Pacific Neogene events in time and space as well as the analysis of their nature and interrelationship. Forthcoming events sponsored by IGCP-246 include an international symposium at Nanjing, China, on June 2-3, 1988, under the title of "Pacific Neogene: Continental and Marine Events"; an international seminar for committee members at Shizuoka, Japan on October 3-5, 1988; and an international symposium at Vina del Mar near Santiago, Chile on August 7-10, 1989, on "Pacific Neogene: Stratigraphic, Paleocceanographic and Andean Events."

A 131-page volume containing 100 extended abstracts from the Berkeley meeting can be purchased for US\$10 from Dr. J.R. Blueford, U.S. Geological Survey, MS 144, 345 Middlefield Road, Menlo Park, California 94025, U.S.A. □

after reports by R. Tsuchi and others

Neotectonics in Taiwan

Taipeh, Taiwan, 31 March - 11 April 1987

Papers given at international conferences may take years to appear in print. Some of them never make it. A splendid exception is provided by the proceedings of the first Sino-British geological conference to be held in Taipei on geotechnical engineering and hazard assessment in neotectonic terrains, and published a mere six months later. The meeting was attended by 85 earth scientists from Taiwan, 20 from the U.K. one from F.R.G. and one from Norway.

Figure 1: Conference delegates inspecting a Quaternary section.

The relief of Taiwan has developed since the Pliocene, following collision of an island arc on the Pacific plate with the margin of Asia. Slope instability and frequent large earthquakes have been countered with great skill, but a rapidly growing population and the demands of industry mean that pressure is increasing for development in areas hitherto untouched by the engineer. Conference sessions dealt with the geological and geomorphological setting, the problems that face engineers in areas affected by neotectonic deformation and the attendant instability, and case studies from Taiwan and elsewhere.

The field trip, led by Christopher Fong and Hao-Tsu Chu, gave the delegates a very clear view of the diversity of terrains to be encountered in Taiwan, including many features of manifestly tectonic origin, notably the active faults of the Longitudinal Valley and the many areas subject to rapid uplift during and since the Pliocene. That the geomorphological problems and the associated issue of seismicity can be successfully circumvented was illustrated by two spectacular dams, the Tchiu and the Feisui, and several impressive examples of highway engineering.

The ideas and information exchanged at the conference reflect the lively interest among all the participants in the problems created by neotectonic processes, their explanation and their geotechnical solution. Indeed, plans are well under way for a second conference to be held in the U.K. in June 1989. This meeting will be open to others interested in the interaction between tectonics and engineering. □

C. Vita-Finzi
Department of Geography
University College
26 Bedford Way
London WC1H 0AP
U.K.