Deposit Models for Bolivia

La Paz, 4 - 16 September 1988

A mineral deposit model is the systematic arrangement of the essential characteristics of a group of mineral deposits. There are strong parallels in methodology that exist between the search for mineral deposits and the practice of diagnostic medicine. In both fields, the models consist of large assemblages of "symptoms," each of which may be misleading if not evaluated in the light of other symptoms in the assemblage. Again in both medicine and geology there is a logical sequence to follow because the outcome of the search for one symptom determines other symptoms that should be sought or lines of inquiry to be pursued.

The advantage of this approach in geology is that quite reliable tentative determinations for most ore deposits can be made using existing data and simple field and laboratory observations, provided there is a framework of models against which the results can be evaluated. For the developing world, a knowledge of models and how to use them is the most valuable asset a country can have in conducting a search for mineral resources. It allows widespread exploration by low-cost methods and permits technologically complex and high-cost techniques to be reserved for deposits that show real promise. Too many nations make the mistake of leaping indiscriminately into advanced exploration, thereby squandering scarce resources. Furthermore, a fully developed portfolio of deposit models allows a nation to appraise its mineral potential and to construct mineral asset management strategies even before all its holdings in resources have been discovered.

Existing deposits are the products of expensive research (\$30 million in U.S. alone) by both the public and private sectors during the last decade or two. Some are published, some are described in computer data bases, and others remain unpublished concepts. Just as in medical diagnosis, "clinical" training is required to take full advantage of mineral-deposit modelling expertise.

The IUGS/Unesco Deposit modelling program is designed to facilitate the formulation, compilation, and transfer of mineral-deposit modelling expertise to developing coun-

tries. It promotes more efficient and effective use of models in both resource assessment activities and exploration by sponsoring mineral-deposit modelling workshops, which bring international experts on a particular type of deposit or tectonostratigraphic terrane to work with geologists from the mining industry, universities, and geological surveys of a country or region. Each workshop focuses on a limited number of deposit types of known or potential economic importance to the host nation. The purpose is to provide a training format in which the characteristics of world-class deposits can be related to the deposits or prospects in the country.

Figure 2: Jose Pinto of the Servicio Geologico de Bolivia showing the relationships between altered rock and ore at the Pulacayo deposit.

Figure 1: Participants in the Bolivia workshop with the famous Cerro Rico deposit of Potosi in the background.

Four deposit modelling workshops have now been held, in Khartoum (1985), Manila (1985), Belo Horizonte, Brazil (1986), and Santiago, Chile (1987), as earlier reported in *Episodes*. The most recent was a workshop in Bolivia on volcanic-hosted, epithermal precious-metal deposits. This included an eight-day, 2,200-km field trip, which examined deposits first mined for precious metals during Colonial times, some of which, like Cerro Rico, have been world class silver producers. The formal sessions in La Paz were attended by 145 registrants from around the world, though more than 200 people attended most sessions. As the La Paz newspaper *Presencia* reported "The Bolivian professionals explained that a visit to a deposit in the company of the international leaders was like attending a three month course at the University."

The workshop theme was selected because of the newly recognized importance of the spatial and genetic relationships of epithermal gold-silver deposits to eroded volcanic centers in the central Andes. There is a growing awareness that many mineral deposits of this region are located in eruptive centers marked by remnants of stratovolcanoes,

Conference Reports

calderas, dome complexes, diatremes, and fossil geothermal systems. Many mining companies are currently exploring for precious metal deposits in the Neogene-Quaternary volcanic complex of the central Andean region of southern Peru, western Bolivia, and northern Chile. This young volcanic complex covers about a third of the central Andean mineral province, where most known mineral deposits are Miocene in age or older. New mineral deposits are being discovered within this complex, and many others of Miocene and pre-Miocene age almost certainly exist beneath the younger cover.

The workshop strongly confirmed that there is excellent potential for discovery of new precious metal deposits in Bolivia. An appropriate geological setting for these deposits is provided by the many young, magmatically evolved volcanic rocks, which are little eroded and which have associated widespread, locally intense, argillic to advanced-argillic alteration zones. Indeed, some of these deposits are already known, and there are many other attractive targets for exploration.

Most of the known precious-metal deposits that were examined during the workshop are related to volcanic domes of a variety of shapes, sizes, and composition. Some domes

have well-developed tuff rings and explosion breccias, others have none. Some were formed by viscous flows, some developed an immiscible fluid phase, some were cut by breccia pipes, and others were intruded by younger rhyolites. An important challenge is to determine which features (if any) are unique to domes that developed ore-forming systems, and which hydrologic/structural features localized ore.

The workshop leaders included C.G. Cunningham (U.S.A.), M. Bernstein (Chile), R. Henley (Australia), A. Panteleyev (Canada), R. Sillitoe (England), and A. Wallace (U.S.A.). Publications from the workshop included extended abstracts, a field guide, workshop notebook, and a Selected Bibliography of Volcanic-hosted, Epithermal, Precious-Metal Mineralization.

after reports by C.G. Cunningham, A.T. Ovenshine
and G.H. Allcott
U.S. Geological Survey
Reston, VA 22092
U.S.A.

Precambrian Dyke Swarms: Status and Progress of IGCP Project 257

Lund, Sweden, 8 - 10 August 1988

Mafic dyke swarms, some of enormous size, are found cutting Precambrian Shields on every continent, having been injected at intervals throughout Proterozoic time. An understanding of these major swarms in terms of mantle and tectonic processes requires coordinated and cooperative studies on a global scale. IGCP Project 257 (Precambrian Dyke Swarms) has five main scientific objectives: to study how the dyke were emplaced, to correlate swarms with effusive products, to define dyke geometry in three dimensions, to correlate Precambrian swarms on a world-wide basis and to test if times of major dyke injection were globally synchronous during the Proterozoic, and to further understanding of the origin of Precambrian swarms in terms of crustal tectonics and mantle processes.

In order to make Project 257 a realistic and meaningful enterprise 16 working groups have been formed, based on national or thematic lines. Several of these working groups, such as the Scandinavian, Canadian, Indian and one concerned with the North Atlantic Craton have already made significant advances in terms of scientific goals. New research projects on dykes have been started in Australia, Brazil, Canada, China, Finland, India, Norway, Sweden, U.S.S.R., U.S.A. and Zimbabwe.

Project 257 held its first major meeting recently in Sweden when some 50 scientists from 13 countries attended a symposium on "Mafic Dykes and Related Magmatism in Rifting and Intraplate Environments." The symposium provided an excellent blend of regional reviews, new field and laboratory data, and discussion of major problems both old and new. The Lund conference was particularly important in bringing together the Scandinavian and Soviet working groups, an essential step if comparisons are to be made between dykes in the western and eastern Baltic Shield. The following

Figure 1: Cenozoic dykes cutting Mesozoic sandstones, Monastery National Park, Utah.

account attempts to present a state-of-the-art overview, derived both from papers presented at Lund and from recent publications (especially Geological Association of Canada Special Paper 34, 1987, "Mafic Dyke Swarms," edited by the writer and W.F. Fahrig, 503p.).

One major issue was whether dykes and swarms are fed primarily by vertical or lateral flow of magma. Studies that are addressing this problem are being undertaken on the two largest radiating swarms in Canada, the Matachewan and

Conference Reports

the Mackenzie. Geochemical and petrological data suggest that dykes in the former were fed from magma chambers situated at relatively shallow depths of less than 10–12 km (W.A. Phinney and D.A. Morrison, U.S.A.). These magma chambers were fractionating, while being continually replenished from the mantle. The observation that individual dykes in other Precambrian swarms retain remarkably constant compositions for hundreds of kilometres along strike, yet are compositionally distinct from neighbours, suggests that each was derived from a single magma chamber. The length of these dykes compared with the shallow depth of the assumed magma chamber requires an overwhelmingly lateral transport of magma through the crust.

The observations that many swarms appear to radiate from a focal region, that dykes tend to branch away from, rather than towards, these foci, and that near-focus ones tend to be more mafic in composition suggest that crustal magma chambers are concentrated in the focal region and that dykes propagate outwards from the focus. Swarms associated with Deccan volcanism in India appear to radiate from an area known to be the location of a migrating hot spot when the dykes were formed (S. Bhattacharji, U.S.A.). It has long been held that in Iceland magma is transported laterally in dykes during episodes of caldera activity associated with shield volcanoes such as Krafla. However, A. Gudmundsson (Iceland) suggests that the dominant mode of injection, which is not associated with individual volcanic centres, is vertical.

Most mafic swarms are assumed to have been formed in rifts associated with intraplate or plate margin activity, but new evidence suggests that some Precambrian swarms were injected during times when the adjacent crust was in active compression. An example of this behaviour is the 1.235 Ga Sudbury dykes in Canada that were emplaced when the nearby Grenville Province was developing, according to A. Davidson and K.M. Bethune (Canada). Another example presented by H.C. Halls and H.C. Palmer (Canada) is the 2.04 Ga Kapuskasing swarm, which appears to have been

formed during crustal uplift and tilting along the eastern thrust fault of the Kapuskasing structural zone in northern Ontario. A search for geochemical and petrological differences between these dykes and those known to have formed in extensional environments may thus provide general criteria for the recognition of the tectonic setting of dyke swarms.

Concerning the three-dimensional aspect of dykes, those of the Sudbury swarm that can be traced 5 to 10 km into the Grenville Province have been uplifted about 20 km. They become significantly thinner and also more curved and bleb-like, a possible consequence of their greater depth of emplacement, according to Davidson and Bethune. The recognition of dykes that have been emplaced at large crustal depths is also relevant to studies of crustal rheology as a function of depth and to the question of lateral magma transport. Kuehner and Green have recently elsewhere that the petrological composition of dykes emplaced at large depths can be reproduced by equilibrium phenocryst assemblages of chilled margins. However, Halls and Palmer showed that there may be other manifestations, one of which may be low levels of deuteric alteration and in thin section, the presence of blue-green amphibole rimming pyroxene and a tea-coloured cloudiness in plagioclase.

This phenomenon is seen in two dyke swarms that cross amphibolite to granulite facies rocks within the Kapuskasing structural zone. Outside the zone the dykes have clear plagioclase, often intensely speckled by low temperatures hydrous alteration products, and secondary amphibole of a green fibrous variety. Because the dykes within the Kapuskasing zone are fresh and undeformed, their anomalous mineralogy may be the result of crystallization under conditions of higher pressure and slower cooling. It is perhaps significant that the hyperite dykes of Sweden reviewed by Z. Solyom (Sweden), which are notable for their darkly pigmented plagioclase, occur in the Protogine Zone, a region of faulting that forms the eastern boundary of the Sveco-Norwegian Province in the Baltic Shield and across which there is a westward increase in metamorphism from amphibolite to granulite facies.

A topic for continuing debate is the relationship of mafic dykes to rapakivi granites. Both in Finland and Sweden, mafic dyke swarms appear to be spatially related to rapakivi plutons and commonly on strike with them. For example, I. Laitakari (Finland) described the Häme swarm in Finland, which is composed of more than 400 members and is about 150 km long. The dykes may have been emplaced into fractures opened by the rapakivi plutons because maximum widths gradually decrease from 1000 m to 10 m away from the plutons. Some of the dyke-parallel fractures near the plutons are filled with rapakivi magma, and sometimes there are composite dykes. It seems that mafic magma was available both before and after granite emplacement.

Anorthosite bodies associated with the granites and the presence of plagioclase megacrysts in the dykes suggest that the production of anorthosite is a factor common to the genesis of both diabase and granite. In southeastern Sweden, a set of northeast-trending dykes are geochemically similar to dykes in southwestern Finland about 500 km to the northeast, which appear to be spatially associated with rapakivi granites.

R. Gorbatshev (Sweden) discussed recent advances in mapping Swedish dykes. Sm-Nd studies by L. Johansson and A. Johansson (Sweden) have shown that a member of the hyperite dykes that closely follow faults in the Protogine Zone of southern Sweden is 1177 ± 38 Ma old, contrasting with 1500 Ma hyperite dykes farther north, with which they were previously correlated. A new Sm-Nd date of 924 ± 26 Ma has also been obtained by Johansson and Johansson for the N-S

Figure 2: Examining dyke swarms on aeromagnetic maps of Sweden.

Conference Reports

trending Blekinge dykes, most of which occur immediately to the east of the Protogine Zone. This swarm follows a gently arcuate path about 750 km in length, more or less independent of local faulting, but parallel to the Protogine Zone.

Dykes of comparable age parallel the Grenville Front on its northern side in Canada and may have a similar origin. Solyom reported that the 1.2 Ga Central Scandinavian Dolerite Group, the 1.17 Ga southern hyperites, the 0.92 Ga Blekinge swarm and the 0.3 Ga dykes along the Tornquist Line all have tholeiitic affinity but high Na, K, Ti and P. Thus despite the age differences, the similarity in chemistry suggests that all the swarms formed in similar tectonic environments and are probably rift-related.

A most important scientific advance in the study of dykes has been the recent acquisition of high precision U-Pb dates from zircon and baddeleyite in mafic intrusive rocks. The

precision, typically a few million years in Precambrian dykes, will have tremendous impact in the study of dyke swarms and may be able to define not only the period of time represented by the igneous event but perhaps even the spatial evolution of a particular swarm.

Future meetings of IGCP 257 (see Coming Events, this issue) are now scheduled for Santa Fe, New Mexico, as part of the IAVCEI Congress (June 25-July 1, 1989), for western Scotland (August-September 1989), and Australia (July 1990). A 106-page volume containing abstracts and a summary of the Lund meeting together with a field guide to the mafic dykes of southern Sweden is available (as IGCP 257 Technical Report 1) from the writer.

H.C. Halls
Geological Sciences, Erindale College
Mississauga, Ontario
Canada L5L 1C6

Pacific Neogene Events: Timing, Nature and Interrelationships

Shizuoka, Japan, 3 - 6 October 1988

The recognition, characterization, and timing of key climatic, tectonic, and evolutionary events recognizable in Neogene strata and fossil assemblages from throughout the Pacific and the eastern Indian Ocean are the concerns of IGCP Project 246. Recent progress was reviewed in October at a meeting organized by R. Tsuchi of Shizuoka University, the project leader, and sponsored by the Japan Society for the Promotion of Science and the Fujiwara Foundation of Science.

Four main types of events were discussed: a mid-Neogene climatic optimum at about 16 Ma; major Neogene cooling and upwelling events at about 15-13 Ma, 6-7 Ma, and 2.5 Ma; the opening of the Sea of Japan at about 15 Ma and related Pacific tectonic events; and events of rapid biotic evolution

that occurred during the middle Miocene for microfossils and during the late Pliocene for molluscs. These events are recognized in strata exposed around the margin of the Pacific Basin as well as in deep-sea sediments from the Pacific Ocean and Antarctic region by changes in fossil assemblages, oxygen isotope data of benthic foraminifers, paleomagnetic evidence, and study of the composition and distribution of sediment types.

Naturally, geological work has progressed further in some countries than others, so both descriptive reconnaissance and detailed specific studies were reviewed in workshop sessions. IGCP 246 is now expanding to include nonmarine investigations, and palynological and paleobotanical data were presented from China, Vietnam, and Japan.

During a field trip to the nearby Izu Peninsula, a terrane that drifted northward from the low-latitude North Pacific and collided with the Japanese mainland in the Early Pliocene, participants saw how tectonics and paleoclimatology can be integrated. This showed that the occurrence of tropical larger foraminifera in post-early Middle Miocene strata of the Izu Peninsula could only be due to their deposition in the low-latitude North Pacific and its subsequent northward drift.

During the workshop, participants were encouraged to write down their evidence for climatic, tectonic, and evolutionary events on time-space charts for the Pacific Basin. These preliminary charts will be carefully edited in the next few months, distributed to other IGCP 246 members, and published in perhaps 1-2 years time.

J.A. Barron
U.S. Geological Survey
345 Middlefield Road
Menlo Park, CA 94025
U.S.A.

Figure 1: Workshop participants compiling time-space charts for Pacific Neogene events.

Global Bioevents: Abrupt Changes in the Global Biota

Boulder, U.S.A., 16 - 23 May 1988

The analysis of global bioevents and, in particular, the dynamics of mass extinction continue to be major topics in international geosciences. This was evident at the third international conference of IGCP Project 216 (Global Bioevents in Earth History), sponsored recently by the University of Colorado and the International Palaeontological Association (IPA). The interaction between the 100 paleobiologists, paleoceanographers, paleoclimatologists, stratigraphers, sedimentologists, geochemists and geologists from 13 nations in attendance and concerned with both terrestrial and extraterrestrial phenomena gave great depth to the discussions. Large-scale biological models and causal mechanisms were reviewed, and bioevent dynamics were traced backward through time, from the Holocene extinction event to the Ediacaran-Tommotian boundary at the base of the Cambrian System, though the major emphasis was on Cretaceous time.

An excursion to the Raton Basin of Colorado and New Mexico gave participants an opportunity to examine and sample in detail the most completely preserved known record of the K-T boundary interval in continental facies (Fig. 1). An integrated record of plant extinction, rare element geochemistry (including Ir enrichment), and shock (impact) metamorphism has been documented here in great detail, allowing interpretation and modelling of the continental K-T extinction event.

Participants also examined the Pueblo, Colorado, exposures of the Late Albian-Middle Turonian Greenhorn Cyclothem, containing the best documented Cenomanian-Turonian (C-T) extinction interval in North America. The evidence is well displayed here of diverse bioevents associated with regional and global marine anoxic events, rapid immigration-emigration of warm southern water masses into the Western Interior Basin, and anoxic bottom water overturning. Emphasis was on the centimetre-scale study of the 2.5 Ma C-T mass extinction interval, including large stable-isotope fluctuations, the Bonarelli oceanic anoxic event, and iridium enrichment layers.

During the conference, major forcing mechanisms for global bioevents were sought in large-scale, short-term, climatic, oceanographic, chemical and tectonic events acting on typically "perched" Phanerozoic biotas with much narrower adaptive ranges than today. This biotic sensitivity reflects adaptation to predominantly warmer, more equable, maritime-dominated climates and higher global sea level, without permanent polar ice, that characterized over 90% of Earth history. Modern climate models provide guidelines for measuring large-scale fluctuations in Earth-ocean systems, but fall short of being able to duplicate precisely very short-term paleoclimate events, such as those related to impacts.

Extraordinarily large-scale and rapidly fluctuating disruptions in oceanic temperature, chemistry, circulation and stratification, as measured through stable isotope and rare element geochemistry, were associated with most global biotic crises throughout time. Possible causes include rapid vertical advection of toxic and anoxic waters from oxic, nitric, and sulphatic redox zones at and below the pycnocline, rapid mixing due to meteorite impact, rapid changes in ocean salinity during exceptional periods of evaporation and salt formation (saline giants), and abrupt large-scale

Figure 1: The K-T boundary claystone (2.5 cm thick kaolinitic layer just below pen point) at the Starkville North site, Raton Basin, New Mexico. Immediately above this layer is the thin dark boundary bed, which contains shocked quartz grains and high concentrations of Ir. Photo by C.L. Pillmore.

redox changes. Widespread, short-term atmospheric changes may also have resulted from major bolide impacts and rapidly emplaced atmospheric dust, smoke and water vapour, as well as from extensive mantle degassing associated with volcanism and emplacement of flood basalts, possibly resulting from impact.

A relationship between impacting and extrusive volcanism was suggested by the correlation of all major flood basalts during the last 250 Ma with cycles or quasi-cycles of clustered impacts and associated mass extinctions. Multiple bioevents of both a destructional character, involving mortality, extinction, ecological and biogeographic loss, and of a constructional nature, involving adaptive radiation, ecological and biogeographic gain, are commonly associated with each major interval of complex climate-ocean perturbation.

Discussions on the dynamics of large-scale bioevents through time focused on the biological responses to major environmental perturbations. Researchers seek signs of

Conference Reports

environmental crisis in the fossil record, and they look for understanding of extinction, survival, origination and evolutionary dynamics during mass extinction intervals. Dynamic models for rapid, large-scale extinction events associated with major environmental shifts were developed from modern three-dimensional models of nuclear winter phenomena involving short-term (weather), longer term (climate), and very long-term predictions (stratospheric ozone change) for major global perturbations affecting life. Whereas giant fire events would most closely fit nuclear winter models, predictions can also be made from them for the effects of large volcanic eruptions and large impacts. New data and models for patterns of survival during mass extinction events demonstrate that older theories, suggesting that only ecological generalists survived global crises, are much too simplistic.

The diversity of genetic, ecological, and morphological groups that rapidly repopulated the Earth after mass extinction strongly suggest that ecologically and genetically diverse groups survived these crises. Survivors utilized many "strategies" related, for example, to adaptive flexibility, reproductive and dispersal modes, migratory abilities, trophic characteristics, population dynamics, changing evolutionary pathways, habitat range, biogeographic dispersion, use and availability of refugia and dormancy. Initial studies of Cretaceous fauna support this assumption and reveal a pattern of repopulation from disaster species-dominated communities, through opportunist-dominated, eurytope-dominated, to increasingly diverse and more equable normal marine communities.

In terms of extinction theory, studies of individual taxonomic groups have yielded a complex picture of extinction dynamics. Thus rampant paludification may bring about terrestrial land plant extinction, change the CO₂ balance in the atmosphere, and cause climatic cooling without other forcing mechanisms. Major plankton and larger benthic foraminifer extinctions may, overall, be out of phase with the Raup-Sepkoski cycles of mass extinction. Indeed, the evolution-extinction history of foraminifera was shown to

correlate more with Earthbound phenomena than with extraterrestrial cycles. Plankton also provide evidence for other, non-extinctional bio- and geo-events; horseshoe-shaped coccoliths, for example, only form in response to global salinity crises.

The largest single group of papers dealt with the patterns and environmental dynamics of the Cretaceous-Tertiary interval (Fig. 2). The concept of a single catastrophe related to the large bolide impact at the Cretaceous-Tertiary boundary was discarded in favour of a more complex, dynamic Maastrichtian-Danian boundary interval, which involved multiple impacts and mantle degassing or emplacement of flood basalts as a possible result of impact. These events were superimposed onto stressed global environments that resulted from a large-scale eustatic fall, regression, and loss of shallow shelf and epicontinental marine habitats. Massive physical and chemical disruption of the oceans resulted in a stepwise mass extinction among both Cretaceous and Early Paleocene taxa.

Evidence was presented to suggest that the Deccan Traps of India, and other flood basalts associated with roughly cyclic mass extinction-impact intervals, might themselves be caused by mantle perturbations associated with giant impacts. This is supported by the discovery of a small impact crater associated with the Deccan Traps, by evidence for a giant tsunami and mantle degassing associated with a large marine impact in the Gulf of Mexico and Caribbean area at or very near to the K-T boundary, and by the occurrence of multiple Ir spikes at and above the boundary in the Brazos River, Texas, sections.

New and more precise analyses of published taxonomic data, and high-resolution (centimetre-scale) sedimentologic and paleobiologic analyses of complete North American, Mexican, and Spanish K-T boundary sections presented at the meeting support a complex stepwise extinction history that began about 2 Ma before the Tertiary Period with the abrupt disappearance of tropical rudistid-dominated reef biotas. This was followed by stepwise loss of ecologically diverse Gulf Coast molluscan-dominated biotas, deep water low-oxygen inoceramid bivalve-dominated communities, and specialized calcareous plankton prior to the main K-T boundary event.

The discovery of progenitor Paleocene molluscan faunas in the subtropical Late Maastrichtian rocks of Mexico, and of abundant Cretaceous species-lineages surviving into the Paleocene from Mexico to Alaska, further supports the concept of stepwise, ecologically graded K-T extinctions spanning the roughly 3 Ma-long Late Maastrichtian to Early Paleocene interval. Studies presented on the recovery of Caribbean Province molluscan biotas following mass extinction events revealed that most of the evolutionary time between extinctions was spent attaining earlier levels of diversity and ecological complexity over a period of 25-30 Ma, even though early radiations were rapid.

Reports on intra-Cretaceous extinction patterns concentrated on the C-T boundary interval and its complex extinction event. This is characterized by large-scale, global stable isotope fluctuations, which reflect rapid shifts in temperature and ocean chemistry, disruption of the carbon cycle, a global oceanic anoxic event (the Bonarelli Event), and massive climate perturbations. An interval of Ir enrichment reported from the latest Cenomanian of North America and at the C-T boundary in Poland contains evidence for both deep mantle and extraterrestrial sources for the Ir. A shocked quartz layer, associated with Fe-P stromatolites and Ir enrichment in Poland, and a major volcanic event in Germany were also documented for this boundary.

Figure 2: The K-T boundary at the base of the dark, lignitic layer (just below knee level) in Buffalo Jump Park, Alberta. Photo courtesy of D.A. Russell.

Conference Reports

New analyses of the C-T extinction event continue to show a stepwise pattern spread over about 2.5 Ma, which began with an abrupt decline in reef-building rudistids in the Caribbean Province during the Middle Cenomanian, correlative with a major temperature drop in Tethys and a possible restriction or disappearance of a warmer, more saline, core tropical climate zone ("Supertethys") favouring rudistid reef development. All later Cenomanian mollusc-foraminifer extinctions are precisely or closely correlated to Ir enrichment levels, to major stable isotope disruptions, or both. This also supports the concept of complex extraterrestrial and Earthbound (oceanographic, climatic) causal mechanisms for mass extinction.

Analyses of Paleozoic-Mesozoic intervals have produced important new insights into the nature of hitherto little-known bioevents and mass extinctions. Relatively abrupt pulses of extinction of Mesozoic dinosaurs and other large reptiles were documented at 10-15 Ma intervals, with each pulse spread out over 2-3 Ma. Reptilian extinctions were characterized by rapid demise of both herbivores and carnivores as a result of competition, disease, and other factors, and they were associated with the immigration of many new species without any slowdown in evolutionary rate. Major biotic crises among larger benthonic foraminifera were closely correlated with large-scale anoxic events, with reproductive strategies determining survival potential. Polish colleagues reported evidence for a bolide impact at the Middle-Late Jurassic boundary, on the basis of an Ir spike, Ni-Fe spherules, shocked quartz, diaplectic glass, and possibly soot. These features are associated with stromatolitic crusts of possible hydrothermal origin and an abrupt moderate-level extinction among, especially, ecologically restricted marine faunas.

The Triassic-Jurassic (T-J) extinction boundary also shows evidence of environmental perturbation - two major negative $\delta^{13}\text{C}$ spikes within an overall negative $\delta^{13}\text{C}$ fluctuation. The lower of these is associated with the demise of conodonts at the Norian-Rhaetian boundary well below, and the second spike occurs 2 m above the T-J boundary extinction.

A review of the best data on the Permo-Triassic extinction interval - one of the most profound of the Phanerozoic - demonstrates that it was not abrupt nor pervasive among all major taxonomic groups. Extinction events (steps?) occurred below, at and above the boundary over several million years. These had different patterns among the major marine taxa involved and seem to be correlated primarily to Earthbound processes such as eustatic fall, or change in shallow marine ecospace, an oceanic salinity crisis, global cooling, nutrient supply changes, and biotic phenomena. Relatively short-term, global changes in Mississippian echinoderm faunas were attributed to biologically regulated phenomena (community reorganization) without major changes in paleoenvironment - an example of pseudo-mass extinction.

Among major Paleozoic bioevents, the Late Devonian mass extinction interval (the Kellwasser Event at the Frasnian-Famennian (F-F) boundary) has received the most high-resolution analyses. New data emerged at the conference that more precisely define the dynamics of this event. Short-term stepwise extinction was defined against a background of increasing extinction rates for 4 Ma prior to the F-F boundary, but speciation rates nearly kept pace with extinction until just prior to the F-F boundary when one or more extinction steps decimated the fauna.

The main Kellwasser extinctions were related to a global anoxic event with subsequent ocean overturn. In addition to the still-debated, stromatolite-associated Ir anomalies previously reported in Australia, Canadian researchers reported

evidence at various F-F boundary localities of an abrupt drop in sea level that produced microkarst in emergent platform carbonates, and a narrow boundary interval with trace-element enrichment associated with a high sulphur isotope anomaly. In some places there is an algaloncolite boundary bed with geochemical features indicating rapid oceanic overturn bringing "old" anoxic, N_2S -enriched bottom water abruptly onto shallow cratonic platform settings. These findings again open speculation about a F-F impact event, possibly in the ocean.

Major Silurian to Middle Devonian bioevents are expressed by abrupt changes in regional and global paleobiogeography. These were driven by rapid changes in ocean temperature and circulation associated with accelerated sea-level rise (e.g. Uralian event, Kacak event, Taghanic Event) and changes in plate configurations. Many smaller Silurian and Devonian bioevents are also recognizable. Ordovician-Silurian (O-S) boundary extinctions within graptolite facies were especially dramatic. Following a minor Late Ordovician event, an abrupt, large-scale extinction of graptolites occurred in association with Late-Ordovician glaciation, probable rapid fluctuations in the marine de-nitrification zone, and nutrient crises. Survivors of the O-S event were graptolites with small colonies and small zooids.

Intra-Cambrian mass extinctions, represented by three well-defined, and two probable biomeres, have very sharp stratigraphic boundaries and are now being recognized globally. These bioevents may represent temperature or oxygen crises, or both, among shelf biotas. Each biotope is characterized by abrupt loss of the most diverse, evolutionarily advanced members of pre-existing trilobite and brachiopod-dominated communities. Extinctions were immediately followed by brief introduction of and population bursts among ecological generalists with no known ancestors, and by opportunists derived from older biotas. Subsequently, more gradual radiation involved new, evolutionarily advanced forms extending up to the base of the next biotope.

The first Cambrian faunas (Tommotian) are separated from their Precambrian (Ediacaran) predecessors by a major mass extinction event, still poorly understood. The relatively abrupt extinction among the Ediacaran fauna of large, thin-walled, passively respiring integumented forms may have been related to a global anoxic event, as evidenced by increased C_{org} levels associated with a positive $\delta^{13}\text{C}$ spike. Early Cambrian radiation among smaller, initially naked, but later shelled Tommotian taxa possibly reflects adaptations to low eH and pH waters favouring small organisms with high relative surface areas, thin body walls, and active systems of respiration. A Gaian feedback loop was proposed to explain this change, enhanced by conditions that caused the Cambrian-Precambrian mass extinction.

From the meeting a series of recommendations emerged that stressed the need for more research and better models in several areas. These include the relative roles and possible relationships of large-scale volcanism and impact in disrupting ocean-climate systems during mass extinction intervals, and the need to explain very rapid shifts in temperature and chemistry that seem to be causal mechanisms for mass extinction. Also required are models for oceanic impact effects, the predominant type of Phanerozoic event leading to biotic crises, a comprehensive classification of extinction types and their relationship to origination, and definition and interpretation of the common denominators among mass extinctions as clues to causal mechanisms.

More work is needed to understand survival mechanisms, the nature of "Lazarus taxa," and processes of origination, radiation, ecological restructuring, and repopulation of event-affected biotopes following mass extinction intervals. The deep ocean record of mass extinction must be documented,

Conference Reports

starting with integration and contouring of various data across the C-T and K-T deep ocean environments, and the major geological effects of large-scale impacts need to be defined and related to mass extinction history. More emphasis should be placed on sampling strategies and very detailed interdisciplinary studies of bioevents at all scales, and on mass extinction intervals not already investigated in detail. Alternatives to impacts must be sought for bioevents, which still operate on a regional to global scale within very short time intervals.

The meeting concluded with a unanimous endorsement for continuation of the research being conducted under Project 216, and for close integration of this work with IPA activities. Possible new directions for IGCP-216 would include consideration of the past as a model for the present and future, and integration of bioevent research within the forthcoming Global Change program.

The next international conference of Project 216 will be held at Oxford, U.K., in early July 1990, and in 1989 several symposia and workshops will take place during the Washington Geological Congress and during several other international meetings. The proceedings of the conference are being compiled as a series of research papers to be published in 1989. Copies of the abstracts (\$10) and field guides (\$5) from this conference are available from the first author.

Erle G. Kauffman
Department of Geological
Sciences
CB-250 University of Colorado
Boulder, CO 80309, U.S.A.

Otto H. Walliser
Institut für Geologie und
Paläontologie
Universität Göttingen
Goldschmidt-Strasse 3
D-34 Göttingen, F.R.G.

The Kibaran of Central Africa

Lubumbashi, Zaire, 29 August - 9 September 1988

Until recently, the middle Proterozoic Kibaran belt of East and Central Africa was considered to be an intracontinental orogen that evolved through a cycle of rifting and sedimentation (about 1400-1200 Ma ago) followed by compressive phases between 1200 and 1100 Ma. Granites emplaced within the belt around 1000-900 Ma were thought to be the source of most of the region's economically important mineralization (Sn, W, Be, Li, Ta/Nb, and Au/Pt). These granites were regarded as generated in a Pan-African mobile belt that overprinted the Kibaran. Now, however, French and Zairean workers including J.-P. Burg, J.-P.H. Caron, A.B. Kampunzu, D. Kapenda and B. Rumvegeri propose a model involving evolution of the belt through a true oceanic stage with an Andean type margin followed by continental collision. This controversy considerably enlivens current discussions on the Kibaran, as was seen at the first field meeting of IGCP Project 255 (Kibaran Evolution and Metallogeny) of which the writer was the initial leader.

About 40 participants from nine countries attended a two-day conference, which was followed by an excursion to the Kibara Mts., the type area of the middle-Proterozoic

Figure 1: Project 255 co-leader A. Ntungicimpaye points to columns of unaltered potassic pegmatite between channels of albitization and greisenization which carry the tin (-niobium-tantalum) mineralization, Manono, Zaire.

Kibaran belt. During the meeting, D.P.M. Hadoto (Uganda) and A. Ntungicimpaye (Burundi) were elected co-leaders of Project 255, in order to strengthen and facilitate African participation.

The Lubumbashi meeting illuminated several aspects of the problem, relating to syn-sedimentary volcanic rocks (A. Ntungicimpaye and L. Tack, Burundi), the intrusive suite including I- and S-type granitoids (A.B. Kampunzu, Zaire) and mafic-ultramafic massifs (D. Kapenda, Zaire), the structural and metamorphic evolution (B. Rumvegeri, Zaire), geophysical models (B.R. Byamungu, Zaire) and the "post-tectonic" tin granite related mineralization (J. Brinckmann, FRG; S. Cauet, Belgium; W. Pohl, F.R.G.). A resolution of opposing interpretations, not surprisingly, was not achieved at Lubumbashi, and must clearly wait for more data.

The field excursion provided a cross section of observations on sedimentology, deformation, metamorphism, granitoids and mineralization of the Kibara Mountains. We travelled about 2500 km on roads and tracks, twice even crossing the upper Congo (Lualaba) River, by ferry. A highlight was a visit to Manono, apparently the largest tin (Nb/Ta) pegmatite being exploited anywhere in the world, and part of the Kibaran tin granite-related mineralization. The question was raised as to whether the tin granites are really independent of the Kibaran as L. Cahen proposed, or whether they were the products of a final Kibaran orogenic phase that affected both Kibaran suites and the molasse deposits. Other targets are the nature of the ubiquitous tectonic control on the mineralization, and the derivation of the gold (Pt) quartz veins. New data on fluid inclusions cast doubt on one interpretation that the gold is the epithermal equivalent of catathedral tin and tungsten.

The first number of IGCP Project 255 Newsletter-Bulletin is now available and can be obtained from the writer. The next meeting of the project will take place in western Tanzania in September 1989. There, the tin granites, related mineralization, mafic-ultramafic massifs and the eastern margin of the Kibaran will be discussed.

W. Pohl
Institute of Geology and Paleontology
Technical University
P.O. Box 3329
D-3300 Braunschweig, F.R.G.

Trends in Experimental Mineralogy, Petrology and Geochemistry

Bochum, F.R.G., 7 - 9 April 1988

What are the depths of magma genesis in the mantle? How hot are migmatites formed by partial melting in the continental crust? And how deep was the subduction of former sediments in zones of continental collision? These and many other questions of this kind are continually being asked by igneous and metamorphic petrologists throughout the world. Only laboratory studies on Earth materials or on simplified chemical systems can provide reliable answers.

These studies now require expensive, high-pressure equipment and maintenance, as well as often tedious and meticulous work. The days of quick, spectacular results are over. Thus many petrologists, especially in the U.S.A., are turning to more rapid methods of calculating mineral equilibria from thermodynamic data. On the other hand, and perhaps reflecting a more conservative attitude, scientists in Europe have been more hesitant to give up the newly gained techniques.

It is thus no surprise that the First Symposium on Experimental Mineralogy and Geochemistry was organized in France in 1986. This met with much interest and showed so much potential that the Second Symposium was held two years later in Bochum, West Germany, attended by about 200 scientists from around the world. This was organized by the Mineralogy Institute of Ruhr-University and sponsored by the French and the German Mineralogical Societies, by the newly founded European Mineralogical Union (EMU) and by the IUGS Commission on Igneous and Metamorphic Petrogenesis (CIMP).

The question of the interaction between fluids and rocks and minerals drew much attention. Although fluids are rarely preserved in rocks and minerals, and then only in very small quantities, it is quite evident that these highly mobile phases (gases, melts, or supercritical fluids) must have played a paramount role for many processes of petrogenesis. They may be a source of, a solvent for, or merely a coexisting phase of solid earth materials. In contrast to studies on natural products, experimental research can exert direct controls on the type of interaction between solids and fluids and can directly approach problems such as element fractionation or solubility.

In this connection M. Lagache (France) summarized present knowledge on the distribution of elements between silicates and hydrothermal solutions, and I.D. Ryabchikov (U.S.S.R.) pointed out that the hydrous fluids coexisting with K feldspar and jadeite at high pressures have a peralkaline chemistry. The close interrelationship between experimental geoscientific research and urgent problems of nuclear waste disposal became evident from the study by A. Godon and colleagues (France) on the reactivity of glass in contact with geological materials and in the presence of fluids.

In a session on the properties of fluids themselves, E.U. Franck (F.R.G.) reported for the first time on a synthetic system of geologic importance ($H_2O-NaOH$) for which thermodynamic and other properties were determined over the whole compositional range up to 400°C and 4 kilobar. J.D. Frantz (U.S.A.) presented a new method of analyzing minute fluid inclusions within minerals by X-ray fluorescence using synchrotron radiation.

Despite the fact that chemical equilibrium has quite often not been attained in natural rocks, its definition and determination as a function of pressure and temperature in multicomponent natural systems remain very important tasks for experimental petrology. R.C. Newton (U.S.A.) summarized equilibrium data bearing on the origin of granulites in the lower crust, and H.J. Massonne (F.R.G.) presented data of interest for medium and high-grade metamorphism of pelitic rocks in the upper crust. The stability limits of ellenbergerite, a spectacular new high-pressure mineral were discussed by C. Chopin (France) who had discovered it in the sensational coesite-pyropite rocks of the Western Alps.

Figure 1: Martine Lagache, President of the newly founded European Mineralogical Union (EMU), during her opening address at the Bochum Symposium.

The dramatic influence of only traces of water on the mechanical behaviour of quartz was discussed independently by M.S. Paterson (Australia) and P. Cordier (France). New results were presented by H. Kern (F.R.G.) on the experimental deformation of fine-grained limestones, which can be used to interpret preferred crystal orientation in naturally deformed rocks in terms of their strain history.

Using synchrotron radiation, R. Boehler (F.R.G.) has measured the density of a heated iron wire in a diamond anvil cell under the P-T conditions of the core-mantle boundary. S.L. Webb (Australia and F.R.G.) discussed a new model for the anomalous pressure dependence of the elastic moduli of orthopyroxene, and also the frequency dependencies of compressional and shear viscosities of melts. The title of the symposium might well have included, therefore, experimental geophysics, as the various branches of geoscience continue to merge.

The contributions to the topic of crystal chemistry were mainly confined to experimental studies on silicates, the most common of the rock-forming minerals. C.M. Graham (U.K.) and W.V. Maresch (F.R.G.) reviewed recent progress

Conference Reports

in the knowledge of amphiboles, perhaps the most complicated group of geologically important silicates. They emphasized the dramatic influence of structural defects (Realbau) on crystal chemistry and stability relations, as well as the difficulty of controlling these defects in synthesis work. These conclusions are obviously also of interest to the consideration of reaction kinetics that followed.

Whereas mineralogists and petrologists have in the past emphasized the importance of equilibrium studies for solving petrogenetic problems, they now realize that the time factor is a new dimension for understanding rock-forming processes and thus for geodynamics as a whole. Again, silicate systems are the main focus of interest for work on kinetics. C. Giampaolo (Italy and U.K.) reported on the dehydration kinetics of Mg-cordierite, while N. Le Breton (France and F.R.G.) presented most surprising results on the experimental introduction of CO₂ into the same mineral. She appears to have found, for the first time in experimental mineralogy, a case of an oscillating chemical equilibrium, with the CO₂ contents of cordierite fluctuating with time at constant pressure and temperature. O.W. Flörke (F.R.G.) discussed the kinetics of exchange of traces of Al for Si in the structures of high and low quartz; D.C. Rubie (U.K.) reviewed the kinetics of partial melting of a quartz muscovite schist, in a discussion that made a convenient link to the last topic.

Although the main granite controversy is over, much remains to be done in understanding the origin of granitic and other melts within the lithosphere. W. Johannes (F.R.G.) discussed the experimental constraints on the formation of granitic melts, especially as a function of variable volatile activities. E.S. Persikov (U.S.S.R.), on the other hand, stressed the effect of volatiles on the properties of magmas. P.J. Wyllie (U.S.A.) concentrated on igneous processes in subduction zones by emphasizing diffusion and

reaction contamination resulting from interactions among magmas, vapours and rocks brought together essentially by tectonic mechanisms. D.L. Hamilton (U.K.) presented new data on the distribution of trace elements between immiscible silicate and carbonatite melts.

One sensational finding reported at the meeting by Sobolev (U.S.S.R.) with implications for global geodynamics was the discovery of inclusions of minute diamonds in garnets of a metasedimentary gneiss from northern Kazakhstan. He interpreted the diamonds as representing carbon of former organic matter present in the original sediment. On the basis of the known diamond/graphite equilibrium curve, this implies that sediments, probably of continental crustal origin, may have been subducted to mantle depths of at least some 130 kilometres.

Another highlight was the video presentation by A. Henk and V. Lorenz (F.R.G.) of laboratory experiments modelling fluidization and explosive volcanism that lead to the formation of maars and diatremes.

The symposium left an impression of an extremely active and powerful, modern branch of the earth sciences, and we now look forward to the third symposium, which will take place in Edinburgh, Scotland, during the spring of 1990. A selection of papers given at the Bochum symposium will be published in volume 1 of the new European Journal of Mineralogy to appear in 1989.

Werner Schreyer
Institut für Mineralogie
Ruhr-Universität
Bochum, F.R.G.

Lake Sediments Reviewed in China

Beijing, 31 July - 4 August 1988

IGCP Project 219 was established to foster international communication between geoscientists studying both modern and ancient lake deposits and to develop a data base on lake deposits worldwide. As part of the first major international conference on sedimentology to be held (by IAS) in China, IGCP 219 organized recently a workshop for a large group of Western specialists to hear the details on numerous important lacustrine deposits in that country. Chinese workers presented 71 papers and all major lake deposit research groups in China were represented at the meeting, including strong contingents from the Ministries of Petroleum and Geology, the China University of Geosciences (Beijing and Wuhan), and various geological institutes of the Academia Sinica.

There is great interest among Chinese geologists in lacustrine sedimentology and stratigraphy, because most of the current fossil fuel exploration and production in China occurs in nonmarine Mesozoic and Cenozoic basins. Discussions of the coal-bearing Mesozoic Ordos Basin and the Tertiary Bohai and Jinyang Rifts with their extensive oil and gas reserves focused on models of facies geometry that could be applied towards future exploration. The recognition that many Chinese petroleum reservoirs occur in sub-lacustrine fan turbidites has generated much research into subaqueous gravity flow processes and deposits. Li Sitian, Yang Shigong and Xia Wenchen showed how gravity flow events of various magnitudes in the Early Fuxin Basin can be recognized through their 3-D geometry.

Figure 1: Examining carbonate-gypsiferous paleosols of the Paleogene Guanzhuang Formation, Dongjiazhuang, China.

Conference Reports

The growing debate among rift-lake sedimentologists about the relative importance of axial as opposed to lateral sources of fluvial sediments for late syn-rift lake infilling was addressed by Wang Defa, Zheng Junmao, Deng Hongwen and Pang Ming. They argued that for the majority of Mid-Jurassic to Early Tertiary Chinese basins, most sediments can be shown to be derived from transverse proximal sources as opposed to axial sources. Localized sand sources for deep water channels was also supported by the work of Qi Xingyin and Shou Jianfeng in the Dongpu Basin of the Bohai Rift.

Studies by Dong Zhenuan in the Northern Sichuan Basin, Tang Zeyao and Chin Jigao in the Central Sichuan Basin and Huang Xingzhen, Yang Xiaorong and Lei Zhenfeng in the Qaidam Basin showed the importance the Chinese attach to developing sedimentological models of lacustrine carbonates as reservoir facies. Some of the Sichuan coquinas show remarkable similarities to the productive Lagoa Feia Formation coquinas of the Campos Basin of offshore Brazil.

Papers on isotope geochemistry fluid inclusion research, clay mineralogy and laser-induced fluorescence illustrated that Chinese sedimentology has moved far beyond granulometry. Xue Shuhao, Wang Keyou and Ying Xudong showed how local bedrock lithology and depositional environments within the Dongpu Basin controlled the distribution of trace and rare-earth elements. Chinese studies have enormous potential for resolving some of the controversies surrounding the interpretation of biomarker data in paleoenvironmental (and particularly paleochemical) analysis, because of the tremendous variety of both modern lakes and lake deposits in China. Luo Binjie integrated data on carbon isotopes with pristane/phytane ratios as an index of paleosalinity. In collaboration with Meng Qianxiang and Wang Youxiao, he was also able to show that the distribution of various bi-cyclic sesquiterpanes can also be used to distinguish sediments deposited in waters of differing salinity.

Several speakers made broad syntheses relating the evolution of lake basins in eastern Asia to regional tectonics. For the Ordos Basin, Chen Channing, Wang Shousong, Ke Baojian, Chen Zhiming, Chen Jingshan and Jiang Haoren argued that the periodic reactivation of the marginal thrust-fold belt controlled both the cyclic sedimentation pattern observed in the basin and the apparent migration of the depocenter towards the North China Platform. They summarized the differences between this foreland basin and those of the North American Cordillera, proposing the term "collateral foreland basin" to describe Ordos-type basins.

Rift basin tectonics and sedimentation in the Bohai Bay region were summarized by Xin Quanlin, Zhu Xiaomin, Zheng Qing and Zhang Jie. They illustrated the asymmetric "dustpan-shaped" half-graben configuration common to many of the early Tertiary lake basins of northeastern China. Structural provinces within these lakes are associated with predictable lithofacies associations in a very similar fashion to those in the modern, large rift-lake basins of Central Africa, despite some significant differences in tectonic style.

The attention being paid to modern Chinese lakes as paleoclimate indicators, as analogs for ancient lakes and because of their evaporite mineral potential is particularly interesting. The closed lakes of Qaidam Basin and Lake Qinghai have been under investigation by Chinese sedimentologists for some years. As shown by Yue Changshuo and Qu Zheng for Gas Hure Lake, the evolution of western Chinese closed basin lakes is closely analogous to those of the Great Basin of the U.S.A., both resulting from a combination of fault block segmentation and increasing late Quaternary aridity. G. Lister (Switzerland) illustrated the complexities in predicting lake level changes due to changes in climate. He

showed for Lake Qinghai that lake morphometry strongly controls the equilibrium level under a given set of precipitation and evaporation conditions. Lister, Chen Kezao and others established a lake-level curve for Lake Qinghai for the late Quaternary using oxygen isotope data derived from ostracodes.

Qaidam Basin provides a natural laboratory for understanding the deposition and diagenesis of economically important evaporites. Sun Dapeng showed that borates in Xiao Qaidam Lake (primarily ulexite and pinnoite) form during early diagenesis, possibly related to periodic freezing of groundwaters. For Dabsan Lake, also in Qaidam Basin, he traced the brine concentration pathway leading to the formation of extensive potash deposits.

Talks by foreign participants at the workshop sparked some fascinating discussions comparing the lacustrine sedimentology of rift and foreland basins elsewhere with the data presented by the Chinese. E. Robbins and D. Textoris (U.S.A.) interpreted the environments of deposition of Late Triassic Deep River (rift) Basin coals of North Carolina. They proposed that lake-level fluctuations regulated the water table on the lake margin, which secondarily controlled the distribution of Triassic wetlands. The case for a non-marine-lacustrine origin for parts of the Mid-Proterozoic Belt Supergroup of Montana was presented by D. Winston (U.S.A.) who interpreted certain fine-grained, rhythmically bedded sequences as playa-lake deposits, in opposition to earlier tidal-marine interpretations.

For Swiss lakes, K. Kelts (Switzerland) presented some provocative, initial data, which demonstrated that Sr isotope stratigraphic methods might be applied to solving single-basin correlation and basin interconnection problems in the lacustrine rock record, as has been previously done in marine strata. P. Allen (U.K.) showed that most of the Devonian Orcadian Basin lake deposits previously suggested to indicate deep-lake conditions were, in fact, laid down under very shallow-emergent conditions.

Summarizing the results of the large working group, on Spanish lake deposits, L. Cabrera (Spain) showed how the African-European collision generated numerous lake basins, both foreland and rift on the Iberian Plate, many of which are petroliferous or coal-bearing. P. Homewood (Switzerland) and L. Reggiani utilized a thrust-loading model for the edge of the Alpine Foreland Basin to explain small scale (tens of metres) cycles in Oligocene lake deposits. In his summary of the Pliocene-Recent transtensional basin deposition of the Dead Sea region, I. Zak (Israel) demonstrated that cyclic sedimentation of evaporites and clastics have characterized deposition within the basin during both marine and lacustrine intervals.

Structural differences between recent stromatolites from Lake Tanganyika and those of East African saline lakes, according to A. Cohen and S. Awramik (U.S.A.), may be due to differences in microbial communities, which affected the proportion of sediment binders to calcite precipitators. T. Kreuser reviewed Permo-Triassic Karoo basin deposition in Tanzania. Karoo coal swamps formed on prograding delta plains and were subsequently preserved within a series of half-graben basins.

A post-meeting field trip to the Paleogene rift-lake deposits of the Shengli Oilfield in the Dongying Depression, which yield 23% of China's oil production, and other Tertiary lake deposits of Shandong Province gave foreign participants a brief opportunity to examine some of the lacustrine strata discussed during the meeting. The diatomaceous and richly fossiliferous paper shales of the Shanwang Formation are closely analogous in many respects to the Oligocene Florissant Beds of western North America. Future comparative

Conference Reports

Figure 2: Discussing oncolitic packstones of the Guanzhuang Formation near Pingyi.

taphonomic studies from these types of exceptional fossil assemblages could provide valuable insights into the process of lacustrine fossil preservation. In Pingyi Basin, participants had an opportunity to see an exceptionally diverse suite of early Tertiary lacustrine carbonate lithologies, broadly correlative with the Dongying Depression deposits.

Chinese sedimentology is unquestionably entering a new era. Many of the younger sedimentologists are very sophisticated in their approach to science, and many want the organizational as well as the technical tools to pursue their craft more openly and effectively. The appearance of interdisciplinary studies such as seismic facies analysis at this meeting was both noteworthy and heartening. Zhang Jie, Xin Quanlin and Yang Yunling showed, how thin sand bodies deposited as delta-front, lacustrine turbidites could be recognized from seismic reflection data in the Dongying Depression.

The development of greater scientific interchange among scientists and between institutes is certainly a positive sign for Chinese geology. Meetings such as the IAS conference are extremely important at the moment not only for developing cooperation between Chinese and foreigners but also for allowing the Chinese geoscientists to exchange data among themselves. While it is too early to talk of a Chinese "school" of sedimentology, this may well develop in time, given a continuation of the kind of exchange that marked this successful conference.

A.S. Cohen
Department of Geosciences
University of Arizona
Tucson, Arizona 85721
U.S.A.

African Magmatism: Mesozoic to Present

Giens, France, 26 - 28 September 1988

To follow-up its 1986 Lubumbashi conference on magmatic rocks in extensional zones, IGCP Project 227 (Magmatism and Evolution of Extensional Regions of the African Plate) recently held its second international meeting, in Giens by the Mediterranean. Organized by the Laboratory of Magmatic Petrology of the University of Aix-Marseille III, and jointly sponsored by Unesco, the French IGCP Committee, CIFEG, IUGS and the Ministry of National Education (Paris), the meeting was attended by some 60 geoscientists from 12 countries.

Setting the tectonic framework for the discussion, J. Angelier (France) presented a comprehensive review of the structural and mechanical characteristics of rifting processes. These were illustrated by a variety of examples from continental to oceanic stages: Basin and Range, Rhine graben, Gulf of Suez, Gulf of California, Red Sea, Iceland. The important relation between the direction of extension and the rift axis was stressed as was the interaction between tectonics and magmatism.

The tectonic setting and magmatic evolution of the Rio Grande Rift were taken by R.N. Thompson (U.K.) and co-workers as a guide to a better understanding of the African extensional magmatism. They emphasized that the occurrence of a linear zone of epeirogenic uplift above an asthenospheric upwelling, at least 1,200 km in length beneath the Rio Grande Rift, is analogous to the linear zone of regional updoming along the Ethiopian and Kenyan segments of the East African Rift system. Thus, linear mantle upwelling was favoured rather than a plume mechanism.

One of the most striking points raised during the Conference was the existence of "two Africas" as far as the rift system is concerned: West and Central Africa (the Mid-African Rift

System), characterized by little or no magmatism, and East Africa, where Cenozoic magmatism is widespread. The former system extends from Nigeria (Benue Trough) through Niger and Chad, eastwards to Sudan and to the Mediterranean Sea. It is linked to the fracturing of Gondwanaland and opening of the South Atlantic Ocean. This rift system was analysed by J.D. Fairhead (U.K.) who suggested possible reasons to account for the lack or scarcity of volcanism within it: thermal state of the upper mantle beneath the rift, degree of crustal extension, and the role of isostasy in developing crustal structures. M. Popoff and P. Baudin (France) gave an overview of the different stages of the tectonomagmatic and sedimentary evolution of the Benue

Figure 1: Questioning models for African rift magmatism at Giens.

Conference Reports

Trough from Late Jurassic to Tertiary times. Despite the scarcity of magmatism, each igneous episode is significant in terms of the successive extensional tectonic phases (rift faults, transfer faults).

Another good example of structural control on magmatism was provided by A. Pouclet (France) and co-workers in Eastern Niger (Air and North Termit areas). In these areas, Cenozoic volcanism is related to the Hoggar-Lake Chad structural axis, and the magmatic types show a temporal succession (from alkaline to transitional lavas) clearly controlled by the evolution of the extensional tectonic regime.

Some contributions dealt with more recent volcanics from Western Africa, such as the Quaternary alkali lavas from the Air massif (I.F. Hazell et al., U.K.) and from the Cameroon Line. In the latter, new geophysical and geochemical investigations were presented by J.D. Fairhead (U.K.) and B. Deruelle (France). The distribution of hygromagmaphile elements indicate, according to R. Kambou and others, a slight heterogeneity of the magma source. They do not correspond to any space-time evolution usually invoked to support conventional mantle plume models.

The East African Rift session also offered some excellent contributions analysing igneous activity in relation to its structural setting. G. Feraud (France) presented new ^{39}Ar - ^{40}Ar dates showing that the magmatism of the western border of the Arabian Plate is related to the early opening (28-22 Ma) of the Red Sea. Extensional episodes (accompanied by normal faulting and tilting of blocks) and development of dyke systems are closely linked.

On the basis of new tectonic and magmatic data, P.J. Vellutini (Djibouti) stressed the important role played by the Manda Inakir rift zone in the geodynamic evolution of the Djibouti region. This active rift zone is linked to the Asal rift by a north-south strike-slip fault system. It was proposed that a triangular "microblock" existed here, which would have been displaced toward the northeast in relation to the movement of the Danakil Horst. This new interpretation of this key area of the East African Rift was completed (A. Anis, Djibouti, and J. Demange, France) by recent information from three geothermal drill holes in the Asal rift. It has now been established that the main faults are vertical and reach more than 2,000 m in depth, and that oceanic crust has not been generated here. Asal is thus interpreted as the result of rifting in a continental environment and does not correspond to the emergence of an oceanic ridge.

G.R. Davies (U.K.) gave a detailed geochemical account of the Ethiopian Flood Basalt Province, showing that these lavas are mainly derived from sub-continental lithospheric mantle, with all but the earliest volcanism undergoing extensive crustal interaction. In his account of the volcanism of the South Afar Rift, G. Yirgu (Ethiopia), pointed out that southwards the amount of extension decreases, the alkalinity of the basalts increases, and the crust progressively thickens.

The complexity of magma evolution in the central Kenya rift was examined by R.A. MacDonald (U.K.) using three types of Quaternary to Recent associations: peralkaline trachytes, peralkaline rhyolites and their associated mildly alkaline basalts. He outlined the diverse mechanisms in operation here: partial melting of a subcontinental lithosphere combined with assimilation of continental crust for the basalts; crustal melting, facilitated by halogens and CO_2 in the case of the peralkaline rhyolites; and crystal fractionation and partial melting of basalt for the trachytes. A detailed study of the evolution of the peralkaline trachytic magma at Longonot volcano, central Kenya was provided by S. Scott (U.K.), who showed that the magmatic evolution of this volcano reflects the existence of

a zoned trachytic magma chamber where several differentiation and mixing events (between trachytic, phonolitic and basaltic magmas) occurred.

Farther south in the East African Rift, the Virunga chain (Zaire) was the subject of other presentations. Thus, P. Lestrade (France) described the petrology and geochemistry of the lavas (K-hawaiites) of a new parasitic cone (August 1986 eruption) belonging to the Nyamuragira magmatic system. For the ultra-alkaline lavas from the Nyiragongo, he also showed that a detailed sampling of the upper half of the crater wall allowed the distinction between different units made of melilitites and bergalites (Lower Unit), nephelinites (Middle Unit) and feldspathoidal aggregate lavas (Upper Unit). All these lavas are linked by complex petrogenetic processes (fractional crystallization, crystal flotation, gas transfer, magma mixing).

In contrast with previous models, it was proposed that the adjacent Nyiragongo and Nyamuragira magma systems evolved from different primary magmas. For the Kivu Rift, R.T. Lubala and M. Kanika (Zaire) focussed their attention on the space-time arrangement of the magmatic types with increasing rifting and on the consequences of this temporal change on the modelling of the asthenosphere-lithosphere dynamics during rifting.

For the basalts of the Kivu Lake area, investigations by C. Dupuy (France) and co-workers demonstrated mineralogical constraints concerning the mantle source (presence of phlogopite and apatite) and its degree of heterogeneity. P. Mohr (Ireland) and A.B. Kampunzu (Zaire) reviewed the problem of magmatic evolution and petrogenesis in the East African Rift System, emphasizing the differences between the Western and Eastern branches of the Rift (crustal extension, abundance of volcanism). They distinguished four major volcanic suites (ultra-alkaline lavas and carbonatites, alkali basalts, transitional basalts, and tholeiites) and discussed the mantle components that can be identified under the East African Rift. Estimates of the volumes of volcanic products provides evidence to the essential role of convective circulation in the underlying asthenosphere.

The characteristics of the source of the African extensional magmatism was a "leitmotiv" underlying many of the papers. D.K. Bailey (U.K.) stressed that no two continental rifts are alike in either volumes or sequences, thus ruling out the hypothesis of a massive upwelling of convecting asthenosphere. J.B. Dawson (U.K.) concentrated on mantle xenoliths from northern Tanzania and reviewed the question of upper mantle heterogeneity. These xenoliths show evidence of metasomatism through the presence of pargasite, biotite, and veins of katungite-bulk composition. A similar topic was raised by J.M. Dautria and M. Girod (France) whose study was devoted to the mantle xenoliths enclosed in basalts from the Tahalra massif (Hoggar); they concluded that the carriers of metasomatism were to be found in the magmatic veins of basanitic, kimberlitic and carbonatitic nature, occurring in the mantle peridotites. Carbonatite magmatism was discussed by B.T. Ngwenya (Malawi) who analysed a possible genesis of natrocarbonatites.

H. Bertrand (France) stressed the difficulties encountered when applying geochemical discriminant diagrams to the identification of the tectonic setting of the African continental tholeiites. He concluded that the observed dispersal of these diagrams reflects the diversity of the mantle source and that of the magmatic processes, rather than the tectonic setting.

Among the fundamental questions evoked were: the scale of the asthenospheric diapirism; the influence of old subducted oceanic lithosphere relocated beneath the African Rift System; the mechanism of rifting (primary or secondary role of asthenosphere?); the origin of silicic lavas (mantle,

Conference Reports

anatexis of crustal rocks or of underplated gabbros, fractional crystallization, involvement of CO_2 ; and the degree of crustal contamination. Others queried the existence of mantle plumes in different parts of Africa and the size and origin of diapiric mantle "mushrooms" under the stretched crust. Also unclear is the degree of interaction between lithosphere and magmas of asthenospheric origin, and the relation between the thermal state of mantle, stretching of the crust and magmatic production. General models accounting for a large spectrum of magmatic systems are now available, as opposed to the concept of uniqueness of each volcanic province.

Knowledge of the African extensional magmatism and of the rifting mechanisms has clearly been considerably enlarged during the last few years, thanks to the close cooperation among geoscientists from different disciplines. African rift

studies have now reached an exciting stage and additional progress will be made if the multidisciplinary approach is maintained and expanded, based on a well-established field geology.

The extended abstracts of 40 communications were published in a bilingual (French and English) volume of 250 pages by the CIFEG (International Center for Training and Exchanges in Geosciences). The next meeting of the IGCP 227 will be held in Washington DC, in July 1989 during the 28th International Geological Congress.

C. Coulon
Laboratoire de Pétrologie Magmatique
Faculté des Sciences de St-Jérôme
13397 Marseille Cedex 13
France

Tectonic Setting of Proterozoic Volcanism and Ore Deposits

Finland and Sweden, 15 - 22 August 1988

To bring together scientists actively working on the geochemistry, volcanology, metallogeny and tectonics of the Svecofennides, and similar problems elsewhere, IGCP Projects 217 (Proterozoic Geochemistry) and 247 (Precambrian Ore Deposits and Tectonics) held a joint meeting recently that began in Stockholm. Thirty-seven scientists from Sweden, Finland, Canada, The Netherlands, Zambia, Spain, Brazil, and the U.S.S.R. attended the field meeting, which consisted of a six-day field trip in Sweden (R. Gorbatschev - leader) and Finland (G. Gaal - leader); and a one-day symposium in Turku, Finland convened by C. Ehlers (Finland).

The first three days of the excursion were dedicated to the Bergslagen Region of Sweden, which is the name given to the ore-bearing segment of the Svecofennides in central Sweden. The geology of the region is complex, but basically comprises thick successions of variably altered and weakly metamorphosed, 1.9-1.86 Ga old volcanic and sedimentary rocks, which are cut by a variety of granitic plutons. Although the rocks were apparently erupted and deposited in a series of basins, the nature of the subsidence, the basement upon which the supracrustal rocks were deposited, and the regional tectonic setting are controversial. Nevertheless, the presence of granitic plutons and abundant siliceous volcanic rocks clearly indicate that there was pre-existing continental crust within the region: the fundamental questions concern its age and origin.

R. Hellingwerf (The Netherlands) showed us alteration, mineralization and breccias of the Alvilangan-Vikern stratiform Zn-Pb-Fe-Mn deposit and I. Lundström (Sweden) demonstrated the well-preserved primary volcanic textures of the region. Subsequent discussions focused on whether the volcanic rocks were subaerial or subaqueous, the origin of the widespread alteration of the region, and the nature and tectonic setting of the extension. A special issue of *Geologie en Mijnbouw* (v. 67, nos. 2-4) on the structure, stratigraphy and ore-forming processes of the Bergslagen Province, edited by J. Baker (The Netherlands) and R. Hellingwerf, was published in time for the excursion and provided useful background information between stops.

Near Hällefors where we focused on volcanic rocks and their alteration by high-level intrusions, J. Baker demonstrated variable Na metasomatism related to a hypabyssal, probably

sub-cauldron granitic complex, and I. Lundström once again demonstrated the excellent preservation of primary textures in the region by showing us a section containing cooling units of ash-flow tuff, air-fall deposits containing accretionary lapilli, and a variety of intercalated epiclastic rocks. B. Lagerblad (Sweden) then dealt with the Gyllingen massive sulphide deposit from which 10^5 tons of high-grade (10-14%) Zn-Pb ore were mined during the 1970s, and the historic Oster Silvberg mine, which was mined intermittently from as early as 1367 until the First World War.

A visit to metasedimentary and granitic rocks to the east and south between Uppsala and Stockholm led by G. Stalhös (Sweden), took us into a region where the metamorphic grade is generally upper amphibolite and the rocks penetratively strained, in marked contrast to those we had seen the previous two days. This part of the trip was particularly controversial and evoked tremendous discussion on the nature of volcanism and temporal relationships between plutonism, deformation, metamorphism, and their overall relationships to the lower grade and less strained rocks of western and central Bergslagen. For example, the preservation of low-grade volcanic and sedimentary rocks in the

Figure 1: P. Ward (Finland) leads a discussion on the outcrop.

Conference Reports

Bergslagen region clearly demonstrates that the crust was abnormally thickened, yet to the north and south much higher grade rocks are exposed at the surface, indicating considerable crustal thickening in those areas. To my knowledge no model adequately explains the spatial distribution of metamorphism and crustal thickening.

Following a one-day symposium in Turku, Finland, we moved to southern Finland, where the Svecofennides comprise E-W-trending belts of 1.895-1.887 Ga old volcanic and sedimentary rocks in which an interesting story of northward-directed folding and thrusting is emerging. The deformation appears to predate granitoid intrusions dated at 1.885 Ga. Microcline granites were emplaced in the region at about 1.8 Ga. The region was later refolded about north-easterly trending axes and then cut by the well-known rapakivi suite at about 1.7 Ga. The younger northeasterly trending cross-folds are rather broad, open structures which provide considerable structural relief through the earlier deformed region, and they await construction of down-plunge projections, as has recently been usefully applied in several Proterozoic orogenic belts in Canada.

Rocks on the islands of Nagu and Korpo, shown to us by C. Ehlers and A. Lindroos (Finland), were chosen to demonstrate the idea of northward-directed thrusting, and we saw

well-exposed and variably strained gabbros, amphibolites, and mafic volcanic rocks, in places structurally overlain by highly strained cordierite-garnet-biotite gneiss. Of particular interest were occurrences of highly strained rocks, probably mylonites, although this interpretation was hotly contested. M. Väisänen (Finland) clearly demonstrated the syntectonic nature of metamorphism in the Orijärvi-Aijala area, and he led the group to strained volcanic rocks and mylonitized limestone, which once again provoked considerable discussion of and the mechanisms for producing them.

Overall, the excursion served its purpose exceedingly well, bringing together a broad spectrum of geologists to examine regional problems in the Svecofennides. Not surprisingly more problems were generated than solved, but the positive atmosphere among workers in the region will undoubtedly yield exciting results in the years to come.

R.S. Hildebrand
Geological Survey of Canada
588 Booth Street
Ottawa, Ontario
Canada K1A 0E8

West African Orogens in a Circum-Atlantic Setting

Nouakchott, Mauritania, 8 - 20 December 1987

Late last year 71 intrepid geologists from many countries all around the Atlantic made their way to Mauritania, to participate in a rare experience, an IGCP Project 233 conference and field trips to a region not often visited by so many foreign geologists.

In his introductory conference talk on terrane analysis, the writer proposed a revised definition of a terrane applicable to circum-Atlantic Paleozoic orogens as "an area characterized by an internal continuity of geology that is bounded by faults, mélanges representing a trench complex, or cryptic suture zones across which neighbouring terranes (i) may have a distinct geological record not explicable by facies changes (i.e. exotic terranes), or (ii) may have a similar geological record (i.e. proximal terranes) that may only be distinguished by the presence of the terrane boundary representing telescoped oceanic lithosphere." Keppie also showed how the dating of accretionary fabrics can provide better constraints upon the initial docking of terranes, though it is necessary to distinguish between fabrics produced by subduction, docking and post-accretionary telescoping.

The geology of North Africa was then outlined by R. Black and G. Rocci (France) in terms of two Archean-early Proterozoic cratons surrounded by late Proterozoic orogens, which are amenable to terrane analysis and plate tectonic modelling. The West African craton has acted as a rigid core since 2 Ga ago and is surrounded by thrust-bordered late Precambrian Pan-African orogens: the Trans-Saharan Belt on the east and the Mauritanide Belt on the west, in places reactivated by Variscan and Alpine events.

R. Caby (France) showed that a west-to-east section across the Trans-Saharan Orogen in the Tuareg Shield (central Sahara) consists of the following tectonic elements: the West African craton, Pan-African foreland nappes derived

from a passive margin miogeocline, a dismembered ophiolite, a Pan-African magmatic arc complex, and reworked Archean-Proterozoic rocks. To the south, the Pan-African and ophiolite units are represented by a cryptic suture between the West African craton and the Benin-Nigerian craton, as summarized by M.A. Rahaman (Nigeria) and E.K. Agyei (Ghana). However, A.N. Boullier and J.M. Bertrand (France) showed that stretching directions in the Trans-Saharan Orogen are mainly subhorizontal N-S to NE-SW, suggesting that the accretion was oblique and probably sinistral. The orogen is also cut by numerous dextral N-S faults suggesting a complex kinematic history.

For Morocco, E. Wallbrecher (Austria) and Y. Bassias (France) documented a Late Pan-African, dextral ductile shear zone along the northwestern margin of the West

Figure 1: Field convoy approaching the foreland.

Conference Reports

African craton, also implying oblique accretion. Turbidites in the Volta Basin may be interpreted as foredeep deposits. Pan-African events ended with alkaline magmatism around 560 Ma ago. Geochronological data were presented by B.N. Ekwueme (Nigeria) and E.K. Agyei (Ghana), J. Manu (Ghana) and R.L. Armstrong (Canada). Ekwueme argued that the new Rb-Sr whole rock isochron ages of 1289 ± 153 Ma and 1313 ± 37 Ma provide evidence for the Kibaran Orogeny in West Africa. Pan-African ages are recorded by Rb-Sr and K-Ar isotopic ages between 667 ± 9 and 547 ± 2 Ma.

Figure 2: J.P. Lecorche explaining the geology of the Central Mauritania.

A. Pique (France) divided the Variscan terranes of Morocco into four units: the African craton (including the Reguibat Shield overlain by the Paleozoic Tindouf Basin, the Anti-Atlas, Coastal Block, North-central and North-eastern Morocco); the Western Sahara terrane consisting of nappes of probable Cambro-Ordovician sandstone and pelite (lower nappe), and schists and charnockites (upper), which were thrust over the Reguibat Shield after Frasnian time; the Sehoul terrane composed of thick Cambro-Ordovician turbidites and Late Ordovician deformation during which it was thrust upon the Meseta; and the Internal Rif, which is correlated with the Kabylia and Inner Betic all involved in Alpine orogenesis.

In contrast, J. Boulin, M. Bouabdelli and M. El Houicha (France) divided the African craton of Pique from northwest to southeast into a foreland in the Coastal Meseta, an external zone in the central Meseta made up of an early Paleozoic passive margin, an internal zone in the Midelt characterized by Devonian-Namurian, calc-alkaline magmatism and back-arc deposition in the Meseta basin, and allochthonous units resulting from collision between Coastal Meseta and the African craton during Late Carboniferous-Permian times.

As regards the Mauritanide Orogen, J.P. Lecorche (France), J. Roussel (France) and R.D. Dallmeyer (U.S.A.) proposed an eight-stage sequence as follows:

1. Mid-Late Proterozoic (c. 1100-700 Ma) deposition of West African Supergroup 1, continental sedimentary sequences on crystalline basement of the West African craton.
2. Late Proterozoic (c. 700-680 Ma) intracontinental rifting leading to the formation of oceanic lithosphere and just preceded (c. 720-700 Ma) by a tectonothermal event in the rift facies.

3. Late Proterozoic (c. 680-640 Ma) convergence, producing the western ensialic magmatic arc and imbrication of the oceanic and continental lithosphere during Pan-African I orogenesis.
4. Late Proterozoic-Early Cambrian (c. 640-550 Ma) uplift and cooling of the Pan-African I Orogen and deposition of basal tillite and flysch sequences in the foreland.
5. Cambrian (c. 550-525 Ma) Pan African II deformation and metamorphism ranging from intense in the Rockelides, decreasing in intensity northwards in the Bassarides and central Mauritanides and dying out in the northern Mauritanides. At the same time, molasse was deposited in the foreland.
6. Cambrian-Late Devonian (c. 525-360 Ma) deposition of clastic successions in the Taoudeni and Bove basins on the craton.
7. Early Carboniferous (c. 325-300 Ma) Variscan reactivation of the Mauritanide Orogen by folding and thrusting accompanied by variable grades of metamorphism. This was related to the collision of Gondwana and Laurentia which caused further convergence between the western continental block and the West African craton.
8. Early Permian (c. 300-275 Ma) local ductile deformation along faults bordering the western coastal block during the terminal stages of collision.

M. Kane (France) characterized the ultramafic-mafic complexes of the central Mauritanides as ophiolitic on the basis of their chemistry. Progressing still farther south, M. Villeneuve (France) reviewed the Pan-African Bassaride and Rockelide Orogens, which are similar to the Mauritanides except that the Pan-African orogenesis appears to have lasted here until c. 500 Ma, and Variscan deformation dies out southwards. Finally, R. Venkatakrishnan and S.J. Culver (U.S.A.) presented a model for the Mesozoic breakup of Africa and North America based upon tectonic fabrics in Sierra Leone.

The conference was followed by a field trip across the central Mauritanide Orogen led by J.P. Lecorche, R.D. Dallmeyer and O. Dia. This involved many 4-wheel drive vehicles with drivers, a water tanker, and food and cooks for eight days in the desert. The excursion began in the foreland to the east and progressed westwards across the para-autochthonous units, imbricated ophiolitic units, a continental margin rift facies and a poorly exposed calc-alkaline igneous complex in the west. The material assistance of the Mauritanian Ministère des Mines et de l'Industrie and the Direction des Mines et de la Géologie Mauritanie was greatly appreciated.

For those unlucky to have missed this memorable field trip, a vicarious experience may be acquired by reading the field guide - the first English summary of Mauritanide geology - available, as is the Abstract volume, from R.D. Dallmeyer, Department of Geology, University of Georgia, Athens, Georgia 30602, U.S.A.

J. Duncan Keppie
Department of Mines and Energy
P.O. Box 1087
Halifax, Nova Scotia B3J 2X1

The Cadomian Orogeny Reviewed

Oxford, U.K., 6 - 16 April 1988

The term *Cadomian*, taken from the Roman name *Cadomus* for the present city of Caën, is generally used to describe the late Precambrian magmatic, metamorphic and deformational events recorded by the Armorican rocks of north-western France and the Channel Islands. These events are broadly coeval with the late Precambrian evolution of the Avalonian area immediately to the northwest, and to events recorded by the Monian rocks of Wales and Ireland.

The Geological Society of London and IGCP Project 233 (Paleozoic Terranes in the Circum-Atlantic Area) recently held a conference to review these events. The meeting provide a forum for discussion between French and British geologists working on the Precambrian to Cambrian rocks of Cadomian affinity that crop out in France and Britain (especially the Channel Islands), and there were also contributions from workers on coeval rocks elsewhere in Europe and North America.

In a session on terrane analysis of southern Britain, W. Gibbons (U.K.) described a series of fault-bounded suspect terranes in northwestern Wales and southeastern Ireland, accreted by sinistral transposition around 540-550 Ma ago. The presence of steep basement lineaments in southern Britain, such as the Welsh Borderlands Fault System, may indicate transcurrent terrane boundaries in the Precambrian basement of the southeast. Thus the boundaries between Monian, Avalonian, and Cadomian areas may separate quite different displaced terranes, a view supported by J.C. Pauley (U.K.) in a contribution on Longmyndian rocks that emphasized the likelihood of Early Cambrian sinistral strike-slip faulting along the Welsh Borderlands.

T. Pharoah (U.K.) and colleagues suggested the proposed differentiation of three terranes on the basis of geochemical variations between late Precambrian inliers separated by major, long-active fault systems. Emphasis was also placed upon the presence of an earlier (c670 Ma) metamorphic-magmatic event in southern Britain, and a later phase of magmatism and low-grade metamorphism around the Precambrian-Cambrian boundary. The Precambrian rocks of the Rosslare area of southeast Ireland, as described by F.C. Murphy (Ireland), is notable for extensive exposures of steep, ductile, pre-Arenig mylonite belts that are likely to correlate broadly with those bounding Monian terranes in northwest Wales. The history of these shear zones is prolonged and complex, involving polyphase reworking of gneissic basement and a younger (?) cover sequence of low-grade metasediments.

The discussion then turned to France and a review by M.M. Shufflebotham (U.K.) of the Penthièvre region in NW Brittany, which is the "type-Pentevrian" as defined earlier by J. Cogné. Many lithologies are present here and interpretation of the relationships between them is often equivocal. A particularly controversial issue was whether or not an ancient pre-Brioverian "Pentevrian" crystalline basement existed at all in this area, the majority view being that it did not.

A refinement of Brioverian stratigraphy by D. Rabu (France) and others emphasized the existence of a major tectonic boundary between two non-coeval successions previously

interpreted as superimposed upon one another. E. Denis (France) showed variations in Brioverian sediments between northern and western Brittany, supporting the generally accepted idea that the northern rocks were closely related to a Cadomian island arc system active during the late Precambrian. The Brioverian in western Brittany must, he argued, have been deposited in much more passive conditions. A geodynamic model involving southeast-dipping subduction in late Precambrian time was presented by B. Cabanis (France).

Figure 1: A folded, syn-tectonic granite sheet at Pointe de la Garde, seen on the field trip to northwest France.

The continental red-bed sediments that form an overstep sequence resting unconformably upon Cadomian rocks in France and the Channel Islands were described by D. Went (U.K.), who contrasted these with true regional "molasse" deposits associated with major mountain chains. The term Cadomian Orogeny should not be misconstrued as suggesting some kind of Alpine or Himalayan collisional event.

The mid-Channel magnetic anomaly was interpreted by J.P. Lefort (France) as a mafic body at depth corresponding to a plate suture resulting from subduction under Brittany. Lefort interprets seismic reflectors in the area as thrusts transporting flakes of crust to the north, synthetic to southward subduction. Some higher level, northward dipping faults may be antithetic to Cadomian subduction whereas others are Hercynian in age.

There was much discussion of Cadomian magmatism in the Channel Islands. This is calc-alkaline in character, ranging in composition from the layered St. Peter Port Gabbro in Guernsey, through commonly hybridized dioritic and granodioritic intrusions, to true granites. As pointed out by M. Brown (U.K.) and colleagues, although Cadomian magmatism

Conference Reports

is extensive throughout the area, the continuity of crustal units across the North Armorican region has not been demonstrated and cannot be assumed: the original Cadomian subduction system may well have been dispersed by lateral fault movements. A particular problem lies in the fact that the tectonostratigraphy of this area was originally established in the Baie de St. Brieuc where there is now much doubt about the ages and relationships of key exposures.

Timing of Cadomian magmatic events remains a major problem, with ages varying from about 650 Ma to Silurian, clearly indicating a long and complex magmatic history, despite difficulties in interpreting radiometric data. Cadomian events were compared by M. Brown and his co-workers to the prolonged Mesozoic magmatism of the Chilean Andes, and a similar subduction model was developed by B. Auvray and P. Graviou (France), based on geochemistry of Cadomian granitoids. Some questioned whether the Mancellian granites in Normandy were truly 'S'-type crustal melts.

Figure 2: Examining syn-tectonic granite sheets in the St. Cast Shear Zone.

An important contribution to the tectonic evolution of the region came from G.J. Lee (U.K.) in his account of the Jersey Main Dyke Swarm. This is dominantly basaltic and of a mature calc-alkaline type and indicates a period of post-Cadomian crustal extension parallel to the proposed active continental margin.

A grand overview of Cadomian events was provided by J. Cogné (France), who emphasized that the Armorican area had grown by subduction-related crustal accretion in the late Precambrian (Cadomian), incorporating relict nuclei of ancient (c2000 Ma) crust within Gondwanaland. This process of accretion continued through the Phanerozoic with the Variscan and Alpine Orogenies.

A contribution dealing with supposedly volcanoclastic sediments in the Tregor and Petit Tregor areas of northern Brittany by N.H. Griffiths and R.A. Roach (U.K.) illustrated the problems of identifying certain shear zone lithologies, for some of these rocks may be mylonitic rather than tuffaceous. An interesting study of Nd isotopes by C. Guerrot

and J.J. Peucat (France) suggested that old lower crust influenced significantly the generation of Cadomian granites. Despite several uncertainties, especially concerning the interpretation of radiometric data, the geology of granitic rocks in the Cap de la Hague, according to G.M. Power and others (U.K.), bears many similarities to that of Guernsey, forming the northeastern end of a narrow, linear terrane characterized by the presence of ancient (c2000 Ma) "Icartian" basement.

The final day of the conference emphasized the tectonic assembly of the Cadomian Orogen. In a keynote address J.-P. Brun and P. Bale (France) presented a model that illustrated the composite nature of the area, its overall NE-SW structural grain, and the importance of sinistral shear zones, SW-verging thrusts, and gently NE-plunging stretching lineations. The importance of ductile shear zones within the Cadomian area was further emphasized by R.A. Strachan, G.M. Power and P.J. Treloar (U.K.), who made it clear that an understanding of these mylonite belts, several of which are likely to represent terrane boundaries, is crucial to further analysis of the area.

In northern Brittany these shear zones are sinistral. Some are younger than the St. Malo Migmatite belt and thus about 540 Ma in age. They may therefore be related to the same terrane accretion event recorded by the Menai Strait Fault System in northwest Wales. Available palaeomagnetic data on the Cadomian, reviewed by G.K. Taylor and J.D.A. Piper (U.K.), are as yet too inadequate to aid terrane analysis, but an important tectonic event at about 540 Ma appears to alter dramatically polar wandering curves across the area.

A major problem exists in attempting to correlate Cadomian terranes with those in Spain, the Bohemian Massif and the Appalachians of Canada, though R.D. Nance (U.S.A.) made a key observation in noting that the Cadomian of northwest France appears to have more in common with the geology of New Brunswick than with the "Avalonian" of southern Britain and Newfoundland. What is clear from these comparative studies is that late Precambrian events affecting the Gondwana area are recorded throughout Europe and eastern Canada and are presumably linked to Pan-African events farther south.

From this conference several points were of particular value. First, the majority of French and British workers have abandoned the term "Pentevrian." The c2000 Ma gneisses that run from Cap de la Hague through Guernsey (and probably Sark) to northwest Brittany can be referred to as Icartian, after the dated Icart Gneiss on Guernsey.

Second, the igneous rocks of this area are indisputably of calc-alkaline character and related to plate subduction, the polarity of which is unproven but generally believed to be towards the southeast. Third, the area is dissected by steep, commonly sinistral, high strain zones, which are likely to separate different terranes. The accretion of these terranes took place at least in part around 540 Ma ago. Future work is likely to concentrate upon the analysis of mylonitic rocks across the area. Finally, radiometric data are poor, and many more dates are needed, especially of mylonitic rocks within the shear zones. The conference proceedings are being published by the Geological Society of London.

W. Gibbons
Dept. of Geology
University of Wales
Cardiff CF1 3YE, U.K.

G.M. Power
Dept. of Geology
Portsmouth Polytechnic
Portsmouth PO1 3QL, U.K.