Global Catastrophes in Earth History

Snowbird, Utah, 20 - 23 October 1988

The geological record of mass extinctions and the causes and significance of such sudden disappearances were the focus of a recent interdisciplinary conference attended by 180 scientists. Sponsored by the Lunar and Planetary Institute and the National Academy of Science, this meeting provided an unusual opportunity for extended interdisciplinary discussion. Paleontologists were able to hear the varied views of specialists about shocked quartz, and planetary dynamicists could hear contrasting opinions of stepwise extinction.

Both this conference and its 1982 predecessor (GSA Special Paper 190) have their roots in the discovery of an anomalously high iridium concentration at the Cretaceous-Tertiary (K/T) boundary and its interpretation as the result of the impact of an extra-terrestrial body by Luis Alvarez and his colleagues from Berkeley. This second conference was organized because continuing research in diverse fields has heightened interest in possible causes, mechanisms and implications of global catastrophes in Earth history.

Both patterns and periodicity in extinction proved central issues. K. Flessa (all speakers from U.S.A. unless otherwise noted), in a keynote address, drew attention to the incompleteness of the fossil record and to under-representation in even the best collections. An example was P. Ward's presentation on his continuing study of ammonite extinctions at the K/T boundary on the north coast of Spain. Earlier collections indicated step-wise extinction, but five years of continued collection from numerous sections showed that disappearances are more likely to have been simultaneous. There remain differences of interpretation of the record. G. Keller reported step-wise disappearance of foraminifera from a Moroccan section across the K/T boundary, but J. Smit (The Netherlands) finds simultaneous extinction in the same section.

In a review of the Phanerozoic history of extinctions, J. Sepkoski discerned a period of 26 Ma for the last six major extinctions but found no regularity for the pre-Tithonian Phanerozoic. For D. Raup, catastrophic mass extinctions may play a larger role in evolution than the drum-beat of the Darwinian mode. This theme was expanded in a closing address by S.J. Gould who suggested that sudden, catastrophically induced mass extinctions may have a winnowing effect on life, with survival determined not so much by the accumulation of advantageous adaptations but by such relatively fortuitous circumstances as small size, protective habitat or by resistance to specific environmental effects of a catastrophe.

F.J. Swanson reviewed the complex and varied biological responses over the last eight years to the effects of the Mount St. Helens eruption. It was clear to the audience that modelling the response to a catastrophe is very difficult because it is not easy to predict before the event which features will prove dominant in the environmental response.

Because impacts have been implicated in mass extinction, there is much interest in their size and frequency. On the basis of observations with the 46 cm Schmidt telescope at Mount Palomar, G. Shoemaker and his colleagues, provided new estimates of the flux of Earth-crossing asteroids and comets. Impacts that yield craters 100 km or more across are thought to be capable of causing mass-extinctions. For objects this large, the estimated flux is about 2 in the last 100 Ma.

Dynamic effects of impacts were addressed using a variety of modelling methods: D. O'Keefe and T. Ahrens refined earlier models and considered CO₂ production from impact into a thick limestone terrain. The massive volume of CO₂ produced would result in global warming. A. Vickery and J. Melosh discussed the potentially lethal effects of partial loss of the atmosphere resulting from a major impact. P. Schultz and D. Gault showed that an oblique impact would produce a much smaller and less detectable crater than a vertical one, but a larger and more extensive vaporized mass. Several modelers, including D. Roddy, considered secondary volcanism at impact sites. On most interpretations, such volcanism would be significant only at the very largest craters. R. Prinn and B. Fegley expanded their model of the environmental effects of impacts to include a range of life-threatening events: inhibition of photosynthesis; foliage damage, toxicosis from soil trace metals and inhibition of marine calcification.

Figure 1: A quartz grain (0.5mm across) from the K/T boundary layer at Strakville South in the Raton Basin of New Mexico. The multiple sets of planar features with specific crystallographic orientations are equivalent to those observed at known impact sites. Photo courtesy of R. Grieve.

Multiple impacts of two kinds were considered. A perturbing mass passing close to the solar system could bring numerous comets from the Oort cloud into the inner solar system over an interval of 2 to 3 million years. Step-wise extinctions at various times (including the K/T and Eocene-Oligocene boundaries) and evidence of more than one impact event closely spaced stratigraphically such as that presented by J. Hazel have been considered to represent such a short-lived relatively intense impact flux. The suggestion that the K/T event might represent multiple simultaneous impacts, however arises from the singular nature of the iridium anomaly and the spatial variations in composition of the boundary clay constituents. A. Hildebrand and W. Boynton pointed out that a single impact in an area where continent-derived sediments overlay oceanic crust might do the trick. V. Sharpton and co-workers however, demonstrated

Conference Reports

the absence of ocean crustal mineral grains in the North American K/T sites, an absence that favours multiple impact sites instead.

The Manson structure in Iowa is of K/T age and may be the source of the shocked minerals that are particularly abundant along the eastern flank of the Rocky Mountains according to J. Hartung and associates. This structure, 40 km in diameter, however, is not large enough alone to account for all K/T effects. The Kara and Ust-Kara impacts in northern Russia were thought to be possible K/T events but new ^{40}Ar - ^{39}Ar dates presented by C. Koeberl (Austria) and co-workers from the U.S.A. and the Soviet Union have shown that the structures are actually approximately 75 Ma old.

The role of exceptionally intense volcanic activity in global catastrophes remains unresolved. J. Morgan showed the coincidence in time between the Deccan flood basalt eruption and the great K/T extinction. There is an intriguing possibility that the Siberian trap eruption may have been synchronous with the giant Permo-Triassic biological extinction, although the precise age of that eruption is not well established. Altogether six major extinction events over the last 250 Ma may be correlated with eruption of flood basalt. M. Rampino and R. Struthers also emphasized the possible coincidence in time of other flood basaltic eruptions, with significant marine extinctions during the last 250 Ma. They described a "grand universal hypothesis," somewhat like that of Brochwicz-Lewinski in which periodic large impacts at about 32 Ma intervals might have induced both flood volcanism and extinctions.

J. Sigurdsson, reviewing the volatile-rich eruptions of basaltic material from hot spots, pointed out that as in the 1783 Laki fissure eruption in Iceland, elevated CO_2 concentration lofted above the tropopause could result in a reduction of surface land-mass temperature of 1 to 3 degrees for less than a decade. Both Sigurdsson and T. Simkin, who considered the historic record of explosive volcanism, emphasized the potential of the as-yet-unrecognized, great volcanic event as at least a contributory cause in global catastrophes.

An expected consequence of the impact of a sizeable extra-terrestrial body in the oceans is a tsunami. An event layer that could be the result of a tsunami at the K/T boundary in Texas was discussed by J. Bourgeois. There, a section of mid-shelf muds, estimated to have been deposited at depths of 50 m or more, is interrupted by graded, coarse sand with large intraclasts that in turn is capped by laminated and rippled, fine sand. To produce this event layer, Bourgeois and her colleagues estimate that a wave some 50 m high is required. Modelling suggests that such a wave could be produced by a body 10 km in diameter affecting the deep ocean some 5000 km from the Texas site, or by a smaller body hitting closer.

Is the shocked quartz that is present in numerous sections of the K/T boundary the result of impact or could it also have come from volcanic eruptions? This crucial issue was the focus of intense discussion. B. Bohor, R. Grieve (Canada), V. Sharpton, and G. Izett who have studied the material from the K/T boundary as well as some of the specimens considered to be shocked quartz from volcanic deposits, insist that the volcanic examples lack the well-developed, multiple sets of planar deformation lamellae characteristic of impact-generated shocked quartz. Proponents of the volcanic origin, led by A. Huffman and C. Officer, countered by emphasizing the need for a thorough review of the criteria for identifying shocked quartz and underlined the uncertainties and incompleteness of study of volcanic deposits.

E. Anders and his colleagues drew attention to aspects of a postulated global firestorm at the K/T boundary. They had characterized in considerable geological and isotopic detail sections in New Zealand and Denmark that contained high concentrations of soot.

Could the K/T mass extinction be a model for other similar extinctions throughout the Phanerozoic? One answer has come from analyses of the Ir concentration found at various horizons of mass extinction wherein Ir spikes comparable to those from K/T boundary are sought. F. Asaro and associates reported an Ir anomaly from two widely separated ocean sites that both coincide in time with the mid-Miocene extinctions. This example is the third known from the Mesozoic/Cenozoic boundary; the others are at the K/T and the latest Eocene boundaries. On the other hand, C. Orth and M. Attrep, Jr. found only moderate Ir concentrations at several Paleozoic horizons of mass extinctions and at the Cenomanian-Turonian of the Western Interior Seaway, North America. They concluded that the present concentrations can be attributed to Earth surface processes including reactions with oceanic waters and spreading-center, gaseous magmas.

H. Geldsetzer and D. McLaren (Canada) described the knife-sharp disappearance of many taxa at the Frasnian/Famennian boundary, but no shocked quartz has been found and the only horizon with a moderate concentration of Ir in Western Australia is not at the boundary and could well be endogenous. The terminal Triassic extinction lacks both an Ir anomaly and shocked quartz according to A. Hallam (U.K.), and the same appears to apply to the end-Permian extinction event, the largest in the entire geological record. Taken together, the conclusion seems inescapable that the K/T event is unique in geological history.

Plans were discussed for joint field and laboratory examination of material from key sections of the much-studied K/T mass extinction. There is a need to find and study additional first class sections of this boundary and to explore for additional examples of presumed tsunami layers. Future studies of the boundary should also include increased attention to sedimentological aspects along the lines of the work by A. Preisinger (Austria) and his colleagues in Austria. Beyond the continued emphasis on the K/T events, there is a mandate for equal attention to other mass extinctions. And finally, there is a need to involve specialists in various branches of biology to explore the actual mechanisms of mass extinctions and their possible selectivity.

G. Shoemaker ended his review of astronomical observations of possible earth-crossing asteroids by showing photographs of an object for which the IAU had accepted the name "Alvarez." Shoemaker presented a framed copy of photographic prints of "Alvarez" to Walter Alvarez in paying tribute to his father's memory. He suggested that it is difficult to overemphasize the immensely stimulating influence of the "Alvarez hypothesis" for the earth sciences and for students of biological evolution.

R.N. Ginsburg
Rosenstiel School of Marine
and Atmospheric Science
University of Miami
Miami, FL 33149
U.S.A.

K. Burke and V. Sharpton
Lunar and Planetary Institute
3303 NASA Road 1
Houston, TX 77058
U.S.A.

Proterozoic Mobile Belts

Tianjin, China, 6 - 10 September 1988

Last September over 180 geoscientists representing 23 countries met in China, partly under the sponsorship of IGCP Project 217, Proterozoic Geochemistry, to discuss the geochemistry, tectonics and mineralization of Proterozoic mobile belts. Geological and geochemical evidence continues to support the existence of plate tectonics during the Proterozoic.

Many speakers used geochemical discriminant diagrams to constrain ancient tectonic settings. However, K.C. Condie (U.S.A.) and T.C. Pharaoh (U.K.) emphasized that numerous examples of modern basalts are classified in the wrong tectonic setting by such discriminant diagrams. Only when data from many different sources (such as lithological association, sedimentary environment and provenance, metamorphic and structural histories, mineral deposits and geochemistry, etc.) agree can we hope to identify ancient tectonic settings. However, C-Y. Chen (U.S.A.) concluded that relatively immobile trace element ratios can be used to identify original compositions and tectonic settings of spilites and keratophyres. Likewise, T.C. Pharaoh and Cheng Yanjing (all speakers from China unless otherwise indicated) confirmed the importance of bimodal volcanics in Proterozoic arc-like successions.

An in-depth summary of the geochemical and tectonic development of the Precambrian of the North China craton was given by Sun Dazhong. Because of Phanerozoic plate collisions, Proterozoic tectonic reconstructions in China are difficult and correlations must be done with great caution. However, many new Sm-Nd and U-Pb zircon ages (Li Shuguang, B. Hanson, F.R.G., Xia Yuliang) are beginning to provide a sound basis for unraveling the complex Archean and Proterozoic history of the North China craton. The widespread distribution of Proterozoic continental rift and craton successions discussed by Guo Hongxue, Yu Jianhua, Tang Min, Li Zenghui and others demonstrate that the craton was intact by the Early Proterozoic. In contrast, the Baltic Shield and North America were still undergoing arc

accretion in the Early Proterozoic. Some of the oldest known blueschists occur in central China. Dong Shengbao and Kang Weiguo showed that these rocks, which are at least 1 Ga in age, record a Late Proterozoic continent-continent collision along the southern margin of the North China craton.

R.T. Pidgeon (Australia) and O.M. Rosen (U.S.S.R.) presented summaries of U-Pb zircon chronologies in SW Australia and Siberia, respectively. It is clear that more than one zircon population can be found in both plutonic and volcanic rocks and that individual zircon or ion probe analyses are necessary to unravel complex geochronologic histories. Li Shuguang also presented convincing evidence that Sm-Nd whole rock isochrons can be reset during high-grade metamorphism.

Proterozoic anorthosites continue to attract interest. R.J. Wardle (Canada) presented data that suggest some anorthosites and associated granites may be syntectonic rather than anorogenic, and Hu Shiling showed that $^{40}\text{Ar}/^{39}\text{Ar}$ dating may be useful in dating anorthosite emplacement. A. Elueze (Nigeria) presented data that Proterozoic rifting may have been important in Nigeria, and R.A. Fuck and M. Figueiredo (Brazil) summarized geological and geochemical results that emphasize the importance of arc accretion in central and southern Brazil both during the Early and Late Proterozoic.

The probable presence of pre-Pan-African continental basement in Egypt was indicated by G.M. Salloum (Egypt). C. Picard (Canada) argued that the Ungava Trough in Canada may represent a small ocean basin. Norite dikes that are common in the Proterozoic are probably derived from enriched mantle sources and do not represent crustally contaminated komatiites, according to D.J. Hughes (U.K.). Much of the Limpopo Belt in South Africa that has previously been described as metasedimentary is probably of granitic origin (M. Boryta, U.S.A.). And Li Ziyang showed that heavy REE commonly follow U during U mineralization.

Several papers on Proterozoic mineral deposits described little-known Chinese occurrences. Wang Kuiyuan reviewed the Proterozoic Pb-Zn deposits hosted in carbonates and shales along the northern border of the North China craton. Luo Yaoxing presented data suggesting that the Proterozoic magnesite deposits in Dashiqiao may be evaporites that were later altered and metamorphosed.

Figure 1: IGCP 217 field trip participants examine pillow breccias of the Early Proterozoic Xiyanghe volcanics in the Zhongtiao Mountains of east-central China.

K.C. Condie
Department of Geoscience
New Mexico Institute of Mining
& Technology
Socorro, New Mexico 87801
U.S.A.

Sun Dazhong
Tianjin Institute of Geology
& Mineral Resources
Chinese Academy of
Geological Sciences
Tianjin, China

Applying Geoscience to Development

Nottingham, U.K., 26 - 30 September 1988

An international conference on the application of geology in developing countries was held recently at the University of Nottingham under the joint sponsorship of the Institution of Geologists, the Institution of Mining and Metallurgy, the Geological Society of London, and the Association of Geoscientists for International Development. Support from these organizations and the Overseas Development Administration, the British Council, the Commonwealth Foundation, and the Royal Society enabled a wide geographic participation - over 200 geoscientists from 38 countries.

The main aims of the conference were to assess past experience and discuss future trends in the application of the broad range of geosciences to developing countries. In his keynote address, Prof. P.G. Cooray (Sri Lanka) maintained that development means providing for the basic needs of people - food, water, shelter, clothing, energy, health, education, employment and a safe environment - thus improving the quality of life for all (see article by P.G. Cooray in *Episodes*, vol. 11, p. 123-129, 1988 - Ed.). Geoscientists have an essential and important role to play in this, and they were urged by Cooray to give their best.

R.L. Johnson (U.K.) opened a session on the nature and funding of geoscience aid programs with a paper on British technical cooperation overseas. Government bilateral aid had evolved from the former colonial geological survey days to encompass a range of service, technical, research and development and training schemes, in which individual specialists are seconded on short or long-term contracts to programs run and operated mainly by the host geological survey or other government department. G.O. Kesse (Ghana) discussed the nature and development of the successful Ghanaian-German cooperative project for gold prospecting in southwestern Ghana, clearly a useful model with much wider application.

A multilateral approach to geoscience aid was outlined by R.N. Aitken (Brussels), who referred to the "equal partnership" concept between Europe and the developing countries notably in the context of the forthcoming Lomé IV Convention negotiations, and by I.N. Nichol (Canada) who discussed the success of the U.N. Revolving Fund for Natural Resources Exploration in exploring for and identifying potentially economic mineral deposits in Latin America and Africa, and in locating a geothermal energy supply on the Caribbean island of St. Lucia (see Tomita and Nicol in *Episodes*, v. 11, p. 23-29, 1988 - Ed.). The concept of collaboration for mineral resource development within regional groupings of developing countries such as the South African Development Coordination Conference (SADCC) aroused much interest, as did one of the few papers devoted to the application of oceanography with regard to the 200 mile Exclusive Economic Zones, which cover some 25% of the Earth's surface but are poorly known in terms of their economic potential.

Considerable emphasis was placed on the topic of geoscience education and institution building. J.C. Chatupa (Malawi) spoke of the comprehensive geological data base that had been inherited by his country and of how the Malawi Geological Survey was managing the data under an appropriate geoscience development policy. C.A. Kogbe (Nigeria) referred to the current need of programs for the training of geoscientists in Africa, in contrast to those applicable in former colonial times, and P.E. Baker (U.K.) reviewed the need for and status of geoscience education

for Third World students in the United Kingdom, particularly in the light of the "rationalization" of earth science education currently underway in the U.K.

There was considerable support for the concept of institution twinning, and a general recognition of the importance of training technicians and of increasing the flow of scientific literature towards developing countries. Also stressed were the mutual benefits of participation by scientists from these countries in global geoscience programs such as the Ocean Drilling Programme. The basic concepts associated with the establishment of the International Training Centre in The Netherlands and, quite recently, that of the School of Mines Regional Centre for Asia and the Pacific (University of New South Wales, Australia) appeared to be particularly relevant, and examples of university education in Colombia, Jamaica, Tanzania and India provided insight into several new schemes and models. A special session was convened by C.A. Kogbe to reactivate the IUGS Commission on Geology Teaching and to draft a working list of objectives.

Growing demand for adequate supplies of water for drinking and irrigation purposes, coupled with the serious problems of rapid desertification in some parts of the world, has sharply focussed attention on the location and management of water resources. In many developing countries remote sensing, geographical information systems, cheap but effective well-drilling and groundwater modelling are all widely used. However, as V. Ricaldi (Bolivia) pointed out, the organizational infrastructure for water resource management is often the key obstacle to progress.

The need for effective communication and interaction between geotechnical engineers and geologists, construction engineers, planners and decision makers involved in urbanization, water control and transport sector development was stressed by Yudhbir (India). Considerable discussion was generated about the control of major floods, prediction of explosive volcanic eruptions and mega-earthquakes, and the nature of global sea-level rise and its effect on low-lying areas. An important contribution to the forthcoming UN Decade of Natural Hazard Mitigation will be the collation of data on and the publication of natural hazard maps for many developing countries. There was also a proposal that geological surveys place a major effort on urban geology, following the UN ESCAP lead for Asian cities.

The role of LDC geological surveys in the natural resource and other sectors was discussed by N. Money (Zambia). G.M. Clark (U.K.) stressed the importance of industrial mineral development in a country's overall economic development, taking into account the diverse and disordered nature of the marketplace. W. Chesworth (Canada) focussed attention on one relatively low-cost, widely applicable, industrial mineral resource - phosphate rock and mineral additives such as limestone, volcanic scoria, malachite and pyrite. Indeed, the interest in agrogeology reflects the renewed and growing attention being paid to in the potential use of locally available agrominerals.

New approaches to geological mapping in the developing countries were reviewed by E.R. Peters (U.K.), and other speakers discussed the increased use of remote sensing techniques, the integration by computer of exploration data, and electronic data processing. Emphasis should be placed increasingly on the compilation and management of comprehensive data banks rather than primarily on the publication of geological maps, a theme that was pursued also in the energy/metalliferous resources workshops, particularly in

Conference Reports

relation to exploration and geological work in remote terrains. R.P. Foster (U.K.) gave a stimulating report on research carried out in recent years on gold mineralization in Zimbabwe.

The importance of small-scale mining and mineral industries in rural development was emphasized by J. Spiropoulos (U.K.). He and others reported case studies including, for example, simple exploration techniques for gold, mineral concentration methods, gemstone mining in Zambia, and mining rights in the Philippines. There were reports also on the UN-sponsored seminar on small-scale mining held in Turkey just prior to the conference, the new AGID-IDRC initiative Small Mines International, and the AGID-ITDG project on the compilation of a directory of small-scale mining equipment.

There were, not surprisingly, numerous specific recommendations and suggestions made during the conference, some of which are alluded to above. To single out but a few, there were general proposals for:

- strengthening of databases within developing countries on all geological resources (metallic, energy, industrial, and water), on remote sensing, geophysical, and other data, and assistance in the field of geoscience information technology.
- active participation in global environmental and scientific programs, especially those concerned with sea-level rise and its effects on low-lying coastlines, and with such problems as natural hazards, deforestation and desertification.
- the establishment of linkages between institutions in developed and developing countries, as well as with East Bloc nations, and to foster also better East-West communication within the geosciences.
- major efforts to establish recognition among policy makers, economists and planners of the essential and growing value of the geosciences in development.

after report by D.A.V. Stow (Nottingham) and
A.J. Notholt (Uxbridge) U.K.

The Holocene in South America

Paraná, Argentina, 21–26 November 1988

The first scientific activity of the recently created INQUA Commission on the Quaternary of South America was an international symposium on the Holocene of South America co-organized with CADINQUA, the Argentine National Committee for INQUA. The scientific sessions were dominated by papers on Argentina, but interesting contributions on Uruguay, Paraguay, Chile and Brazil were also presented. Most came from the traditional universities and institutes of La Plata, but papers from emerging scientific centers, such as Mendoza, Rio Cuarto (Argentina) and Montevideo (Uruguay), were also conspicuous.

Papers on soils of Patagonia (M. Camilion, M. Osterrieth and J. Chionchi - all speakers from Argentina, unless otherwise noted) and Pampa (M. Osterrieth and A. Karlsson) pointed out the existence, in wide areas of Argentina, of buried and truncated soil profiles. For a climosequence in the Paraguayan Chaco, M. Hurtado and C. Camilion, concluded that the E-W variation in precipitation and evapotranspiration regimes produced deeper and finer soils in the eastern (humid) region than in the west.

Papers presented by J. Codignotto and colleagues, F. Isla and G. Martinez, N. Weiler, R. Violante and J. Clausen on littoral and marine processes, demonstrated different ages for the maximum Holocene transgression, and contrasting coastal and marine mechanisms for various sectors of the Argentine coast. Codignotto and co-workers showed that of the two largest barrier spits of Patagonia, the Valdes Bight spit (42°30'S), with a maximum age of 5700 years BP, is composite, whereas the San Sebastian Bay spit (53°S) is younger (around 4000 years old) and complex. In both cases, the growth of the spits advanced in distinct pulses.

North of Mar del Plata (38°S), a system of barriers with deposits of transgressive and regressive phases was described by R. Violante as comprising lagoons, tide plain, beach ridges, fluvial and aeolian sediments. The maximum level, attained 4600 years ago, was 4 m above present sea level. J. Cavallotto reported on a geomorphological study on the bed of the Rio de la Plata, in which he found two main environments, one in equilibrium with the present

dynamics, the other inherited from the Mid-Holocene transgression. On the other hand, in Bahia Anegada (40°S) N. Weiler has found evidence of a maximum sea level 6560 years ago 5 m above the present sea level. For the subsequent regression, three minor transgressive pulses have been registered, at 6190, 5200 and 4380 yBP. The ^{14}C ages were obtained from shells of *Tagelus plebeius* found in living positions.

From a study of 231 published dates on sea levels along the Argentine coast, Codignotto and colleagues concluded that in most cases the maximum Holocene transgression was attained between 6500 and 7300 years ago, with an ingressive peak at 9400 BP in the Rio Colorado area (39°30'S).

A more complicated history for the coast of Uruguay was described by D. Panrio and O. Gutierrez (Uruguay), with three coastlines at different levels. The highest, with a tentative age of 35,000 years, is characterized by a bed of molluscs. The second level is represented by the Villa Soriano and Vizcaino data at 5800 years BP. A third level, placed tentatively at 4000 years BP, is found in the Rio San

Figure 1: Excursion participants taking a break on the upper delta of the Paraná River. Photo by J. Rabassa.

Conference Reports

Salvador area, where it is represented by tide channels and pedogenetic concentrations. The whole pattern suggests a significant tectonic influence.

In an interesting session on paleoclimates, R. Villalba used dendrochronological data from Chile and Argentina to reconstruct latitudinal variations in the southeastern Pacific anticyclone for the last 500 years. The variations in thickness of annual rings in *Austrocedrus chilensis* and *Araucaria araucana* correlate strongly with the positions of the anticyclone. In the calculations, the monthly mean latitudinal position, the index of austral zonal circulation and the index of zonal central circulation of the anticyclone were considered. The results show large and irregular latitudinal variations of the system, over a total range of 15 degrees of latitude.

A study of paleowind directions in dune fields led M. Iriondo to propose the existence of an anticyclone on the Chaco-Pampa plain during the late Holocene. Paleoclimatic reconstructions for several areas of western Argentina were presented by D. Cobos, J. Boninsegna, R. Villalba and other dendrochronologists. They showed that in the province of Mendoza (32°S to 37°S) a continuous desertization of the climate over the past 400 years has been recorded, whereas in Rio Negro, Patagonia, the Frias glacier (41°10'S) registered its maximum Neoglacial expansion at 1638 AD.

Advances in understanding the local stratigraphy of several areas in the provinces of Cordoba and Buenos Aires were also reviewed. Four allostratigraphic units, described by M. Zarate and A. Blasi in the loessic La Postreta Formation of Late Pleistocene to Holocene age near Tandil (Buenos Aires) are each limited by erosion surfaces. The mobilization of the materials was produced by saltation and short-lived suspension related to local dust storms. M. Villegas and M. Sanchez reported their discovery of a system of aeolian and fluvial deposits in the middle and lower basin of the Cuarto River (Cordoba), which are Late Pleistocene to Holocene in age. The river belt comprises an old terrace, a modern terrace and the present flood plain; the surrounding plain is divided into an undulating aeolian unit, old dunes and modern dunes.

Fifteen contributions were presented on paleobotany and zoology. G. Cuadrado and J. Neiff described the palynology and composition of "embalsados" (rafts), a type of floating tropical peat, which today covers some 5000 km² of swamps in northern Argentina and large areas in Paraguay. Forty-four pollen taxa have been identified, most pertaining to species living at present in the same environments. S. Martinez (Uruguay) reported 27 species of molluscs in the Villa Soriano Formation near Merin Lagoon (eastern Uruguay) dominated by *Heleobia australis*, *H. charruana* and *Erodona mactroides*. He concluded that the temperature of the water was higher at the time of deposition, some 5800 years BP.

In terms of sedimentology, D. Poire and others described the significant deposition models for the Holocene of southern Cordoba and San Luis provinces in several papers. The general pattern suggests a very recent, and hitherto unsuspected, uplift (probably Late Pleistocene to Present in age) for the southernmost Sierras Pampeanas. In a provocative contribution, G. Irion (F.R.G.) suggested a damming-up of the Amazon waters during the last glacial maximum, resulting in the formation of a giant lake in the Amazon valley and the lower reaches of its tributaries. In an interesting study of the upper delta of the Paraná River, A. Sanchez D'Alotto demonstrated that the intensity of sedimentological modification was proportional to the age of the deposit. A comparison of the geomorphology before and after the largest historical flood of the river (1983) showed

Figure 2: The cliffs along the Paraná River. Photo by S. Sanguinetti.

that the modern fluvial units were significantly modified, the older ones slightly altered, and the middle Holocene littoral units of the area completely unchanged.

A new theory for the pre-Hispanic cultures of northeastern Argentina, presented by C. Ceruti, showed that the changes from dry to humid climate during the late Holocene was the main factor causing the dispersal and replacement of the dominant inhabitants at that time. With regard to land use, N. Abelli and co-workers described a system of management and conservation of soils now successfully applied in a wide area near Rosario (Pampa). The procedure is based on the preservation of the very friable upper thapto horizon (an aeolian silt) by means of limited plowing and other conservative practices. The addition of chemicals has been shown to be unnecessary.

Participants in the conference field trips visited the principal Holocene depositional environments of northeastern Argentina. On the excursion to Toma Vieja, a classic locality visited by D'Orbigny, Darwin and other famous naturalists, participants examined the imposing eastern cliff of the Paraná flood plain, which at 1200 km long and 30-50 m high is probably the largest single Holocene feature of the continent. Mass movements (especially rotational slumps) and large scale gullying are actively modifying the slope. The gullies, in turn, are colonized by tropical forest-type vegetation because of the special microclimatic conditions developed in these places. As a general consequence, the forest has advanced in this way some 600 km to higher latitudes.

Martin H. Iriondo
CONICET
Casilla de Correo 487
3100 Paraná
Argentina

Variscan Terranes in Western Europe

Montpellier, France, 26 - 31 August 1988

The latest meeting of IGCP Project 233 (Terranes in Circum-Atlantic Paleozoic Orogens) brought some 100 people from 14 countries together recently in France. An exhibit of a Map of pre-Mesozoic terranes in Circum-Atlantic Phanerozoic orogens compiled by many of the participants provided an opportunity for the writer to present an introductory overview of some of the results of the project.

Archean and Early Proterozoic rocks have now been recognized in all the major circum-Atlantic cratonic terranes, and Middle Proterozoic (Grenvillian) rocks are now known in all except the West African craton. Pan-African orogens are developed across all of these older basements but are limited to central and western Europe, the southern British Isles, the southeastern Appalachians, northwest Africa and northern South America. Accretionary deformation associated with the Pan-African orogens overlaps the birth and spreading of Iapetus' suggesting that these parts of the present circum-Atlantic lay in different oceans in Precambrian-Cambrian times.

The abundance in the Pan-African orogens of magmatic arc and peri-arc terranes, as compared with the circum-Atlantic Paleozoic orogens may be related to the absence of continent-continent collision in some of the former. Late Precambrian-early Paleozoic disrupted terranes composed of tectonic mélanges occur along the margins of many of the major cratonic terranes and represent imbrication of the continent-ocean transition. Most ophiolites, magmatic arcs and peri-arc basins in the vestiges of Iapetus are Late Cambrian to Early Ordovician, except in central and western Europe where they range from Silurian to Devonian. The ophiolites generally appear to represent small ocean basins.

The main accretionary events appear to be early Paleozoic in the Appalachians-Caledonides, persisting into the late Paleozoic in the Variscan Orogen. Late Paleozoic events in the Appalachians-Caledonides may be related to progressive transpression while convergence was still taking place in Europe. Faunal provinciality evident in the Cambro-Ordovician and cosmopolitan in the Siluro-Devonian supports closure of Iapetus by the end of the Ordovician. The Project 233 terrane map will be the base for other thematic maps and provides one tool in the search for a palinspastic model for these orogens. Current models based purely on paleomagnetic data provide widely divergent views and need to be integrated with terrane analysis.

J.F. Slack (U.S.A.) and H.S. Swinden (Canada), pointed out that whereas metallogenic data can be used to aid terrane identification, complications arise when mineralization is repeated at different times within one terrane, when similar mineralization occurs in different terranes, when similar but tectonically unrelated deposits are structurally juxtaposed, and when post-accretionary mineralization superimposes epigenetic ores on previously formed deposits.

Swinden, Slack and A.P. LeHuray (U.S.A.) showed how lead isotopic compositions from volcanogenic massive sulphide deposits provide a fingerprint of the host terrane. Deposits in most of the Avalon composite terrane plot in a restricted field close to crustal growth curves - an exception is from southeast Newfoundland. Deposits in the northern and central Exploits Subzone of Newfoundland have moderately

radiogenic lead, whereas those in the southern Exploits Subzone are consistently higher in radiogenic lead, and those of the Notre Dame Subzone have relatively non-radiogenic lead. The data support some correlations along the Appalachians, while some plot in diffuse fields overlapping those defined in Newfoundland. However, caution was urged because basement-derived lead isotopic compositions may be contaminated by more radiogenic lead in sedimentary depositional environments.

Figure 1: Terranes for sale. Anchor is presumably for docking the terranes. IGCP 233 participants on excursion in France.

R.A. Ayuso (U.S.A.) and M.L. Bevier (Canada) showed that Pb isotope variations in northern Appalachian granites define four regionally distinct groups of plutons indicative of different source regions. For the northern Appalachians, Keppie demonstrated close correlation between mineral deposit type and tectonic environment prior to accretion, but this was replaced by widespread, superimposed, poly-metallic mineralization related to Devonian (-Carboniferous) "stitching" plutonism following accretion.

On the application of geophysics to terrane analysis, H. Perroud (France) reviewed Variscan paleomagnetism, concluding that reliable data for the Paleozoic is far from complete and that limitations of the method may not allow distinctions between neighbouring terranes to be made. Other speakers used gravity, magnetic, and seismic data to trace terrane boundaries beneath overstepping sequences both on- and off-shore, the most comprehensive of which by J.P. Lefort (France) is to be published shortly in book form.

Much work by French geologists was reviewed in a session on terranes in the Variscan belt of France. Only in the northern part of the Armorican Massif are the Cadomian (late Precambrian) terranes preserved without the obscuring effects of the Variscan orogenesis. Here, from northwest to southeast are a late Precambrian magmatic arc built upon a 2.0 Ga old basement bordered by a back-arc basin and all thrust southwards over a continental margin sequence, which is alongside a continental block with a turbidite basin on its southern side. The southern part of the Armorican Massif and all of the Massif Central were involved in strong Variscan orogenesis, involving mainly east to west nappe transport and cut by large, generally dextral transcurrent shear zones and faults.

At least six major nappes have been recognized in the Massif Central (P. Ledru and A. Autran) that may correlate with nappes in the southern Armorican Massif (M. Colchen and others, P.L. Guillot and J.P. Floc'h, France; P. Matte). From bottom to top these consist of an early Paleozoic

Conference Reports

volcanic arc, an ophiolite, late Precambrian-(?)Paleozoic miogeocline, late Precambrian-early Paleozoic continental rise, an early Paleozoic peri-arc basin, and Carboniferous rift volcanic and sedimentary rocks. Dismembered ophiolites occur at the base of some nappes (G. Dubuisson and others).

D.S. Santallier and co-workers showed that the metamorphism associated with the Variscan progressed from high pressure during the Ordovician and Silurian, through medium pressure in the Devonian-Carboniferous, to low pressure in Late Carboniferous time. The high pressure metamorphism is attributed to subduction processes, and the latter two are accompanied by widespread plutonism.

A selection of papers from the conference is to be published in *Tectonophysics*, and others will appear in book form. It is a tribute to the linguistic capabilities of French speaking geologists that the papers at the conference were presented in English. This added greatly to the effective dissemination of information on all regions of the circum-Atlantic.

J.D. Keppie
Department of Mines and Energy
P.O. Box 1087
Halifax, Nova Scotia
Canada B3J 2X1

Metalliferous Black Shales

Beijing, China, 31 July - 4 August 1988

The second international meeting of IGCP Project 254 on Metalliferous Black Shales took place last year in China as a part of a major symposium on "Sedimentology Related to Mineral Deposits." The 370 attending scientists (280 of them from China) were presented with forty scientific papers concerning metallogeny of black shales.

Project 254, organized only two years ago, has already produced new information on the genesis of ore deposits connected with black shales. For example, in a survey of more than 4500 chemical analyses of black shales from all known Soviet gold deposits, the geochemical characteristics of potential gold-bearing black shale can now be identified. Based on the chemical composition, including major and micro-elements, it is now possible to formulate a hypothesis about the type of gold ore to be expected in quartz veins, in sheet-like zones of veinlets, associated with black shale formations, or in disseminated sulphides in fine-grained black shales. In general, the typical characteristics of gold-bearing black shale are increased FeO content, and increased values of $\text{FeO}/\text{Fe}_2\text{O}_3$, C_{org} and S. Furthermore, increased content of Na and lower content of Ca and Ti can be traced. This geochemical model, which is very important in prospecting for gold ores associated with black shales, is being verified in other regions as well.

Copper-ores in black shales of the "Kupferschiefer" type constitute one of the most important sources of Cu in the world. Significant progress has been made especially on understanding the genesis of these ores, which have long been classified as a typical example of syndimentary ore formation. Accumulating clay-carbonate muds, rich in organic matter, generated hydrogen sulphide and produced a reducing environment. However, the amount of metals concentrated during deposition seems to be responsible solely for the geochemical anomaly typical of almost all types of black shales. Most of the sulphide mineralization was formed during diagenesis, from early burial up to post-lithification, and is not directly controlled by sedimentation. The fact that Kupferschiefer ores were formed during multi-stage processes will also influence new prospecting techniques for these important ore deposits.

New data were also presented on the diverse roles of organic matter in the genesis of U-V, polymetallic, Mn and Hg deposits. A lithological control apparently exists in U-bearing black shales, and the metal enrichment is usually syngenetic to early diagenetic in origin.

IGCP 254 has now launched a group to correlate chemical analyses. This has distributed samples of the USGS Devonian oil shale from Kentucky (standard SDO-1), which is now the subject of the international interlaboratory comparative studies. The characteristics of SDO-1 will be published through 1989 in the *Geostandard Newsletters*, and a USGS Bulletin on SDO-1 characterization will be prepared.

During the meeting, national coordinators met with the management of Project 254. The main goals of the project were discussed - to characterize the carbonaceous formations in various regions, and the processes leading to the formation of the economically significant accumulations, especially of Au, Cu and accompanying elements, U-Mo-V, base metals, PGE, and manganese ores. Four international workshops are planned for 1989: Leoben, Austria; Krakow, Poland; Washington D.C.; and Sofia, Bulgaria.

Jan Pasava
Zbynek Gabriel
Geological Survey
Malostranské nám. 19
118 21 Praha 1
Czechoslovakia

Figure 1: At the Tangganshan manganese ore deposit.

Early Organic Evolution and Mineral and Energy Resources

Maria Laach, F.R.G., 19 - 23 September 1988

The final meeting of IGCP Project 157, which had focussed on the influence of the biosphere on processes of economic mineral and hydrocarbon formation, was convened recently as the 9th Alfred Wegener Conference, with support from the Alfred Wegener Foundation, the Deutsche Forschungsgemeinschaft and the Max-Planck-Society.

As pointed out by M. Schidlowski, Conference Chairman and Project Leader, Project 157 was driven by the fundamental premise that life is a powerful geochemical agent responsible for quantitatively important transformations in the exogenic cycle. For example, one of the sub-project topics, fossil soils or "paleosols" may be aptly described as reaction rims at the contact between the Earth's crust and the ancient atmosphere that had fixed various liquid and gaseous reactants during their development.

Reporting on Lower Proterozoic paleosols of the Elliot Lake region, C.E.G. Farrow and D.J. Mossman (Canada) emphasized their polygenetic development under relatively oxygen-deficient conditions and their significant role, directly and indirectly, in processes of ore formation. T. Reimer (F.R.G.) described carbonaceous high alumina shales in the Transvaal Supergroup, resembling Precambrian paleosols, but which developed as a result of intraformational karst-type weathering over a vast area. From his work in coastal Venezuela on a recent ferri-ferrous oolite, M.M. Kimberley (U.S.A.) hypothesized that soil development has been irrelevant to iron formation, a view that generated considerable controversy. Among other presentations, two papers dealt with experimental work designed to simulate weathering in possible Archean atmospheres. A key participant at all levels, G.J. Retallack (U.S.A.) startled some with his tales of Martian soils held in deep freeze for the past 2000 Ma and of carbonaceous chondrites as the oldest abiotic paleosols in the solar system.

The role played by life forms is in many instances more obvious, especially as regards the organic constituents of Precambrian sediments, the topic of a second sub-project. An example is the 3.8 Ga isotopic record of autotrophic carbon fixation from carbon in sedimentary rocks, presented by M. Schidlowski (F.R.G.). Yet within this record of isotopically light sediments, H. Strauss and co-authors (F.R.G.) reported major secular variations of $\delta^{13}\text{C}$ in late Proterozoic organic carbon correlative with variations in carbonate carbon. They conclude that the record encodes valuable information concerning the onset and evolution of biochemical pathways. Several other workers, notably A.J. Kaufman and colleagues (U.S.A.) described secular variations of carbon isotopic ratios in carbonates and organic carbon-rich rocks. Consistent enrichment of ^{18}O in the micritic phase vs whole rock carbonate, led them to conclude that the former more accurately reflects primary isotopic values.

Several contributions dealt with determination of thermal maturity of organic matter and with state-of-the-art procedures in organic geochemical analysis of kerogens and oils of various ages. Emphasis was on the molecular chemistry of early organic matter. On the early Earth such reactions may have been significant. B. Nagy (U.S.A.) and D.J. Mossman dealt with the paragenesis of kerogens in Lower Proterozoic metasediments of the Elliot Lake region,

Canada, beginning with cyanobacterial mats, from which extracted hydrocarbons underwent limited migration and subsequent resolidification as a result of various hydrous-hydrothermal episodes.

On a related theme, the importance of biomarkers as potential environmental indicators was recorded by several groups, notably F. Pu (China) and co-workers, who described the extensive distribution of late Precambrian cyanobacterial dolostones in north and southwest China, and by D.M. McKirdy and R.J. Willink (Australia) who used isotopic and biomarker geochemistry to identify numerous controls on the distribution and composition of organic matter preserved in the pre-Devonian sedimentary record. Particularly instructive was a summary by T.G. Powell and R.E. Summons (Australia) of the geochemistry of Precambrian source rocks and oils using Australian and Siberian examples. They report surprisingly different geochemical signatures, with a marine carbonate source rock implicated for the Siberian oils, in contrast to mixed lacustrine-marine characteristics for the McArthur basin of Australia.

As regards the antiquity of bacterial sulphate reduction, another sub-project, H. Ohmoto (U.S.A.) focussed attention on the sulphur isotopic composition of Archean marine sediments and on the geochemistry of Archean oceans. He presented a model in which sulphate-rich seawater and intense activity of sulphate-reducing bacteria help to account for decreased sulphate-sulphide isotopic fractionation in Archean marine sediments compared to those of younger age. However, the presence of large quantities of pyrite and hematite in banded iron formations evidently requires that the Archean oceans contained appreciable amounts of free oxygen. Reviewing the case for bacterial sulphate reduction, P.A. Trudinger (Australia), maintained that the evidence all indicates that sulphate reduction evolved in the early Precambrian, before the development of an oxic atmosphere.

In an effort to draw the "bottom line" regarding the antiquity of dissimilatory sulphate reduction, M. Schidlowski noted that the problem remains to explain the disappearance of bacteriogenic sulphide patterns in the sulphur isotopic record from ca. 2.8 Ga, a phenomenon clearly incompatible with the fossil record of prokaryote differentiation and the existence, as early as at least 3.5 Ga, of a marine sulphate reservoir. An explanation may lie in the peculiarities of Archean seawater geochemistry and (by implication) mantle-driven processes.

I.B. Lambert and colleagues (Australia) used examples of Lower Proterozoic sulphide deposits to demonstrate how easily the role of biological sulphate reduction and hydrothermal processes can overprint each other, resulting in a great range of $\delta^{34}\text{S}$ values. Curiously, $\delta^{34}\text{S}$ values obtained from Proterozoic sediments on a world-wide scale show a positive $\delta^{34}\text{S}$ imbalance, interpreted by Lambert as evidence for a Proterozoic supercontinent.

The evolution of microbial processes in the sulphur cycle through time was discussed by H.G. Trüper (F.R.G.). The phylogenetic picture is currently being focussed using the technique of 16S ribosomal RNA sequencing, which permits recognition of subtle relationships hitherto unsuspected

Conference Reports

between prokaryotes. Trüper reports the property of dissimilatory sulphate reduction occurring in archaebacteria and also in several species of early eubacteria.

The fourth sub-project topic, fossil microbial eco-systems of the "stromatolitic" type and their recent analogs drew much attention at the conference. W.E. Krumbein and associates (F.R.G.) presented an overview of geophysiology - the science of the phenomena and normal functions of Earth as a living material body. There followed several interesting "case histories" of some early Earth biosphere activities including: Lower Precambrian stromatolite associations of Karelia by V.V. Makarihin (U.S.S.R.) and a review by J. Bertrand-Safarti and R. Milandou (France) of giant subtidal stromatolites in the upper Proterozoic of the Congo. The latter appear to form part of a complete facies sequence from basin to carbonate ramp, lagoon and sabkha, not unlike that of the present day Persian Gulf. Another highlight was J. Paul's (F.R.G.) presentation on stromatolites, bioherms and biostromes of the Central European Zechstein basin, itself the subject of a popular 3-day field trip led by the speaker after the close of the conference.

Several papers dealt with the description and interpretation of modern stromatolites as possible models for ancient analogues. J. Trichet and colleagues (France) described a modern type of calcareous stromatolite from the Tuamotu Archipelago in which calcification occurs by *in situ* precipitation, under biological influence, of high-magnesium calcite. The Australian team of R.V. Burne and L.S. Moore presented fascinating details of modern microbial communities from various localities world-wide, emphasizing their potential environmental facies interpretation.

S.M. Awramik (U.S.A.) made a good case for treating stromatolites as trace fossils, because in most instances fossils of the various organisms normally found in a living cyanobacterial mat are not present. However, in those rare exceptions that disprove rules, stromatolites provide key evidence for 3500 Ma - old life on Earth. According to S.

Golubic and A.H. Knoll (U.S.A.), cyanobacteria are one of the morphologically most complex and diverse groups of prokaryotes. Most astonishingly, judging from comparisons between modern and ancient cyanobacteria, prokaryotes diversified rapidly and at high taxonomic levels on the early Earth. In this sense, modern cyanobacteria contain the largest number of "living fossils" known.

Results of experimental work with stromatolites and stromatolite environments were reported by two main groups of researchers. Y. Cohen (Israel) found that primary carbon production and sulphate reduction is most efficient at salinity ranges up to twice the concentration of seawater. This results in rapid recycling of virtually all of the photo-assimilated CO₂ by SO₄ reduction. B. Lazar and J. Erez (Israel) reported that the extreme carbon isotope depletions found in hypersaline microbial mats results from intense photosynthetic activity in the mats, which produce a positive CO₂ gradient across the atmosphere - brine interface. M. Schidlowski, H. Gorzawski and I. Dor (F.R.G. and Israel) account for the exceptional enrichment of ¹³C in the biomass of stromatolite-type microbial ecosystems from experimental solar ponds as a response to high productivity sustained under conditions of eutrophication. The trend to use heavy carbon stems, in effect, from largely indiscriminate metabolism of all available CO₂-supplies.

In sum, the meeting was an occasion to draw up a balance sheet of a decade's work pursued under the auspices of IGCP Project 157 in those blurred areas where biogeochemistry, economic geology and evolutionary biology tend to blend or overlap. Judging from the conference proceedings (to be published by Springer-Verlag), it may be safe to say that the field as a whole owes much to the stimulus and coordinating efforts that emanated from Project 157.

D.J. Mossman
Mt. Allison University
Sackville, N.B.
Canada E0A 3C0

M. Schidlowski
Max-Planck-Institut
Postfach 3060, D-6500 Mainz
Federal Republic of Germany

Protecting Ancient Works

Athens, Greece, 19 - 23 September 1988

If you were planning the ideal geological conference, you would probably want to have the opening ceremonies at some classic location associated with the theme of the conference, followed by a site visit to a type area, followed by a plenary session devoted to what the attendees just saw in the field. That is exactly how Professor Paul Marinos and his organizing committee scheduled the first day of the recent International Symposium on Engineering Geology and Protection of Ancient Works, Monuments and Historic Sites.

The opening ceremonies were held in the 2,150 year old Odeon of Herod Atticus (Fig. 1), at the base of the Acropolis of Athens. These were followed by an engineering geological tour of the Acropolis and its classical monuments: the Parthenon, the Erechtheion and the Propylaea. The afternoon was devoted to a plenary session in which geological aspects of the physical and chemical deterioration of the Acropolis and its monuments, including methods for structural restoration and protection, were presented.

During the sessions that followed there were many papers on the effects of slope instability and on foundation problems. Case histories indicate that a worldwide method of remedying foundation problems associated with ancient building

Figure 1: The Odeon of Herod Atticus - site for the opening ceremonies.

Conference Reports

seems to be jet grouting. Other presentations dealt with the geology of ancient construction materials, and the quarries from which these materials were taken. A topic of wide interest continues to be the range of physical and chemical processes leading to the deterioration of building stones and monuments, including bacterial alteration of limestone. Many different types of synthetic resin, applied and/or injected, are now being used to inhibit and stop the deterioration processes.

Many sessions at the conference dealt with the application of geophysical techniques to archeological exploration, with earthquake hazards (the downfall of many ancient cities and indeed civilizations), and with early construction techniques and materials. Even paleogeography was a theme as speakers reviewed the gradual, and sometimes not so gradual, changes in the environment, such as sea-level fluctuations that had profound effects upon ancient cities and past civilizations.

Figure 2: The results of slope movement at a monastery near Mount Athos.

The post-conference field trips attracted a great deal of attention. One through the Peloponnese, in southern Greece, began with a boat tour through the historic Corinth Canal to examine bedrock faulting, jointing and earthquake related rockfall damage. A highlight of this trip was the visit to the Temple of Apollo Epikourios at Bassae (5th century B.C.), which is suffering from severe differential settlement and physical and chemical weathering of its limestone columns - the oldest known examples of Corinthian style.

Figure 3: The ruins at Dion at the base of Mount Olympus, where a rising groundwater table is threatening the archeological site.

During other excursions, there were many stops of both historical and geological interest including Olympia, the ancient birthplace of the Olympics, which had its low moments in history, due to natural hazards. A severe earthquake, which destroyed the city, was followed by major flooding in the 6th century AD that buried, but preserved, the ruins in thick deposits of alluvial mud. Near Patras, participants viewed the suspected location of Eliki, an ancient Ionian city believed to have been swept into the sea by an earthquake-triggered landslide. Another highlight was a visit to the monastic state of Mount Athos, where the dramatic effects of slope instability on these 800 and 900 year old structures could be seen (Fig. 2). A visit to the ancient city of Dion at the base of Mount Olympus (Fig. 3) provided an opportunity for a discussion on methods of lowering the pervasive regional groundwater table that has risen several metres since the classical Greek period and is now threatening the Temple of Isis.

The conference was a large one with an interdisciplinary participation of 520 from 39 countries. These included engineering geologists and members of the International Association of Engineering Geology the sponsoring society, and soil and rock mechanics, engineers, geologists, geophysicists, geochemists, archeologists and architects representing the International Society of Soil Mechanics and Foundation Engineering, the International Rock Mechanics Society, and the International Council on Monuments and Sites. The proceedings of the symposium have already been published in three volumes by A.A. Balkema.

after report by D.F. VanDine
Victoria, B.C., Canada

Marine Geology in Asia

Shanghai, China, 7 - 10 September 1988

Recent years have seen a rapid growth in hydrocarbon exploration and geological investigations off the Asian coasts, resulting in the accumulation of vast amounts of valuable data. These were reviewed recently at the First International Conference on Asian Marine Geology held on the campus of Tongji University in Shanghai, and attended by 130 scientists from 15 countries. The broad spectrum of topics included the tectonics and evolution of the western Pacific and its marginal seas, deep-water changes and paleogeography, dynamic sedimentation on the continental shelf, deep-water and near-shore mineral resources and marine micropaleontology. Some years ago "marine geology" for most Asian countries meant nothing more than "coastal geology," but this conference was dominated by papers on deep-water and off-shore geology. Research on Asian marine geology has clearly spread from surface sediments to deep structures, from near-shore to deep sea, from pure description to a consideration of dynamic processes and causal mechanisms.

As stated by IUGS Past-President E. Seibold (F.R.G.) in his key-note lecture at the opening session, marine geology can be defined as "geology between land and sea," and continental margins are one of the bridges in between. This is particularly true for the Asian off-shore areas where the active continental margins of the western Pacific with their series of arcs, trenches and associated marginal seas directly bridge the continent and the ocean. Of great interest were papers describing the results of detailed surveys of trenches around Japan; these significantly improve our understanding of the geodynamic phenomena related to the plate subduction. For example, A. Taira (Japan) described a variety of lithologies in the basement of forearc regions, including accretional wedges, old continental basement and ophiolitic rock assemblages.

Many papers were contributed to tectonic evolution of marginal seas and arc systems. V.P. Zinkevich (U.S.S.R.) stressed the role of accretion of the continental margins in the tectonic evolution of the Bering Sea Basin, and several taphrogenic stages were distinguished by H. Gribidenko (U.S.S.R.) in his model of tectonic evolution proposed for the Okhotsk Sea plate. An example from the Japan Sea presented by H. Tokuyama (Japan) showed that crustal thickening occurred at the final stage of back-arc spreading and was caused by huge intrusions of igneous rocks into the oceanic crust. Xue Weiling (China) demonstrated and interpreted a geophysical profile of 2,160 km in length, running from east of Taihang Mountain and the North China Plains, through the Bohai Gulf, Yellow Sea and East China Sea to the Ryukyu Trench. Emphasizing the significance of microplates, H. Kagami (Japan) discussed lateral movement of the forearc sliver of the Nakai Trough during the past 500,000 years and its tectonic consequences. A series of arcuate structures ("epi-continental arcs") along the East Asian coastline behind the island arcs was pointed out by Xue Jiawei (China).

Asia, the largest and highest continent in the world, is also the largest contributor of terrigenous detritus to the ocean system. Accordingly, the dynamics of sedimentation on delta and shelf was one of the major topics of the conference, and the role of such factors as tidal currents, storms and shelf sea circulation were evaluated and reconsidered. Zhu Erqin (China), for example, holds that the surface sediments of the East China Sea shelf are in

equilibrium with recent hydrodynamic conditions, and that the widespread sands there are of autochthonous origin, rather than relict sediments of late Pleistocene age.

Figure 1: Examining Ordovician turbidites in Zhejiang province, near the western border of the East China Sea.

Attention was also drawn to mud patches surrounded by sands in the outer shelf of the East China Sea and ascribed to cyclonic gyres there. Duan Weimin (China) discussed characteristic features of early diagenesis of fine-grained sediments in these patches. Suk Bong Chool (South Korea) discussed the deposition history on the East China Sea shelf and its adjacent seas during sea-level fluctuations since the late Pleistocene. Distribution patterns of fine-grained sediments in the Yellow Sea were shown by Yong Ahn Park (South Korea) to indicate that the mud in its central portion is mainly supplied by the Yellow River, whereas that in the southeastern Yellow Sea is supplied mainly by other sources, such as the Korean rivers.

On the topic of deltas A.S.R. Swamy (India) showed that the rate of sedimentation varies from 0.1 to 0.9 cm/year on the east coast of India, and he identified five sets of Holocene sand ridges in the Krishna delta, extending 14 km inland and 27 km offshore from the present coast. Discussing the dynamic sedimentation in the Gulf of Thailand, P. Menasveta (Thailand) reviewed its oceanographic conditions as related to sedimentation, and A. Siripong (Thailand) reviewed the surface distribution of suspended sediment in the Phang-nga Bay, together with its variations in time and space, using remote sensing and water sampling.

On the topic of sea-level change and coastline migration, there was debate about the existence of "Holocene high beach-rocks" and the frequency and nature of Quaternary sea transgressions in East China. Shao Shixiong (China), for instance, listed eight transgressions in the east China Coastal Plains since the Gauss/Matuyama boundary and attributed all of them to eustatic changes. R. Shaw (Hong Kong) related the deposition and erosion history of the sea floor around Hong Kong to the Holocene transgression on the basis of offshore geological mapping.

Conference Reports

Among the scientific highlights were papers on paleoceanography of the South China and adjacent seas. For example, J. Kennett (U.S.A.) stated that the Kuroshio Current intensified as the result of effective closure of the Indonesian seaway by the late Miocene. On the basis of foraminifera, their transfer function paleotemperature, and oxygen isotopes, for example, marine micropaleontologists from Tongji University showed that during the last glaciation drastic changes in configuration of the South China Sea and its circulation pattern caused a significant decrease in winter sea surface temperature (about 9°C lower than during the interglacial), a remarkable increase of fine-grained terrigenous detritus on its northern slope, and a reduction in dissolved oxygen in deeper waters. T. Oba (Japan) proposed a new method for estimating temperature, oxygen isotope composition and hydrology of the paleocean from isotopic measurements of individual foraminifera and proved its usefulness by application to piston cores taken from the northwest Pacific.

There were also several papers on the sedimentary facies and geological history of coral reefs in the South China Sea and along the Indonesian coasts. As revealed by bore-holes drilled on the Xisha Islands, reefs began to grow in the early Miocene and continue to the present day.

The conference demonstrated the remarkable degree of international collaboration, for example, on cruises organized jointly by Japan and France, China and the U.S.A., and China and West Germany. The participation by Prof. Chen Peiyuan from Taiwan in the conference was the beginning of direct exchanges and cooperation between marine geologists from both sides of the Taiwan Strait, and it was decided to prepare a list of Asian marine geologists and to promote academic exchanges within Asia.

The next conference will be held in Tokyo in 1992. A 207-page volume of extended abstracts from the conference can be purchased for US\$20 from Lao Quiyuan, at the address below.

Wang Pinxian
Dept. of Marine Geology
Tongji University, Shanghai 200092, China

Fluid Systems and Fluid-Rock Interactions

Paris, France, 29 August - 2 September 1988

The past decade has seen many improvements in characterizing and analyzing the chemical and physical properties of modern and fossil fluids, and their interactions with minerals and rocks. These were reviewed at a recent symposium at the University of Paris co-sponsored by the IUGS Commission on Igneous and Metamorphic Petrogenesis. Papers presented at this symposium were of two types: those concerning natural systems and those dealing with experimental work.

A interesting review of the behaviour of REE, with respect to geothermal waters, was presented by A. Michard (France), who also introduced many new data. In general, the REE content of hydrothermal solutions is dominantly controlled by pH rather than temperature, but partitioning of REE between minerals and fluids is still poorly known. From the Lardarello (Italy) geothermal field, E. Petrucci, B. Turi and S.M.F. Sheppard (Italy) suggest, from H and O isotope studies, that two rock units exchanged isotopes with a fluid of probable meteoritic origin.

A model, presented by A.B. Thompson and J.A.D. Conolly (Switzerland), involved dehydration and hydration reactions as the primary control of fluid fluxes and crustal porosity. For large-scale models, the generation of metamorphic fluids is mainly controlled by metamorphic heat sources.

For fluid inclusions rich in $\text{N}_2\text{CO}_2\text{-CH}_4$ from the Montredon thermal dome, the N_2 was shown by C. Guillot and colleagues (France) to have been derived from an organic primary source. N_2 was then trapped as NH_4^+ in micas, and possibly later recycled. Nd and Sr tracers were used to show that at least two different fluids from different sources were involved in this system, according to R. Giere and colleagues (Switzerland). And for the Himalaya, C. France-Lanord and S.M.F. Sheppard (France) showed that whole-rock D values increase with metamorphic grade, though the calculated isotopic compositions of the fluids are rather uniform over the profiles.

Most interest was directed towards experimental studies. These were introduced by E.U. Franck (F.R.G.), who reviewed the experimental determination of binary systems involving water and other volatiles (CO_2 , CH_4 . . .). He also discussed the modelling of these systems by the means of an equation of state based upon a well defined molecular interaction model, and showed that small amounts of electrolytes strongly shift the range of two-phase behaviour to higher temperatures.

Another invited lecture was given by T.M. Seward (New Zealand) on the speciation and solute association in hydrothermal fluids. Recent experiments demonstrate the continuum from "individual" ionic species at room temperature to appreciable ion pairing at 300°C. New data on transition metal chloride complexes (i.e. NiCl^+ , FeCl^+ . . .) indicate stabilities similar to the HCl pair. A model based on the dielectric constant of the solvent was proposed by J. Schott and others (France), which applies to $\text{H}_2\text{O-CO}_2$, $\text{H}_2\text{O-NaCl}$, etc., and works for neutral and charged species. However, the authors' conclusion seems too optimistic, at least for systems involving H_2O , volatiles, and salts.

The knowledge of the speciation of solutes in hydrothermal fluids is of primary interest for fluid-rock interactions. In this connection the importance of using both electronic optical absorption and Raman spectroscopies to characterize the solute species was clearly demonstrated by D. Crerar and colleagues (U.S.A.).

On a different topic, P. Landais and others (France) reported their experimental work on the maturation processes of organic matter, comparing the obtained results with natural systems. The best simulation is obtained using pyrolysis in gold tubes under 1 to 4 kbar pressure.

Zhao Bin and colleagues (China) presented an impressive list of partition coefficients between granitic melts and aqueous vapour for elements such as W, Mo, Ta, Nb, K, Ca, Mg, Fe, F and Cl. Using a radiotracer technique, F. Delbove and others (France) reported the partitioning of Cl between

Conference Reports

alumino-silicate melts and aqueous fluid. The relative contents of Na, K and Ca have a strong effect on the variation of the partition coefficient with pressure.

Feldspar-fluid interactions are still fashionable, as indicated by the number of papers dealing with that topic. Data from literature suggest that the rate of dissolution of minerals in aqueous fluids depends on the density of dislocations. Nevertheless, new experimental data presented by W.M. Murphy (U.S.A.) on the solubility of sanidine (at 95°C, pH3) show no significant difference in the rates of reaction between undeformed and deformed samples. Dissolution is stoichiometric with rate constant at 10^{-13.6} mol. cm⁻² sec⁻¹. Experimental results on anorthite and plagioclase, presented by S.C. Dujon and colleagues (France), show that the physical properties of synthetic anorthite depend on the run conditions. In the plagioclase-water-chlorine system, fluid unmixing occurs up to high pressures, yielding complicated metasomatic evolutions.

The partition of coefficient europium between anorthite and Cl-bearing fluid was shown by P. d'Arco and others (France) to be independent of the Cl content. It decreases from 0.8 to 0.1 with increasing Eu in anorthite. However, the location of the ions is not clear, and defining a thermodynamic partition coefficient does not seem appropriate at this stage.

It is extremely difficult - and often impossible - to estimate the content (or activity) of common solute species, such as fluorine or boron, and particularly H⁺ (or HCl), in paleo-hydrothermal systems. M. Volfinger and M.L. Pascal (France) used specific mineral associations to buffer the

activity of HCl in fluids. The calibration has been experimentally determined and can be used in natural systems as an "HCl probe." V. Manier-Glavinaz and others (France) used the same technique of buffering solute concentrations in solutions by the means of convenient mineral or mineral associations to estimate the composition of aqueous fluids that equilibrated with vein and pegmatite minerals. These include Li, Na and Cs in solution at equilibrium with alkali-rich beryls, Vg and H in solutions at equilibrium with mafic minerals (talc, chrysotile, forsterite, brucite), Ca and Mn in fluids coexisting with Ca and Mn tungstates, Fl and aqueous phase buffered by phlogopite or topaz + andalusite, and B in solutions buffered by tourmaline + cordierite or chlorite + albite + andalusite + quartz.

The effect of water in normally anhydrous minerals has dramatic effects on the physical properties of these minerals, particularly their rheology. J.C. Doukhan and others (France), in their discussion of the incorporation of water into quartz and diopside, reported that in the former it occurs as clusters of water molecules, or as a point defect solution, whereas in diopside, most is associated with minute clino-amphibole lamellae.

Abstracts have been published in *Chemical Geology* and selected papers will be published in the *European Journal of Mineralogy* in late 1989 and early 1990.

A. Weisbrod
Ecole Nat. Sup. de Geologie
B.P. 452
54001 Nancy cedex
France

M. Lagache
Ecole Normale Supérieure
24, Rue Lhomond
75231 Paris cedex 05
France

CANADIAN SOCIETY OF PETROLEUM GEOLOGISTS CALGARY, CANADA

Memoir 14

Devonian of the World

Edited by N.J. McMillan, A.F. Embry and D.J. Glass

Devonian of the World is the proceedings of the Second International Symposium on the Devonian System, which was hosted by the Canadian Society of Petroleum Geologists in Calgary, Canada in August 1987. It contains 2222 pages, bound in three approximately equal-sized volumes.

Volume I: Regional Syntheses

Global Devonian Geology
Canada, U.S.A.
Europe and North Africa
South America and Southern Africa
Australia, New Zealand and Antarctica

Volume II: Sedimentation

Clastics and Tectonics
Black Shales
Carbonates, Reefs and
Evaporites

Volume III: Paleontology

Paleoecology and
Biostratigraphy
Global Events and Boundaries
Paleoecology and Biostratigraphy

Cost in Canadian funds,
including postage and handling:
Canada \$100
International \$115

Send order to:
Canadian Society of Petroleum Geologists
#505, 206 - 7th Avenue S.W.
Calgary, Alberta, Canada T2P 0W7

Please send remittance in Canadian funds with order,
or supply Visa number and expiry date