Computers and Resource Exploration

Helsinki, Finland, 21 - 23 July 1988

Modern methods of resource prediction and resource assessment in exploration were the topics of a symposium cosponsored last year by COGEODATA and the International Association for Mathematical Geology (IAMG), and convened by the writers. Thirty-eight papers were presented to some 75 scientists from 16 countries along with several practical computer demonstrations on such subjects as trend analysis, classification, correlation, expert systems, image analysis, remote sensing, geostatistics, computer mapping, and databases.

Most of the computing reported was done on microcomputers, which now are widespread in academic, governmental, and industrial organizations. Although no new techniques were revealed, the papers demonstrated that predictive techniques and those that optimize are important. It was also shown that modeling is popular and effective and that, above all, the systems approach and data integration are necessary in today's environment.

In their paper on statistical pattern integration, F.P. Agterberg, G.F. Bonham-Carter and D.F. Wright (Canada) offered new and original solutions to three important problems: how to optimize binary pattern for linear features association with mineral deposits, integration of linear structures with geochemical and lithological features, and quantifying uncertainty due to missing information. D.P. Harris and Goucheng Pan (U.S.A.) abandoned the concept of gridding spatial data and introduced a new intrinsic sampling methodology that allows the use of genetic relations to integrate geodata and the optimal discretization of various geofields and deposit-model areas. The gold potential of the Bohemian Massif was evaluated by J. Janatka and P. Moravek (Czechoslovakia) using a combination of empirical approach and multivariate analysis, notably cluster analysis and multiple linear regression.

C. Kliem and Th. Petropoulos (F.R.G.) integrated the exploration knowledge in the assessment of the mineral resources of parts of Greece and Turkey by various statistical methods, and H. Kürzl (Austria) reported on the use of a digital Geographic Integration System (GIS) for investigating regional patterns used to calculate the favourability of occurrence of known mineralization types. A short description of SCANDING, developing out of the REGION package, a fully integrated resource-assessment software used in Eastern European countries was given by E.A. Nemirovsky (U.S.S.R.), who also reviewed, with V.V. Marchenko (U.S.S.R.), the principles underlying the use of geological maps in quantified resource assessment.

The interface between bibliographic and factual data was the topic of a talk by E. Sauzay, H. Teil, M. Vannier and L. Zanone (France) who presented the GEONIX package for data integration by various statistical techniques. Characteristic analysis was applied by S.B. Suslick and B.R. Figueredo (Brazil) to regional geochemical data in part of Brazil for indicating favourable areas for base metals and bauxite. V.G. Zolotarev (U.S.S.R.) formulated an approach to the genesis of granitoid-associated hydrothermal ore deposits for estimating precious and base metal potential.

In a pragmatic approach, J. Aarnisalo (Finland) reviewed the use of image processing techniques by Outokumpu Oy (Finland) in integrating satellite geophysical and, to some extent, geological data for pinpointing exploration targets. A geographic information system (MARICA) developed by BRGM was demonstrated by D. Bonnefoy and A.L. Guillen (France). This is an integrated and interactive package that

unites image processing techniques, statistical treatment and expert system techniques. P. Leymarie, J. Dardel and L. Renouard (France) introduced new techniques of computing digital elevation models from SPOT stereo pairs used for structural analysis in uranium exploration.

Figure 1: Computer specialists at Helsinki.

Conditional simulation techniques in correlating geological structures with known oil occurrences were discussed by H. Burger, M. Eder, A. Mannaa and W. Skala (F.R.G.). With the SEAPUP computer program, D.J. Forman and A.L. Hinde (Australia) simulated drilling and discovery of onshore petroleum traps in Australia and assessed undiscovered crude oil resources, and S.M. Habesch (U.K.) used image analysis techniques to measure the geometrical parameters of porosity networks in thin sections of oil reservoir rocks. Mathematical modelling of sedimentary structures and subsidence history was applied to multivariate data in a microcomputer to predict hydrocarbon resources in part of East Germany by J. Harff, J. Springer, B. Lewerenz and W. Eisenbeck (G.D.R.). Probable sizes of undiscovered petroleum resources were estimated by T. Jasko (U.K.) using the Monte Carlo and Latin Squares methods on a microcomputer.

After compiling an inventory of mineral resources and lithological variations of Brazil and China, J.C. Griffiths (U.S.A.), H. Hu and H.C. Chou (China) compared these two countries on the basis of geological diversity and drew conclusions on mineral resource development of China and Brazil. D.N. Pilant, J.C. Griffiths and M. Smith (U.S.A.) applied the GLOBAL data bank for a similar geological comparison of the U.S.S.R. and the U.S.A., using Atkin's Q-analysis, and B. Saltikoff and T. Tarvainen (Finland) demonstrated the use of explorational data bases in the Geological Survey of Finland.

As regards related statistical techniques, F. Chung (Canada) presented a new regression technique based upon the maximum likelihood method for utilization of geochemical datasets with observations below detection limits, and J.E. Robinson (U.S.A.) pointed out the pitfalls of trend analysis techniques as applied in microcomputer softwares and demonstrated simple tests to avoid them.

The contributions of the conference presented a representative cross section of current applications of computerized mathematical techniques in resource exploration worldwide. The papers presented increased our knowledge in this field of practical mathematical application and will, we hope, facilitate world-wide contacts among experts and users.

Gabor Gaal
Geological Survey of Finland
SF-02150 ESPOO
Finland

Dan F. Merriam
Department of Geology
Wichita State University
Box 153, Wichita
Kansas 67208, U.S.A.

Tectonics of Eastern Asia and the Western Pacific

Tokyo, 13 - 16 December 1988

The sixth in a series of Japan-U.S.S.R. symposia on geotectonics, held recently in Japan, was the occasion for an international forum on new ideas and results from studies of tectonic problems in the Western Pacific and eastern Asia sector. Organized by the Tokyo Institute of Technology and the Japanese National Committee for the International Lithosphere Program (DELP), and sponsored by the Inter-Union Commission on the Lithosphere and the Ministry of Education, Science and Culture of Japan, the meeting was attended by 49 foreign and 101 Japanese scientists.

In a review of the progress and achievements of the Japanese DELP research program, S. Uyeda (Japan) expressed his view that although some of the achievements were spectacular, the overall goal of the program was not reached in part because it was too ambitious, and in part because funds and supporting facilities were insufficient. However, the DELP program undoubtedly served as a common forum for geologists and geophysicists alike and fostered an atmosphere of interdisciplinary cooperation.

In a second invited talk, P. Tapponnier (France) used a vast amount of field evidence to argue that a substantial amount (about 30%) of crustal shortening caused by the collision of Tibet to Eurasia was accommodated by the extrusion of several blocks in southeast Asia along the major strike-slip faults, such as the Altyn Tagh and the Red River, in central and eastern Asia. Photographs taken by the SPOT satellite were most spectacular in uncovering Pleistocene or Holocene displacements along these major faults. Holocene slip rates along the Altyn Tagh and Karakorum faults are between 20 mm/year and 45 mm/year. Tapponnier concluded that the crustal blocks in Tibet and southeast China are presently escaping rapidly eastward.

C. Sengör (Turkey) compared examples of tectonic style from the Cimmerian and Alpine systems of the Mediterranean and Asia to those from the Circum-Pacific and older parts of the Cordilleran orogenic belt of North America. He concluded that many complexities of orogenic belts result from post-collisional disruption of tectonic elements. Sengör warned that the increasingly popular "terrane analysis" which regards most of these disrupted fragments as independent micro-continental objects has led many researchers astray in the interpretation of orogenic collage. He suggested that it is necessary in studying these to distinguish first "primary components" (originally buoyant objects that became accreted after collisions) from "secondary components" (slivers produced by the disruption of already accreted materials or from host continents). This can only be done if the genetic interpretation of terrains (and not terranes) are taken into account.

Other highlights of the meeting included reviews of the opening process in the Japan Sea, Okinawa Trough, and Izu-Bonin Trough; tectonic processes in the Philippines, Taiwan, and Indonesia; new paleomagnetic data from various parts of China (e.g. Tarim, North China, South China, Tibet); and results from detailed surveys in the Shikoku Basin and the Western Pacific.

Two interesting papers were given based on petrological studies, which may have important bearings on the formation of marginal basins. Y. Tatsumi (Japan) argued that, when the Japan Sea opened at about 15 Ma, the opening must have been associated with the injection of "hot

asthenospheric material" chemically similar to mid-ocean ridge basalts (MORB). His data show that chemical composition of the volcanic rocks changed systematically to approach that of MORB, suggesting a mixing of the wedge mantle material with "injected material."

S. Maruyama (Japan) discussed the chemical composition of back-arc basin basalts from the Philippine Sea, Lau Basin, and Scotia Sea. His compilation suggests that these basalts were formed with a much smaller degree of partial melting than in Pacific and Atlantic MORBs, and that physico-chemical conditions for the formation of back-arc basalts are similar to those of Atlantic MORB. Maruyama concluded that back-arc opening is similar to the process occurring in large oceans such as the Atlantic, due perhaps to an uprising asthenospheric diapir, rather than to a passive process such as trench retreat or anchored slab model.

Figure 1: Amused by continental tectonics, T.W.C. Hilde (U.S.A.) on left, Shi Yanshen (China) next to him. B.C. Burchfiel and J. Dewey on the right.

Many of the speakers were aware of the limits of the simple plate tectonic regime in deciphering continental tectonics. In contrast to Tapponnier's views, J. Dewey (U.K.) concluded that crustal thickening in Tibet was the main agent that took up the shortening due to the continental collision of India with Eurasia. In any case there was unanimous consent that the large amount of shortening cannot be accounted for by the classic schemes involving rigid plates.

Plate tectonics also fails to give adequate explanations for the formation of marginal basins, and growth of continental margins and island arcs. It is apparent that "accreted" material plays an essential role in the evolution of both continental margins and island arcs, and that extrusion tectonics and accretion are necessary to account for the complexities of continental tectonics. In this respect B.C. Burchfiel (U.S.A.) suggested that detachment of the upper crust from the lower crust and uppermost mantle is plausible in view of the rheological characteristics of continental lithosphere. This might be one of the reasons why a vast amount of continental area can move along major strike-slip faults.

The proceedings of this symposium will be published in a special issue of *Tectonophysics*.

M. Kono
Dept. of Applied Physics
Tokyo Institute of Technology
Ookayama 2-12-12
Meguro-ku, Tokyo 15
Japan

B.C. Burchfiel
Dept. of Earth Sciences
MIT
Cambridge, MA 02139
U.S.A.

Metamorphism and Crustal Evolution

Changchun, P.R. China, 20 - 23 September 1988

More than 100 geologists from nine countries met recently at the Changchun College of Geology to discuss aspects of metamorphic geology, and especially to evaluate the results derived from metamorphic maps of continental scale.

The participants exchanged experiences concerning the compilation of metamorphic maps at small scales, and the scientific results derived from this international project. Most attention was paid to metamorphic type and its relationship to tectonic setting and crustal evolution. Based on evidence from the Metamorphic Map of China, Dong Shenbao, Lu Liangzhao and others (China) advanced a four-fold genetic sub-division of metamorphic type and emphasized their regular variation in geological age with three megacycles of crustal evolution in China. In contrast, N.L. Dobretsov (U.S.S.R.) argued that the development of specific metamorphic types since Proterozoic times was in accordance with the evolutionary stages of plate-tectonic cycles.

The concept of fault (dynamothermal) metamorphism was presented by Yang Chaoqun (China), who highlighted deep faulting as the main geological process involved in this metamorphic type. And rift metamorphism was also suggested as a new type characteristic of non-orogenic regions.

In his keynote address, J. Zwart (The Netherlands) pointed out that metamorphic maps of first generation had been nearly completed for most continents and countries. The urgent task was now to make detailed studies of representative metamorphic terrains of various type. From this would come metamorphic maps of second generation on the basis of new data and new conceptions.

There was much discussion of the early Precambrian metamorphic geology of the North China and Siberia platforms, as well as of southern India. Speakers generally affirmed that TTG series intrusives predominate over supracrustal rocks in most high-grade terrains, and that the protolith of many layered amphibolitic rocks and basic granulites was mafic or ultramafic intrusives rather than volcanics. Poly-phase metamorphism and ductile deformation of different crustal level are, of course, common in these terranes.

More and more blueschist belts of post-Late Proterozoic age have been found in recent years, including over 20 in China. The characteristics and genesis of those in Central China, Taiwan, the Urals and the eastern U.S.S.R. were reported by Zhang Shuye (China), J.G. Liou (U.S.A.) and N.L. Dobretsov (U.S.S.R.) respectively. Vigorous discussions took place on the sub-division of metamorphic facies and their tectonic settings. Dong Shenbao and co-workers divided blueschists into glaucophane-lawsonite facies, formed in typical subduction zones, and glaucophane greenschist facies, related to the closing of deeply rifted troughs within sialic crust. However, N.L. Dobretsov argued that the wide variations in pressure conditions shown by blueschist mineral assemblages depend only on the maximum depth reached by the subducting plate. Other researchers thought that there may be blueschists of different genesis.

The complex evolution of metamorphic events and tectonics in many orogenic belts, such as Sierra Nevada, California (H. Day, U.S.A.), the Scandinavian Caledonides (A. Stromberg, Sweden) and Qinlin of Central China (You Zhendong, China) was also a topic for discussion. The indications are that most orogenic belts have undergone more than one metamorphic event and orogenic cycle, and there were always several recrystallization and deformation episodes in a single event. In many cases, neighbouring litho-tectonic units of obviously different protolithic formation and metamorphic type may be allochthonous tectonic sheets of different source.

From studies of the PTt path of metamorphic events in the Altay Hercynides, Zhuang Yuxun (China) concluded that progressive metamorphism here was characterized by increase of temperature with decreasing pressure during continuous uplift of the crust. He challenged the use of the geothermal gradient concept in dealing with metamorphism of this kind.

Lu Liangzhao and Yan Hongquan
Institute of Geology
Changchun College of Geology
Geological Palace, Changchun
Jilin, P.R. China

Rudistologists Convene

Belgrade, Yugoslavia, 22 - 29 October 1988

Extinct Jurassic and Cretaceous attached shells, called rudists, have long been known in literature. For example, they were mentioned under the name "Orthocerati" by A. Fortis in the famous book *Viaggio in Dalmatia*, published in 1774. Their systematic study, however, began with Ph.P. Lapeirous (1781) and J.B. Lamarek (1801). Although examined by many renowned paleontologists, there still are many problems associated with rudists to be solved; problems that require closer and more active international collaboration.

The idea of an international conference of "rudistologists" was frequently mentioned by late B. Milovanović (1908-1977) of Belgrade University, one of the most distinguished connoisseurs of rudists. The eightieth anniversary of his birth was the occasion last year in Belgrade for just such a meeting, sponsored by the Union of Geological Societies of Yugoslavia, the Serbian Academy of Sciences and Arts, and the International Paleontological Association. Thirty-eight scientists from 12 countries attended.

An association of middle Cretaceous rudists from central Mexico was described by G. Alencaster (Mexico). Genera were distinguished (*Kinobleia*, *Caprinuloidea*, *Texicaprina*, *Radiolites*, *Toucasia* etc.) that clearly formed a reef core community, with others (*Requienia*) from the lagoonal facies.

Rich rudist biostromes with well-developed specimens were found to have been formed from the Cenomanian to the Senonian on the northeast margin of the Dinaric carbonate platform (western Slovenia), but not farther southward. The northeast part of the platform was wholly destroyed in late Senonian time and redeposited in the form of olistostromes in the adjacent Slovenian basin (S. Buser, D. Pejovic and R. Radoičić, Yugoslavia).

A succession of rudist communities was described from central Italy. *Hippurites* and corals from the open shelf prevailed from the Santonian to the Campanian. They were replaced during the late Campanian and the early Maastriichtian by the forms such as *Bournonia* and *Gorjanovičia* from a shelf of reduced hydrodynamic energy because of its

Conference Reports

Figure 1: On a Rudist reef at Leposavić, southern Serbia. L-R P.W. Skelton, G. Alencaster, M. Sladić-Trifunović, A. Polsak and J.M. Pons.

complicated morphology. Finally, radiolitids from the tidal environment of medium-to-low hydrodynamic energy prevailed in the Maastrichtian (R. Cestari and G. Sirna, Italy).

A new, purely ecological, classification of rudists was proposed by E. Gili (Spain) and P.W. Skelton (U.K.). Three main groups were distinguished: "elevators" characterized by upward growth of commissure; "clingers," which maximized actual contact with the substratum; and "recumbents," which maximize their effective shell diameters. This division is considered important for comparing forms from the same paleoecological group and not from various groups, as has been the case with all taxonomic schemes.

In the Gossau Formation of the eastern Alps, hippuritids built a hydrodynamically controlled colonial association of bioherms, biostromes and patch-reefs. They are distinguished in the form and relation by their cylindrical shells (R. Höfling, F.R.G.)

Based on a detailed study of morphologic and anatomic modifications in shells belonging to the Requeniidae family, J.-R. Masse (France) inferred that their phylogenetic development followed the pattern: from *Matheronia* - *Requenia* - *Toucasia* - *Pseudotoucasia* to *Lovetchenia*. A multitude of new specimens were described of various species of the interesting *Pironaea* genus from southern Valencia, Spain, similar to those found much earlier by Milovanović in eastern Serbia (J.M. Pons and E. Vicens, Spain). Based on a minute study of the outer shell layer of the left valve of the Hippuritidae, Pons and Vicens propose to use the whole canal and pore system for new taxonomic purposes instead of the pore form alone.

The importance for taxonomy was discussed of the ligamental ridge in rudist shells. Because the ligament in rudists suggests a devitalization of function, the ridge should not have a great taxonomic significance. A detailed study of *Pseudopolyconites*, however, by M. Sladić-Trifunović (Yugoslavia) revealed certain regularity in its ontogenetic development that could be used in identification of individual species.

During the Conference a proposal of the Organizing Committee was accepted for a Working Group on Rudists within the International Paleontological Association. M. Sladić-Trifunović, who chaired the Conference, was appointed Chairman of the Working Group, and A. Laviano (Italy) as Secretary, and it was decided to hold the next Conference in Rome in late 1990.

Aleksandar Grubić
Faculty of Geology and Mining
Belgrade University
Yugoslavia

Zuffardays '88

Cagliari, Sardinia, 10 - 15 October 1988

To honour Professor Piero Zuffardi, and to celebrate his 70th birthday, with some delay, a symposium and four field excursions were held in Sardinia, on problems of metallogeny in the western Mediterranean. Prof. Zuffardi was a founding member and former President of the Society for Geology Applied to Ore Deposits (SGA), one of IUGS' more active affiliates, especially in Europe. A former head of the Institute for Mineral Deposits of the University of Cagliari, Sardinia, he later became Professor at the University of Milan where he continued until recently his study of mineral deposits. Zuffardi has long been an authoritative proponent of the syngenetic interpretation of the Pb-Zn deposits of Sardinia. He early recognized the plate tectonic relationship of the Sardinian deposits with the carbonate-hosted Pb-Zn of southern France.

One highlight of the excursions was a visit to the Iglesiente deposit in the SW of the island. Formerly regarded as being derived from Hercynian granite, this is now recognized as a syndimentary, repeatedly remobilized platform deposit of Cambrian age. One topic of general interest was the origin of metals of these carbonate-hosted deposits. V. Köppel (Switzerland) explained these as the result of lateral secretion, deduced by a comparison of isotopes in the ores to those in the rocks of the basement, while H.J. Schneider (F.R.G.) postulated relations between ore and formation water in adjacent rocks.

Figure 1: Professor Piero Zuffardi.

E. Pani (Italy) and R. Valera (Italy) described the economically important fluorite deposits that occur in Ordovician rocks near the Cambrian contact partly as veins and partly as karst fillings. Formed during Late Paleozoic time, they have been mobilized by the Hercynian granites. Valera establishes a group of "intermediate" deposits, formed by endogenic and exogenic processes.

Conference Reports

Another excursion visited the large stratiform bauxite deposits of Olmedo in northwestern Sardinia. They are of the same age (Urgonian to Cenomanian) as the deposits in Provence (France) and equivalent from the point of view of plate tectonics. However, the former are far more regular in form and more homogenous in composition; they are para-autochthonous.

A group of veins with siderite, chalcopyrite and tetrahedrite in the northern Atlas mountains of Algeria was described by V. Glacon (France), as having the same mineral paragenesis as found in the eastern Alps and in the western Carpathians. Though rather enigmatic in origin, these deposits formed in an environment consisting of black shales and mafic rocks. P. Omenetto (Italy) presented a stimulating comparison of the deposits on the carbonate platforms of Gorno and Raibl (southern Alps), Thasos (northern Greece) and Morocco. These were brecciated and impregnated with metal-bearing solutions, partly bituminous.

One of the earliest and most faithful members of SGA, P. Evrard (Belgium) demonstrated the diversity of metallogenic problems by several examples. W.E. Petrascheck (Austria)

pointed to several still unsolved questions of Mediterranean metallogeny: the metal succession in the rifts, the ore deposits in ophiolites, relations between metamorphism and metallogeny, and the question of the original position of the Mediterranean micro-continents. P. Routhier (France) reviewed in witty fashion the evolution of metallogenic theories in Europe, in which Prof. Zuffardi played an important role.

In his concluding speech, Prof. Zuffardi emphasized that there was still a great deal of work to be done to understand properly the metallogeny of Sardinia. But participants left Cagliari most impressed by the high standards of the studies in this area, as illustrated by the excellent geological map of Sardinia. The meeting was ably organized by R. Valeva, the present Head of the Cagliari Institute for Mineral Deposits.

A. Pärtan and W.E. Petrascheck
Commission for Earth Sciences
Austrian Academy of Sciences
Dr. Ignaz Seipel Platz 2
A-1010 Vienna, Austria

Late Glacial and Holocene Environments, Vistula Basin

Poland, 15 - 21 June 1988

Although IGCP Project 158 (Palaeohydrology of the Temperate Zone During the Last 15000 Years) formally ended in 1987, some of its final results were presented at an important symposium organized jointly with INQUA's Eurosiberian Subcommission on the Holocene organized last year by the Polish Academy of Sciences, the National Committee for Quaternary Research and the W. Szafer Institute of Botany and other institutes. The 60 participants from 18 countries were reminded that Project 158 had made great efforts to correlate fluvial, lake and mire events with climatic changes and the impact of human activities. Important changes in fluvial regimes from the full glacial times to the Holocene were established, as were rhythmic phases of higher and lower fluvial activity with a recurrence interval of about 2000 years. The role of other factors, such as ice retreat, glacial rebound, sea-level rise, and anthropogenic activity was also recognized. New approaches had been developed in the reconstruction of hydrological regimes using palaeochannel morphometry and sedimentological parameters.

Data had been collected from over 100 localities for lakes and mires. The studies involved included pollen, diatoms, and the physical and chemical properties of sediments, as well as radiocarbon dates (at least one for each 2000 yr interval), ^{18}O and ^{13}C analyses. Statistical methods, helpful in pollen zonation and in the reconstruction of lake level changes available for studying lake and mire environments had been introduced, and an excellent manual presenting the full range of unified methods was published (*Handbook on Holocene Palaeoecology and Palaeohydrology*, edited by B. Berglund, John Wiley, 1986).

Another result of Project 158 was to foster closer cooperation between European and North American scientists, and such plans were also discussed. The participants generally approved a proposal presented for a new IGCP project on Global Continental Palaeohydrology. The objective of that project would be to establish the nature and causes of hydrological changes on various scales during the last 20,000

years, and to provide estimates of future hydrological conditions by studying past and predicting future water balance estimates.

Figure 1: Laminated sediments from an early Holocene horizon in Gościąg Lake, Poland. Height about 5 cms.

Future activities in palaeoecology and palaeoclimatology in relation to the Global Change Program were also reviewed. The importance of research on key localities was exemplified by preliminary results from the pilot reference site at Lake Gościąg where detailed investigations had already begun on the more than 12,000 annual layers rich in organic remains (Fig. 1). This may serve as a European standard locality for geochronology, as well as for the reconstruction of changes in climate and vegetation.

The symposium field trips to various lake-mire, fluvial and slope deposits, calcareous tufa and archaeological sites in the Carpathians, and in central and northern Poland demonstrated the high level of interdisciplinary paleogeographic studies in Poland and the existence there of a large group of enthusiasts.

- after report by L. Starkel
Polish Academy of Sciences
ul. św. Jana 22, 31-018 Kraków
Poland

Tectonostratigraphic Terranes in China

Nanjing, China, 23 September - 7 October 1988

China has long been recognized as a cluster of large polygonal or lozenge-shaped areas of relatively simple geologic structure separated by intensely folded and faulted boundary zones. In the period before widespread application of the theory of plate tectonics, the large areas were variously referred to as platforms or cratons, and the boundary zones as fold belts or geosynclines. After plate tectonics, more intracontinental mobility was implied by terms such as "blocks" for the large areas and "aulocogens" for the boundaries.

This view of a still relatively immutable China was called sharply into question in 1981 by evidence from M.W. McElhinny and others (Nature, v. 293, p. 212-216) that the various blocks have widely different paleomagnetic declinations and inclinations (see Fig. 1). Over the past eight years, ever-increasing mobility has been ascribed to the continental terranes of China, as illustrated, for example, by the Preliminary Tectonostratigraphic Terrane Map of the Circum-Pacific Region, published by the Circum-Pacific Map Project (IUGS affiliate CPCEMR - Ed.).

To evaluate present knowledge about the terranes in China, and to discuss the latest developments in the principles of terrane analysis, 150 geoscientists from 10 countries recently attended the Fourth International Tectonostratigraphic Terrane Conference at Nanjing University, China. The meeting was organized and chaired by Shi Yangshen (China) and D. Howell (U.S.A.).

Paleomagnetism was once again a primary tool to gain insight into the wanderings of tectonostratigraphic terranes. Lin Jinlu (China) stated that the Sinokorea (North China) and Mongolia continental terranes (Fig. 1) came

together during the Carboniferous, as did the Tarim and Kazakhstan terranes, but neither pair at that time had yet joined with Siberia. Sinokorea and Mongolia converged with Siberia during the Jurassic to Triassic, and Kazakhstan and Tarim continued their relative motion with respect to Siberia until the Cenozoic. The Yangze (South China) continental terrane joined with Sinokorea during the interval Jurassic to Triassic. Lin also stated that the paleomagnetic data restrict the extent of an eastward extrusion of south-east Asia following the collision with India, as proposed in 1982 by Tapponnier (Geology, v. 10, p. 611-616). Indochina rotated 10° clockwise with respect to South China, and South China moved 1000 km eastward with respect to Indochina. This movement is inferred to be the sum of initial left-lateral displacement on the Red River fault during the rotation, followed by greater right-lateral displacement on the same fault during the extrusion of China.

Fang Dajun (China) emphasized that South Korea is part of the Yangze (South China) continental terrane. The North China terrane, including North Korea, moved consistently northward until the Triassic, then rotated counterclockwise. The rotation continued between North and South China during the Jurassic, but by the Cretaceous they were fixed firmly together. Fang believes that North China accreted to northern Eurasia during the Early Cretaceous. A somewhat contrasting view was presented by Yuan Kuirong (China), who inferred that the Tarim and Sinokorea terranes had been joined during the Permian and then rotated 60° counterclockwise to their present easterly trend. The Yangze and Indochina terranes also were initially agglomerated together and have rotated counterclockwise an extraordinary 180° from an original inverted northwest trend to their present northeast position.

D. Howell likened the sutures of China to the boundaries between domains of contrasting age within the Canadian Shield, a kind of horizontal stratigraphy. Accretion at the former sites of the Panthalassa and Tethys oceans is marked by 200 Ma belts, the "graveyards" of the former oceans.

D. Jones (U.S.A.) and C. Sengör (Turkey) engaged in a lively debate on proper scientific method as applied to terrane analysis. Jones emphasized the importance of tectonic-assemblage diagrams and the delineation of multiple suspect terranes, especially during the early stages of research. T. Wiley (U.S.A.) provided an example of Jones' viewpoint by describing two stratigraphically similar terranes separated by 150 km on opposite sides of the Tintina strike-slip fault in Alaska. Previous investigators had considered them parts of the same terrane, dispersed by the fault, but further study showed that they had dissimilar volcanic clasts, oppositely verging faults, differences in organic thermal maturity, and dissimilar adjacent terranes, suggesting that

Figure 1: China at present and during the Permian, showing the rotation and assembly of its constituent terranes.

Conference Reports

Figure 2: The main strand of the Tanlu fault zone, near the town of Juxian, Shandong Province. Subhorizontal slickensides and right-lateral drag are clearly visible at this exposure. This fault marks the boundary between the West and East Shandong terranes. Photo by T. Wiley.

the stratigraphic similarity is only apparent. But Sengör stated that the conservative practice of seeking "facts," and separately naming every suspect subterrane, is a long step backward for tectonic study. Preconceived genetic concepts - hypotheses - even when wrong, are essential to progress in research. Hypotheses are nets, and if you don't cast your nets, you will catch no fish! Theories - templates - are essential to geologic field work.

According to P. Coney (U.S.A.), distant collisions can cause deep-seated deformation and wrenching at the center of a continent that may simulate terrane accretion there. He cited as examples the ancestral Rocky Mountains of North America and the Amadeus-MacDonnell belt in Australia, both perhaps caused by the brief encounter of the supercontinents of Gondwana and Laurasia during the Carboniferous. Likewise, E. Scheibner (Australia) emphasized that continent-continent collision is essential for the development of folded mountains. Aeromagnetic surveys can detect tectonic rotation, according to A. Rapolla (Italy) by a turning of a magnetic source's positive and negative anomalies, and can detect latitudinal displacement by a mismatch with predicted positive and negative sizes.

Lu Huaifu (China) outlined the complicated development of the syntaxis that marks the west edge of the Yangze terrane in central China, near 30°N, 100°E. A key element is the Bayankela terrane, a fragment of oceanic crust trapped between the Yangze terrane to the east and terranes from Gondwana to the southwest. The arrival during the Triassic

of the nearest of these, the Zhongzan terrane, sheared fragments from the Yangze terrane along the Longmenshan thrust and imbricated them over the Yangze. During the Permian, before the Yangze collision, the Zhongzan was itself swept up by the Qiangtang terrane to its southwest, and the two then moved together toward the Yangze. In Jurassic time, the North Tibet terrane, still farther to the southwest, collided with the assemblage, and this, together with the subsequent collision of India, has caused nearly continuous disruption and shearing of the syntaxis ever since.

Seismic profiles described by Weng Shijie (China) show that the deep Qinling suture between the Sinokorea and Yangze terranes dips to the north, whereas at the surface the dip is opposite, and the Yangze overthrusts the Sinokorea terrane. The presence of radiolarian chert and *mélange*, together with other evidence from the Nanhada terrane of north-eastern China, suggested to S. Mizutani (Japan) that the Nanhada terrane is a superterrane with Sikhote-Alin of the Soviet Far East, and with central Honshu, Japan.

As part of a series of papers on the important Tanlu fault zone of eastern China (Figure 2), S. Kumarapeli (Canada), noted that extension along the zone has created grabens containing Cretaceous sedimentary and volcanic rocks up to 9 km thick. Shi Yangshen (China) reported that west of the Tanlu fault zone, the West Shandong terrane contains a fairly continuous section from the Precambrian through the Mesozoic, whereas east of the zone the East Shandong terrane for the most part lacks Paleozoic rocks. The collision between the West and East Shandong terranes seems to have been of pre-Late Jurassic age and followed, during the Late Jurassic and Cretaceous, by transtension. According to Zhu Guang (China) the latter resulted from northerly motion of offshore plates in the Pacific that caused about 1000 km of left-lateral displacement on the Tanlu fault and 3000-4000 km on the Median Tectonic Line of Japan. L. Jolivet (France) agreed with the Mesozoic left-lateral transtension along the Tanlu zone but stated that the motion has now reversed to right-lateral, related to the Oligocene-Miocene opening of the Japan Sea.

In the view of G.W. Moore (U.S.A.), the early Tertiary collision of India with Eurasia, which shoved Eurasia northward, at first caused right-lateral tension at the continental margin, so as to open the Sea of Japan and the South China Sea. Then, during the late Tertiary, the right-lateral slip stepped westward to the Tanlu fault zone. The Pacific plate now moves toward China at 100 km per million years, meaning that, in the 43 Ma since the change in plate motion recorded by the Emperor bend in the Hawaii hotspot track, terranes from Gondwana could have moved by as much as 4300 km toward Asia. In this connection, S. Supriatna (Indonesia) stated that South Borneo (derived from Gondwana) and North Borneo (derived from China and Indochina) converged together along a south-dipping subduction zone during the Cretaceous and finally collided during the early Tertiary. The suture is marked by ultramafic rocks, gabbro, spilite, radiolarian chert and flysch. A post-collision extensional phase continues to the present.

The proceedings from the Nanjing conference will be published by the Circum-Pacific Council for Energy and Mineral Resources, under the editorship of T. Wiley and D. Howell of the U.S. Geological Survey. Plans are now being made for the fifth conference in 1991, probably to be held in South America, the only quadrant of the Circum-Pacific Region that has not yet hosted a conference on the rapidly expanding study of tectonostratigraphic terranes.

G.W. Moore
Department of Geology
Oregon State University
Corvallis, Oregon 97331
U.S.A.

Stratabound Copper in China

Shanxi, China, 12 - 24 October 1988

China is concerned about its copper supplies and would like to increase domestic reserves and production. To review a wide spectrum of stratabound Cu deposits, their metamorphosed equivalents, and exploration techniques, the sixth in a series of workshops sponsored under the IUGS/Unesco Deposit Modeling Program was hosted recently by the Shanxi Bureau of Geology and Mineral Resources, the Zhongtiaoshan Nonferrous Metal Mining Company, and the Chinese Ministry of Geology and Mineral Resources. Lectures by workshop leaders and Chinese contributors were given in Taiyuan and Zhongtiaoshan, and important Lower Proterozoic Cu deposits and host rocks were examined in the field in the Zhongtiaoshan district. Twenty-six participants, including exploration geologists from different provinces and autonomous regions and 16 geologists from Shanxi Province took part.

For this workshop, stratabound Cu deposits were grouped in the following types: sediment-hosted stratiform deposits, volcanogenic massive sulphides, volcanic redbeds (e.g. "Keweenaw-type"), other stratabound deposits (or those that do not fit readily into other classes), and sedimentary exhalative Pb and Zn (Cu, Ag, barite), a few of which contain significant Cu (e.g. Tammelsberg). The general characteristics, geological settings, probable genesis, and economic importance of the various types were discussed, and important deposits and districts in China and around the world were described. Models and exploration guides were presented for the various types of stratabound deposits, and a notebook, including a field guide for the Zhongtiaoshan district and selected bibliographies for various types of stratabound Cu deposits, and a separate volume of reprints were prepared.

Because many important stratabound Cu deposits, specifically those in the Zhongtiaoshan district, are strongly deformed and metamorphosed, experts on structural geology were in attendance. A knowledge of deformation and metamorphism is essential for understanding original features and settings of the deposits, for developing meaningful deposit models, and for evaluating redistribution and possible concentrations of metals during deformation and metamorphism.

Figure 1: Workshop participants, Shanxi Bureau of Geology and Mineral Resources, Taiyuan.

Figure 2: Workshop participants in the field near Zhongtiaoshan. From left to right: Lui Xuequei, Ren Gengshou, Yu Zhijie, and J. Slack.

The workshop also included group efforts to develop alternative working models for the highly deformed and metamorphosed Henglingguan and Hubi-type stratabound Cu deposits in the Zhongtiaoshan district. This exercise was useful for focusing on their principal characteristics, features of the areal geology that might be useful in exploration, and for identifying areas for further study.

Co-organizers of the workshop were Sun Renyi, Huang Chongke and Chenu Ping (China), and R. Kirkham (Canada), who was also workshop coordinator and leader. Other leaders were G. Borg (F.R.G.), K. Maiden (Australia), J. Slack (U.S.A.), and C. van Staal (Canada). Wan Liangguo of the Ministry of Geology and Mineral Resources, Beijing, helped with many of the arrangements.

R.V. Kirkham
Geological Survey of Canada
610 Booth Street
Ottawa, Ontario
Canada K1A 0E8