Classic Paper in the History of Geology

Charles Lapworth's "The Moffat Series", 1878

The investigation of the geology of the Southern Uplands, which form the border region between Scotland and England, has played an important role in the history of geological research, from the time of Hutton, through the nineteenth century, and to the present. The 1878 paper, the 'Moffat Series' by Charles Lapworth (see Figure 1) was a major part in the development of the understanding of the geology of the area. Dealing with a region of complex deformation, he worked out the biostratigraphy and structure, using a hitherto largely neglected fossil group, the graptolites. Twenty-one years after its publication, the history of research of this area included in the Memoir published by the Geological Survey in 1899, stated that: '[t]he publication of this [Moffat] paper marks an epoch in the history of the Silurian Geology of the South of Scotland. It remains the greatest and most original contribution to the study of the life-sequence and structural relations of these highly convoluted rocks'.

Figure 1 Charles Lapworth. Photograph from family archives, reproduced by kind permission of the late Patricia Lapworth.

Background

Charles Lapworth (1842–1920) was born at Buckden, near Faringdon, Berkshire, England. He was educated at the local village school and later went to Culham College, near Oxford, to train as a teacher. His first post was at St Peter's Episcopalian School at Galashiels, in the Scottish Borders, where he and his friends took up the hobby of natural history, especially the study of the local geology. Lapworth became a gifted amateur geologist and acknowledged expert on graptolites, discovering many new species. Between 1870 and 1878 he published a series of papers on graptolites and on the geology of the Southern Uplands. Initially, he worked according to the established ideas on the succession as stated by earlier workers, especially Sir Roderick Murchison, whose Geological Survey officers investigated the area between 1863 and 1873. But Lapworth gradually came to the conclusion that their ideas were untenable, and his work in the area developed on more individual lines, culminating in the Moffat paper. This paper stimulated further research by the Survey and others on Southern Uplands geology. It also enabled Lapworth (1879) to reclassify the Lower Palaeozoic in Britain into the tripartite system of Cambrian, Ordovician, and Silurian. Shortly after the publication of the Moffat paper, Lapworth, until then a schoolmaster, was appointed the first Professor of Geology at Birmingham University.

Lapworth's next piece of research was an application of the detailed stratigraphic procedures he had developed in southern Scotland to the complex region of Scotland's Northwest Highlands (though this time with less dependence on fossils). Here he established the nature of great lateral movements of the crust and conse-

quent repetition of beds. By working in the same meticulous way as he had done in the south, and applying what he had read of discoveries in the Alps, he worked out the succession and structures, and again stimulated the Survey and others to re-map the area and revise their thinking.

The problems in the geology of the Southern Uplands

According to William Watts, writing an appreciation of Charles Lapworth in 1921, the problems in the understanding of the Lower Palaeozoic of the Southern Uplands was similar to those of the Lower Palaeozoic elsewhere. Trilobites and brachiopods had been used for correlation, but had led to contradictions and inconsistencies. There was little or no confidence in graptolites as a stratigraphic tool. The main body of the Southern Uplands were considered to be virtually barren and to have a total thickness of greywackes, with shale bands, of 26,000 feet. It was thought that the graptolites collected from one shale band were comparable to those found in all the others, i.e. the fauna from the lowest band was thought similar to that of the highest. It was assumed that the whole series of greywackes and shales must belong to the same epoch, the Llandeilo, because the few recognizable fossils were dated as that. Because the graptolites were supposedly all found from the same epoch, they were thought to be useless as time indices. The supposition that 26,000 of rock had been deposited in the Llandeilo alone suggested either that deposition had been very rapid or that graptolites had not evolved during this time. In addition, the physical conditions in the Southern Uplands posed further problems. As Lapworth wrote:

[244] [A]ll the rocks of the district . . . [245] dip uniformly to the N.N.W. at high angles. Faults, folds, and inversions are occasionally visible among the greywackes; but in the dark shales they are astonishingly abundant. In every known locality where the Graptolitic beds are exposed, the majority are in this contorted and dislocated condition, and the attempt to ascertain their interrelationship by lithological and stratigraphical evidence has soon to be abandoned as hopeless. Their separation by zoological characters appears to be quite as desperate; for in many localities every trace of their former prolific fauna has been obliterated; in others only one or two fragmentary forms are obtainable, and these are limited to a few inches of thickness of the less altered zones. Even in those bands where the fossils are numerous and well preserved, the neighbouring exposures have frequently not a single fossil in common.

Lapworth's initial mapping

As mentioned, Lapworth's intellectual legacy came from Murchison, who had postulated a relatively simple set of parallel southwest-northeast folds, seemingly open and gently dipping, but showing considerable faulting. Any graptolites that were found were discounted for dating and correlation, and generally the fossil content in these shales and sandstones was thought to be patchy and poor. Dating was done on a macro-scale, using lithology as a basis for correlation with Welsh (rather than Lake District or Scandinavian) rocks.

Lapworth reported that Murchison had thought that the structure consisted of

[246] great folds, the upper arches of which had been denuded. At the same time he emphatically assigned to the whole of the rocks of the district a geological position inferior to that of the Bala Limestone of North Wales.

While acknowledging that this method was adequate for working out a sequence among slightly disturbed rocks, Lapworth stated that it failed where there were structural complexities. Lithostratigraphy alone was not equal to the task of unravelling the structure of the Southern Uplands.

Archibald Geikie, one of the Survey officers, interpreted the structure as being that of a grand anticline running from Dumfries in the west to St. Abb's Head in the east, with the youngest rocks (Caradoc) in a central syncline. However, he found it difficult to determine the succession: 'at the best the decision in most cases rested only on a combination of probabilities' (Geikie 1863, p. 8). Graptolites, thought to be Caradoc or younger, were sometimes found mixed with Llandeilo forms, but this problem was brushed aside using Joachim Barrande's (1799–1883) theory of 'colonies', which suggested that on occasions 'precursorial' forms might be found in strata, out of order with the usual stratigraphic sequence. The Survey published an account of the geology of the area immediately west of Moffat (Sheet 15, Leadhills) in 1871. It had been mapped by Robert Logan Jack (1845–1921) as a syncline of black graptolitic shales, dated as Upper Llandeilo. Nearby grits were dated as Caradoc and were thought similar to ones found elsewhere above the Llandeilo. Two years later (1873), Geikie, working in adjacent areas, found that the Caradoc did not always lie above the same part of the supposed Llandeilo and he proposed an unconformity with

overlap to explain this, not re-examining the earlier work of his colleagues in this highly disturbed area.

At this time, during his early work, which was done close to Galashiels, working with the accepted succession, Lapworth began to find many graptolites. Some were hitherto unknown in the area, while others were completely new species. These came from the black shales that were being exposed by the many new civil engineering works being undertaken in the area. It is possible, indeed probable, that the finding of graptolites in these new, more extensive exposures was the trigger and stimulus for developing his new methodology and there is evidence in his 1872 paper, '*Graptolites of the Gala Group*', that Lapworth was already using graptolites as a stratigraphic tool. He rapidly expanded his fieldwork area from Galashiels to the neighbouring countryside, eventually covering all the part of the Southern Uplands known as the Scottish Borders. The only method Lapworth found trustworthy was the use of palaeontological zones allied to lithology. He found historical justification for what he now termed his zonal approach, and he referred to the work of William Smith, James Hall, Albert Oppel, and Alcide d'Orbigny and their earlier, if cruder, zonal methodology. Other geologists from Britain, Scandinavia, North America, and France had also come to recognize the value of the zonal approach.

By the mid-1870s, Lapworth had become an acknowledged expert on graptolites and corresponded extensively with other geologists who worked in this field. He collaborated with (among others) Henry Alleyne Nicholson of St Andrews and Aberdeen Universities, and with William Swanston of Belfast and John Hopkinson from Hertford, an authority on Welsh graptolites. Lapworth's correspondence with Gustav Linnarsson of the Swedish Geological Survey was particularly important as both men were working on rocks of similar age and type. From the limited correspondence that remains,

Figure 2. Comparative sections of the Southern Uplands, from a letter from Charles Lapworth to Thomas McKenny Hughes, 1877 (coloured in the original). Reproduces by permission of Dr Paul Smith, Lapworth Museum, Birmingham University.

each seems to have told the other about the work he was doing and the results being obtained. Sequences worked out in Britain were tested in Sweden and vice versa, and the zonal work in the two countries proceeded simultaneously. The case for graptolite zonation grew stronger. Interestingly, though, Watts recorded in 1921, on the authority of the surveyor Benjamin Peach, that Lapworth first collected from Dobb's Linn non-zonally, until one day he collected successfully, i.e. with coherent results, in accordance with zonal ideas gathered from the writings of James Hall and Linnarsson.

Lapworth eventually explored the geology of the far west of his region, the valley of Moffat Water, which was known for possessing some of the best exposures of Lower Silurian rocks. This area was only 25 × 8 miles but it contained the most complete, least disturbed and best exposed sections in southern Scotland. One such was at a locality called Dobb's Linn (spelling at the time of Lapworth). This is a deep, branching valley system with large cliff faces of black shales and capped by pale sandstones. Lapworth's mapping and collecting of graptolites revealed a clear sequence of species (Figure 4).

By 1876, Lapworth had worked out a succession and structure for the area, and had published his findings as part of the 'Catalogue of the Western Scottish Fossils' exhibited at the British Association meeting in Glasgow in 1871 (see Watts 1939, p. 283). By 1877, he had completed his investigations of the Moffat area and communicated a brief summary privately to the former Survey officer, Thomas McKenny Hughes (1832–1917), recently appointed to the Cambridge professorship, who had earlier opposed some of Murchison's ideas. Lapworth's section (which was extended southwards to the English Lake District) was compared with that of Geikie, which showed a pronounced unconformity within the rocks of the Southern Uplands (see Figure 2). McKenny Hughes was one of a number of geologists who supported Lapworth's revised interpretation. Lapworth had, by this time, mapped much of Southern Scotland and established the main line of demarcation in the Silurian rocks there, both palaeontological and physical, at the base of the Birkhill Shales, above the Hartfell Shales and Glenkiln Shales. He took only a few months to write up his famous 'Moffat Series' paper, though he later told Watts that he rewrote it half a dozen times to achieve brevity, clarity, and cogency, reducing it to one third of its original size. It was read before The Geological Society of London on 21 November, 1877.

Fieldwork techniques

A remarkable feature of Lapworth's fieldwork was the scale at which he mapped. Dealing with shale bands in great detail, he required base maps at a very large scale. Even six inches to the mile (1:63,360) was insufficient. He had to enlarge these further, drawing his own for the individual key exposures such as Dobb's Linn. Most were drawn at 200 paces to 1 inch. Thus he could record the graptolites he found in the almost paper-thin shales with great accuracy.

There has been a tradition in the Lapworth family that he possessed a 'geological waistcoat', made by his wife Janet. This had multiple pockets down both front panels, in which he placed the carefully collected specimens in the order in which he extracted them from Dobb's Linn, first up one side of the valley and then down the other. The specimens were later decanted, placed in the collecting sequence, and through this painstaking and rather mundane method, allied to an unparalleled knowledge of the fossil group, the stratigraphy and the structure of the rocks was revealed.

In a lecture given in 1879, probably to The Geological Society of Glasgow, Lapworth also stated that he used a set of drawers with specimens placed in the order of their collection. Repetitions or reversals in order indicated folds, overturned beds, and faults. In the discussion that followed the reading of his paper, Lapworth was scathing about any collection of graptolites from the talus at the foot of the cliffs and not found in situ, implying that this practice was not uncommon.

The construction of the Moffat paper

The Moffat paper was divided into four parts:

1. An introduction, stating the geological characteristics of the south of Scotland.
2. A review of previous work done in the Moffat district.
3. A lengthy account of Lapworth's own work on the physical relations of the Moffat Series, with detailed descriptions of the important exposures, followed by an analysis of their palaeontology and stratigraphy.
4. Conclusions.

The physical relations of the area, the geological characteristics

The physical relations of the rocks found in this area were dealt with by describing the main exposures of Dobb's Linn. The reasons Lapworth gave for starting with this area were geologically sound:

[247] The only section of the Moffat Series which allows us to determine with certainty the sequence and palaeontological characteristics of its component beds, and at the same times clearly exhibits the relationship of the group as a whole to the surrounding greywackes, occurs in the centre of the Moffat district.

One of the tributaries of the Upper Moffat Water cuts steep-sided narrow gorge through the black and grey shales (see Figure 3).

[248] [T]he section of the shales and mudstones afforded by the northern cliff [North Cliff] of the lateral gorge . . . is visible from end to end, and, exception being made of a few local contortions, the strata exhibited appear to follow each other in natural and unbroken sequence. The coarse grits and flagstones of the [water]falls dip at 70° to 80° to the east, and plunge visibly beneath a thick series of grey shales with black bands.

Lapworth noted that the 'grey shales' were followed by two further shale bands, separated by mudstone, which passed beneath the greywackes at a low angle. He then traced the exposures of these strata up the south cliff, and across the face of the steep slope and the right-hand side of the longitudinal gorge, and found, crucially, that there occurred:

[248] a complete reversal of its original inclination, until, finally, in a magnificent cliff-section about 100 yards from our starting-place, all the zones we have recognized in the lateral gorge, together with others not there apparent, and inclusive also of the coarse grits and flagstones of the waterfall itself, are now arranged in the opposite order, and dip steadily at a gentle angle to the W.N.W.

This, Lapworth concluded, meant that one of the sections must be inverted. Comparison of the cross-sections, of both the north and south cliffs, printed with the map accompanying the paper (see Figure 3) show this clearly.

[248] [W]e are thus taught at the very outset of our inquiry how utterly futile is the endeavour to determine the original sequence of the deposits merely by attention to [249] their apparent order of superposition in any section, however perfect.

Lapworth supported this claim by dealing with numerous other exposures both to the north and south of the Moffat valley in the same way as at Dobb's Linn. Carefully-drawn sections accompanied these descriptions, illustrating his findings. He concluded this section of the paper with a summary of his observations and, most importantly, his conclusions regarding the physical relations of the rocks. By the simple, if dangerous, technique of following the rock bands laterally across cliffs and rivers he was able to determine the spatial arrangements and changes in the beds. He did not disregard

Figure 3. Map and Sections, from 'The Moffat Series', Plate XII.

lithological characteristics of colour, texture, etc., but these were not the main criteria for identification.

[248] *[T]he order of succession deduced from the evidence afforded by . . . one [section] is completely contradicted by that of the other. It is clear [therefore] that in one of these sections the strata must be inverted.*

Lapworth also demonstrated that a cursory glance at a large rock face could and did mislead. He examined the shale band in extreme detail, looking at [251] 'small flakes about the size and thickness of a finger-nail', and thereby worked out, in detail, a vertical section showing the subdivisions at Dobb's Linn. So he could recognize that:

[252] *[a]t this point [the base of the Lower Birkhill] a sudden and most extraordinary change takes place in the genera and species of the fossils of the Moffat Series. Scarcely a single form of those collected . . . in the strata already described is met with in any of the beds below this line, which are, however, like those above it, linked together by a large community of organic forms. Here, therefore, is the chief palaeontological break in the succession, and here, consequently, are we compelled to draw our main line of demarcation in any natural classification of the Moffat Series of this locality.*

Lapworth had come to the conclusion that, fundamental to his ability to identify the stratigraphic position of any exposure of black shale in the district:

[252] *each species and variety of Graptolite &c. has a definite range in the vertical succession of strata. None pass from the lowest to the highest zone; a few run up through the majority of the beds; several are common to two or three divisions; and the remainder are restricted to even narrower limits. Some (and these are the most valuable for our present purpose) are strictly confined to one or other of the mineralogical zones we have indicated. . . .*

Close analysis revealed divisions based not on lithological changes but on the vertical distribution of individual fossil species and on communities of species.

Not only did Lapworth successfully use zonal techniques for stratigraphic purposes, he was also able to ascertain structural features i.e. a series of reverse faults within a fan-shaped anticline that broke up the strata into a series of wedges, which he compared to the larger but analogous structures in the Alleghenies and the Alps.

[259] *The Moffat Series of this locality, is disposed in the form of a rude arch, which is broken by three longitudinal faults. In the centre of the Glen the plane of the main axis of the anticline is approximately perpendicular, and the beds are shown in their natural position. To the northward the axis dips to the eastward, and the strata upon its western side are all inverted. In the southern portion of the Glen, on the other hand, the axis dips to the westward, and it is the eastern beds which are overturned. . . .*

These three dislocations are all of the puzzling class known as inverted faults—the hade being towards the upthrow side of the break.

It remained for Lapworth to test his findings in highly dislocated strata and also elsewhere in the area. Thus, after this detailed account of his work in Dobb's Linn, he confirmed his findings at nearby Craigmichan Scaurs [scars or crags]. He wrote:

[265] *We have . . . gained a much fuller insight into the thickness and lithological characters of the Glenkiln Shales [the base of the succession in the area], which are seen to have an importance which would never have been suspected from the insignificant exposure of these beds at Dobb's Linn, and we are now in a position to commence the study of the numerous black-shale bands of the district.*

Lapworth then proceeded to describe the black shales found south of the Moffat valley, southwest of St. Mary's Loch, those in the Yarrow valley and in the valleys of Ettrick and Glenkiln. This was followed by similar descriptions of the area north of the Moffat valley, in the basin of the upper Annan river, which included French-

land Burn and Hartfell Spa, the basins of Meggat Water and Moffat Water itself.

The stratigraphical and palaeontological findings

In his next major section Lapworth began with a description of the lowest strata found, the Glenkiln, followed by an analysis of the Hartfell and then the Birkhill Shales. He divided each of these units on the basis of their graptolite assemblages. Key zonal fossils were identified, and the sequence was thus 'labelled'. His analysis was summarized in a table giving the vertical distribution of fossils, which emerged as the result of his meticulous fieldwork (see Figure 4).

[254] *[I]t is evident that the strata of the Moffat Series exhibited in the Main Cliff of Dobb's Linn are naturally arranged in two main divisions palaeontologically distinct. Each of these divisions, again, falls naturally into two well-defined mineralogical subdivisions, each of which is, in turn, composed of several subordinate bands or zones, individually distinguishable by peculiar lithological and palaeontological features.*

In general, the Southern Uplands, with their blanket of Pleistocene 'drift', have poorly exposed bedrock. But after close examination of the few places where the bedrock pierced the boulder-clay cover, Lapworth was able to make correlations between separate localities, and develop new theories regarding the succession and structure. Finally, he put all the new work into the context of the Silurian of Southern Scotland. In other words, he provided a new successional framework, finding, as compared with the results of the Survey officers, a drastically reduced thickness of rocks, which had anything but a simple structure, being folded and faulted, yielding a complex repetition of beds. By contrast, the Survey officers, depended on broad lithological and biostratigraphic generalizations and could not always be sure whether they were ascending or descending a succession. The days of generalized and superficial mapping of an area were now over so far as Britain was concerned, and the meticulous procedures now used by field geologists were shown by Lapworth to be essential for successful analysis of complex areas.

Conclusion

Finally Lapworth summarized his discoveries and in his last section put his detailed findings into a regional and international context:

- [300] 1. *All the black Graptolitic shales of the Moffat district are actually portions of one and the same continuous deposit.*
2. *They owe their repetition to a series of subparallel folds, the upper arches of which have been denuded.*
3. *They rise invariably from below the surrounding greywackes in anticlinal forms, the axes of which are usually inverted.*
4. *The deposit to which they belong is consequently the oldest rock-group in the Moffat district.*
5. *This fundamental rock-group, which is denominated by us the Moffat Series, is composed of a comparatively homogeneous assemblage of dark Graptolitic shales and pale barren mudstones.*
6. *Its collective thickness within the limits of the present district is about 300 feet, but its actual base is nowhere visible.*
7. *It falls naturally into three primary divisions, each of which is characterized by a special fauna, made up almost wholly of peculiar species.*
8. *Each of these primary divisions is again naturally*

Figure 4 Vertical section of the Moffat Series at Dobb's Linn, showing the zonal succession. From 'The Moffat Series', Figure 1, p. 250.

subdivided into several zones, individually distinguished by special mineralogical and palaeontological characteristics.

The small and sometimes boat-shaped outcrops of the Moffat Shales had obviously puzzled geologists in the past. But, said Lapworth:

[301] [t]hey now find their common explanation in the simple circumstance that the dark shales reach the surface of the country along the chief anticlinal lines, the width of the several exposures being dependent merely upon the varying elevation of the crown of the arch.

Also, Barrande's theory of colonies, which, if applied to the graptolite fauna, precluded any coherent succession based on these fossils, was rendered irrelevant.

[301] The extraordinary diversity apparent in the various groups of species yielded by the strata of the same band in the different exposures along its course, and the peculiar localization of some of its most distinctive fossil forms, are quite as easily explained by our discovery of the rigid restriction of the Graptolitic species to definite zones. Not only do the earliest zones of each band make their appearance as a rule only in the widest exposures, but its apparent fauna in any single locality is necessarily that of the collective faunas of the special zones, which the concurrent accidents of plication, metamorphism, and denudation have there left accessible to the investigator. Lastly, we have no longer any room for astonishment at the remarkable similarity in lithological characters, the great lateral extension, persistent north-west inclination, and apparently gigantic thickness of the interminable greywackes of the Moffat district. We are now satisfied that they form in reality a single group of beds of no great vertical dimensions, the same strata being repeated again and again in rapid and partially inverted undulations.

Subsequently, the Geological Survey re-mapped the Southern Uplands using Lapworth's zonal/palaeontological methodology. The result was the 1899 Memoir. Lapworth was asked to review this work for the *Geological Magazine*, which he did in an extensive paper. He stated that

zonal work was probably destined to effect in the history of geological research a revolution as great and an advance as rapid as those brought about by the use of the microscope in the history of biology. This volume does not mark the close of geological investigation in the Uplands. It marks rather the beginning.

Lapworth could have been writing the final comments on his own 1878 paper.

The Moffat paper had a mixed reception. According to Watts it 'attracted the interest and sympathy of the younger men, but the incredulity and even hostility of the holders of older views'. In the discussion that followed the reading of the paper at the Geological Society most of those who had heard it appeared to be impressed by Lapworth's findings and work. On the whole his explanation of the succession and structure of the Southern Uplands was welcomed and judged to be 'a base-line for the correlation of all the Silurian beds of the south of Scotland'. Henry Hicks, John Judd and McKenny Hughes were all supportive. Only Andrew Ramsey, Murchison's successor at the Survey, cited the work of Geikie and his assistants in mapping rocks above the Moffat Series, and suggested that the Scottish beds might represent a gap in the succession in Wales and Shropshire. In reply, Lapworth briefly reiterated the problems of dealing with the apparent order of superposition, and rejected the Survey's postulated unconformities as being without physical evidence.

The favourable reaction of the Geological community to the paper was clear in February the following year. The balance of the proceeds of the Murchison Geological Fund was awarded to Lapworth for his work on the Silurian rocks of the South of Scotland and the graptolites contained in them. Also in 1879, Lapworth put forward his proposal for the tri-partite division of the Lower Palaeozoic into Cambrian, Ordovician, and Silurian. Though this clarified mat-

ters greatly and provided a neat resolution of the controversy between Murchison and Adam Sedgwick about the placement of the Cambrian-Silurian boundary, it was not accepted on the Geological Survey's maps until Geikie's retirement in 1900. The tripartite division has now long been accepted. It rested on the remarkable empirical work of Lapworth in the Southern Uplands and the stratigraphic and structural understanding that this work made possible.

References and selected reading

- Aldridge, R. J., Siveter, David J., Siveter, Derek J., Lane P. D., Palmer, D. and Woodcock, N. H., 2000, *British Silurian Stratigraphy*: Peterborough, Joint Nature Conservation Committee, pp. 149–157.
- Barrande, J., 1881, *Défence des Colonies V. Apparition et réapparition en Angleterre et en Ecosse des espèces coloniales Silurienne de la Bohême*: Prague and Paris, J. Barrande.
- Fortey, R. A., 1993, Charles Lapworth and the biostratigraphic paradigm: *Journal of the Geological Society*, v. 150, pp. 209–218.
- Geikie, A., 1863, *Explanation of Sheet 34: The Geology of Eastern Berwickshire*: Memoir of the Geological Survey of Great Britain.
- Geikie, A., 1871, *Explanation of Sheet 15: Dumfriesshire (north part); Lanarkshire (south part); Ayrshire (southeast part)*: Memoir of the Geological Survey of Great Britain.
- Hamilton, B. M., 1984, The contribution of Gustaf Linnarsson to British stratigraphic geology: *Geologiska Föreläsningar*, v. 106, pp. 185–192.
- Horný, R., 1980, Joachim Barrande (1799–1883), life, work, collections: *Journal of the Society for the Bibliography of Natural History*, v. 4, pp. 365–368.
- Lapworth, C. and Wilson, J., 1871, On the Silurian rocks of the counties of Roxborough and Selkirk: *Geological Magazine*, decade 1, v. 8, pp. 456–464.
- Lapworth, C., 1872, On the graptolites of the Gala Group: Report of the British Association for the Advancement of Science, v. 41, p. 104.
- Lapworth, C., 1872, On the Silurian rocks of the south of Scotland: *Transactions of the Glasgow Geological Society*, v. 4, pp. 164–174.
- Lapworth, C., 1879, The Silurian age, MS Lecture notes: University of Birmingham Archives.
- Lapworth, C., 1899, Review of the Memoirs of the Geological Survey of the United Kingdom, 'The Silurian Rocks of Britain', Vol. 1 Scotland, 1899: *Geological Magazine*, decade 4, v. 6, pp. 472–479 and 510–520.
- Lapworth, C., 1870, On the Lower Silurian rocks of Galashiels: *Geological Magazine*, decade 1, v. 6, pp. 204–209, and 279–284.
- Lapworth, C., 1878, The Moffat Series: *Quarterly Journal of the Geological Society of London*, v. 34, pp. 240–346.
- Murchison, R. I. M., 1872, *Siluria: A History of the Oldest Rocks in the British Isles and Other Countries*, 5th edn: London, John Murray.
- Oldroyd, D. R., 1990, *The Highland Controversy: Constructing Geological Knowledge through Fieldwork in Nineteenth-Century Britain*: Chicago and London, University of Chicago Press.
- Peach, B. N. and Horne, J., 1899, *The Silurian Rocks of Britain*, Vol. 1, Scotland: Memoirs of the Geological Survey of the United Kingdom, Glasgow, H. M. S. O.
- Watts, W. W., 1939, The author of the Ordovician system; Charles Lapworth, M.Sc., L.L.D., F.R.S., F.G.S. *Proceedings of the Geologists' Association*, v. 50, pp. 235–286.

Beryl Hamilton

Kirk Road

New Galloway

Dumfries and Galloway DG7 3RS

UNITED KINGDOM